Sex-specific additive genetic variances and correlations for fitness in a song sparrow (*Melospiza melodia*) population subject to natural immigration and inbreeding

Matthew E. Wolak^{1,2}, Peter Arcese³, Lukas F. Keller^{4,5}, Pirmin Nietlisbach^{4,6}, and Jane M. Reid¹

¹School of Biological Sciences, University of Aberdeen, Aberdeen, Scotland

²Department of Biological Sciences, Auburn University, Auburn, AL, USA

³Department of Forest and Conservation Sciences, University of British Columbia, Vancouver, British Columbia, Canada

⁴Department of Evolutionary Biology and Environmental Studies, University of Zurich, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland

⁵Zoological Museum, University of Zurich, Karl-Schmid-Strasse 4, CH-8006 Zurich, Switzerland

⁶Department of Zoology, University of British Columbia, Vancouver, British Columbia, Canada

* Corresponding authors: matthew.wolak@auburn.edu and jane.reid@abdn.ac.uk

Running head: Additive genetic variance in fitness

AUTHOR CONTRIBUTIONS

MEW and JMR jointly conceived and designed the analyses and wrote the manuscript. MEW conducted the analyses, with contributions from JMR. PA ensured the long-term field data collection. PN and LFK conducted the genotyping. All authors contributed to editing the final draft.

ACKNOWLEDGMENTS

We thank the Tsawout and Tseycum First Nation bands for access to Mandarte and everyone who contributed to the long-term data collection. We thank the European Research Council for funding and the University of Aberdeen for generous access to the Maxwell High Performance Computing cluster. Pierre de Villemereuil, Michael B. Morrissey, and Jarrod D. Hadfield provided enlightening discussions during manuscript preparation. Joel McGlothlin and two anonymous reviewers provided further helpful comments.

DATA ARCHIVING

Data have been archived in the Dryad Digital Repository:

https://doi.org/10.5061/dryad.p7p1jb3 (Wolak et al. 2018).