

Archivos de Bronconeumologia

Changes in control status of COPD over time and their consequences: A prospective international, study.

--Borrador del manuscrito--

Número del manuscrito:	ARBR-D-20-00344R1
Tipo de artículo:	Original
Palabras clave:	EPOC; Control; agudizaciones; desenlaces
Autor correspondiente:	Marc Miravittles Hospital Vall d'Hebron Barcelona, SPAIN
Primer autor:	Marc Miravittles
Orden de autores:	Marc Miravittles Pawel Sliwinski Chin Kook Rhee Richard Costello Victoria Carter Jessica HY Tan Therese Sophie Lapperre Bernardino Alcazar Navarrete Caroline Gouder Cristina Esquinas Juan Luis García Rivero Anu Kemppinen Augustine Tee Miguel Román Rodríguez Juan José Soler-Cataluña David Price
Resumen:	<p>Introducción: El estado de control ha demostrado tener un valor predictivo a largo plazo para las agudizaciones, pero no existe información sobre su valor predictivo a corto plazo o de los cambios de control a lo largo del tiempo.</p> <p>Método: Estudio multicéntrico, internacional, prospectivo para observar el valor pronóstico a corto plazo (6 meses) del estado de control. Los pacientes se clasificaron en controlados/no controlados en la visita basal y en 3 visitas a intervalos de 6 meses utilizando unos criterios de control validados. La tasa de agudizaciones moderadas y graves se comparó entre las visitas clasificadas como controladas y no controladas y entre los pacientes persistentemente controlados, no controlados y aquellos que cambiaron el estatus de control durante el seguimiento.</p> <p>Resultados: Se analizaron 267 pacientes, entre ellos 80 (29,8%) estaban permanentemente controlados, 43 (16%) permanentemente no controlados y 144 (53,7%) cambiaron su estado de control durante el seguimiento. Durante los 6 meses posteriores a una visita en un paciente no controlado la Odds ratio (OR) de presentar una agudización moderada fue de 3,41 (intervalo de confianza (IC) del 95% 2,47 a 4,69) y una OR= 4,25 (IC95% 2,48 a 7,27) para la hospitalización comparado con una visita en un paciente controlado.</p> <p>Conclusiones: La evaluación del control en cada visita proporciona una información pronóstica importante sobre el riesgo de agudización en los próximos 6 meses. La falta de control es una señal de alarma que debe motivar la investigación y la acción para conseguir retornar al estado de control.</p>

Changes in control status of COPD over time and their consequences: A prospective international, study.

Authors: Marc Miravittles (1,2), Pawel Sliwinski (3), Chin Kook Rhee (4), Richard W Costello (5), Victoria Carter (6), Jessica HY Tan (7), Therese Sophie Lapperre (8,9), Bernardino Alcazar (2,10), Caroline Gouder (11), Cristina Esquinas (1,12), Juan Luis García-Rivero (13), Anu Kemppinen (6), Augustine Tee (14), Miguel Roman-Rodríguez (15), Juan José Soler-Cataluña (2,16), David B. Price (17,18), on Behalf of the Respiratory Effectiveness Group (REG).

Center: 1. Pneumology Department. Hospital Universitari Vall d'Hebron; Vall d'Hebron Institut de Recerca (VHIR), Vall d'Hebron Barcelona Hospital Campus, Barcelona, Spain. 2. CIBER de Enfermedades Respiratorias (CIBERES). Spain. 3. 2nd Department of Respiratory Medicine. Institute of Tuberculosis and Lung Diseases, Warsaw, Poland. 4. Division of Pulmonary, Allergy and Critical Care Medicine, Department of Internal Medicine, Seoul St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Seoul, Republic of Korea. 5. Department of Respiratory Medicine, Royal College of Surgeons. Dublin, Ireland. 6. Optimum Patient Care, Cambridge, UK. 7. Department of Respiratory and Critical Care Medicine, Singapore General Hospital, Singapore. 8. Duke-National University of Singapore Medical School, Singapore. 9. Department of Respiratory Medicine, Bispebjerg Hospital, Copenhagen, Denmark. 10. Respiratory Department, Hospital de Alta Resolución de Loja, Spain. 11. Department of Respiratory Medicine, Mater Dei Hospital, Malta. 12. Public Health, Mental, Maternal and Child Health Nursing Department. Faculty of Medicine and Health Sciences. University of Barcelona, Spain. 13. Department of Respiratory Medicine, Hospital Comarcal de

Laredo, Cantabria, Spain. 14. Respiratory and Critical Care Medicine, Changi General Hospital. Singapore 15. Primary Health-care Center Son Pisà. IB-Salut. Palma, Balears. Spain. 16. Pneumology Department, Hospital Arnau de Vilanova, Valencia, Spain. 17. Centre of Academic Primary Care, University of Aberdeen, UK. 18. Observational and Pragmatic Research Institute, Singapore.

Correspondence: Marc Miravittles

Pneumology Department. University Hospital Vall d'Hebron

P. Vall d'Hebron 119-129. 08035 Barcelona, Spain

Phone and Fax: +34 93 2746083

E-mail: marcm@separ.es

Authors' Contributions: *Conceptualization: MM, VC, AK, DBP. Data curation: MM, CE, AK. Data Acquisition: MM, PS, CKR, RWC, VC, JHYT, TSL, BA, CG, JLG-R, AT, MR-R. Formal analysis: MM, CE, AK. Funding acquisition: MM, DBP. Investigation: MM, PS, CKR, RWC, VC, JHYT, TSL, BA, CG, JLG-R, AT, MR-R, JJS-C, DBP. Project administration: MM, AK, DBP. Writing—original draft: MM. Writing—review and editing: PS, CKR, RWC, VC, JHYT, TSL, BA, CG, JLG-R, AT, MR-R, JJS-C, DBP*

ACKNOWLEDGMENTS

The study was designed and coordinated by the Respiratory Effectiveness Group (REG; www.effectivenessevaluation.org; Cambridge, UK) and delivered by Optimum Patient Care (OPC; www.optimumpatientcare.org). **Funding:** The study was funded by an unrestricted grant from Novartis AG.

The following investigators participated in the study: **Spain:** Marc Miravittles, Cristina Esquinas, Miriam Barrecheguren, Alexa Nuñez, Hospital Universitari Vall d'Hebron, Barcelona. Bernardino Alcazar, Hospital de Alta Resolución de Loja. Juan Luis García-Rivero, Karina Hueso, Hospital Comarcal de Laredo, Cantabria. Miguel Roman-Rodríguez, Primary Health-care Center Son Pisà. IB-Salut. Palma de Mallorca. **Poland:** Pawel Sliwinski, Katarzyna Iwan, Jacek Kolakowski, Institute of Tuberculosis and Lung Diseases, Warsaw. **Korea:** Chin Kook Rhee, Esther Ahn, St Mary's Hospital. Seoul. **Singapore:** Jessica Tan, Therese Laperre, Karen Tan Li Leng, Nicole Chia, Ong Thun How, SyifaBinte Shamsuddin, Sherine Lim Shu Gim, Yap Chwee Bee, Soh Rui Ya, Singapore General Hospital. Augustine Tee, Jun Jie Yan, Samuel Hong, William Tan, Changi General Hospital. **UK:** Victoria Carter, Latife Hardaker, Andrew McLaughlin, Optimum Patient Care, Cambridge. **Malta:** Caroline Gouder, Mater Dei Hospital. **Ireland:** Richard W Costello, Royal College of Surgeons. Dublin.

DECLARATIONS OF INTEREST

MM has received speaker fees from AstraZeneca, Boehringer Ingelheim, Chiesi, Cipla, Menarini, Rovi, Bial, Sandoz, Zambon, CSL Behring, Grifols and Novartis, consulting fees from AstraZeneca, Boehringer Ingelheim, Chiesi, GlaxoSmithKline, Bial, Gebro Pharma, Kamada, CSL Behring, Laboratorios Esteve, Ferrer, Mereo Biopharma, Verona Pharma, TEVA, pH Pharma, Novartis, Sanofi and Grifols and research grants from GlaxoSmithKline and Grifols. PS has received speaker fees from Boehringer Ingelheim, AstraZeneca, Chiesi, GlaxoSmithKline, Grifols, Novartis, Roche and Teva, and consulting fees from Boehringer Ingelheim, GlaxoSmithKline, Chiesi, Grifols, Novartis and Roche.

CK Rhee received consulting/lecture fees from MSD, AstraZeneca, Novartis, GSK, Takeda, Mundipharma, Sandoz, Boehringer-Ingelheim, and Teva-Handok.

RWC has board membership with GSK, Aerogen, Novartis, and Teva Pharmaceuticals; consultancy agreements with, Aerogen, GlaxoSmithKline, Novartis, Teva Pharmaceuticals, and Vitalograph as well as grants and unrestricted funding for investigator-initiated studies from Vitalograph, Aerogen and GlaxoSmithKline.

TL has received speaker and consulting fees from Boehringer Ingelheim, GlaxoSmithKline, AstraZeneca and Novartis.

BA reports grants and personal fees from GSK, grants, personal fees and non-financial support from Novartis AG, personal fees and non-financial support from Boehringer Ingelheim and Chiesi, grants, personal fees and non-financial support from Laboratorios Menarini, personal fees from Gebro, Astra- Zeneca, Laboratorios Rovi and Laboratorios Ferrer, outside the submitted work.

JJSC has received speaker fees from AstraZeneca, Boehringer Ingelheim, Chiesi, Esteve, Ferrer, GSK, Menarini, Novartis, and Pfizer, and consulting fees from AirLiquide, Boehringer Ingelheim, Chiesi, GSK, AstraZeneca, Ferrer and Novartis.

DBP has board membership with Amgen, AstraZeneca, Boehringer Ingelheim, Chiesi, Circassia, Mylan, Mundipharma, Novartis, Regeneron Pharmaceuticals, Sanofi Genzyme, Teva Pharmaceuticals, Thermofisher; consultancy agreements with Amgen, AstraZeneca, Boehringer Ingelheim, Chiesi, GlaxoSmithKline, Mylan, Mundipharma, Novartis, Pfizer, Teva Pharmaceuticals, Theravance; grants and unrestricted funding for investigator-initiated studies (conducted through Observational and Pragmatic Research Institute Pte Ltd) from AstraZeneca, Boehringer Ingelheim, Chiesi, Circassia, Mylan, Mundipharma, Novartis, Pfizer, Regeneron Pharmaceuticals, Respiratory Effectiveness Group, Sanofi Genzyme, Teva Pharmaceuticals, Theravance, UK National Health

Service; payment for lectures/speaking engagements from AstraZeneca, Boehringer Ingelheim, Chiesi, Cipla, GlaxoSmithKline, Kyorin, Mylan, Mundipharma, Novartis, Regeneron Pharmaceuticals, Sanofi Genzyme, Teva Pharmaceuticals; payment for the development of educational materials from Mundipharma, Novartis; payment for travel/accommodation/meeting expenses from AstraZeneca, Boehringer Ingelheim, Mundipharma, Mylan, Novartis, Thermofisher; funding for patient enrolment or completion of research from Novartis; stock/stock options from AKL Research and Development Ltd which produces phytopharmaceuticals; owns 74% of the social enterprise Optimum Patient Care Ltd (Australia and UK) and 74% of Observational and Pragmatic Research Institute Pte Ltd (Singapore); and is peer reviewer for grant committees of the Efficacy and Mechanism Evaluation programme, and Health Technology Assessment.

AT has participated as a member of the local COPD advisory board for Astra Zeneca, GlaxoSmithKline, Bayer, Takeda and Novartis and has received meeting/conference travel grants from Boehringer Ingelheim, Novartis, Astra Zeneca and GlaxoSmithKline.

JGR has received speaker fees from Novartis, GSK, Boehringer-Ingelheim, AstraZeneca, Chiesi, ALK, Teva, Menarini, Rovi, Esteve and Pfizer; and consulting fees from Novartis, GSK, Astra-Zeneca, Teva, Boehringer-Ingelheim, ALK and Esteve.

MRR has received speaker fees from AstraZeneca, Boehringer Ingelheim, Chiesi, Gebro Pharma, GlaxoSmithKline, Menarini, Mundipharma, Novartis, Pfizer and Teva, and consulting fees from AstraZeneca, Boehringer Ingelheim, GlaxoSmithKline and Novartis.

Word count: 2573

Abstract word count: 241

1
2
3
4
5
6
7 **Changes in control status of COPD over time and their consequences: A prospective**
8 **international, study.**
9

10
11 **ABSTRACT**
12

13
14 **Introduction:** Control status may be a useful tool to assess response to treatment at each clinical
15 visit in COPD. Control status has demonstrated to have long-term predictive value for
16 exacerbations, but there is no information about the short-term predictive value of the lack of
17 control and changes in control status over time.
18
19

20
21
22
23 **Method:** Prospective, international, multicenter study aimed at **validating describing** the short-
24 term (6 months) prognostic value of control status in patients with COPD. Patients with COPD
25 were classified as controlled/uncontrolled at baseline and at 3,6- month follow-up visits using
26 previously validated criteria of control. **Moderate and severe** exacerbation rates were compared
27 between controlled and uncontrolled visits and between patients persistently controlled,
28 uncontrolled and those changing control status over follow-up.
29
30

31
32
33 **Results:** A total of 267 patients were analyzed: 80 (29.8%) were persistently controlled, 43 (16%)
34 persistently uncontrolled and 144 (53.7%) changed control status during follow-up. Persistently
35 controlled patients were more frequently men, with lower (not increased) body mass index and
36 higher FEV1(%). During the 6 months following an uncontrolled patient visit the odds ratio (OR)
37 for presenting a moderate exacerbation was 3.41 (95% confidence interval (CI) 2.47 to 4.69) and
38 OR= 4.25 (95%CI 2.48 to 7.27) for hospitalization compared with a controlled patient visit.
39
40

41
42
43 **Conclusions:** Evaluation of control status at each clinical visit provides relevant prognostic
44 information about the risk of exacerbation in the next 6 months. Lack of control is a warning
45 signal that should prompt investigation and action in order to achieve control status.
46
47
48

49
50
51 **Keywords:** COPD; control; CAT; exacerbations; outcomes.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

INTRODUCTION

The most recent update of the Global Strategy for Obstructive Lung Disease (GOLD) recommends reassessing chronic obstructive pulmonary disease (COPD) patients after initial treatment based on symptoms and exacerbations (1); however, there is no clear guidance as to how or how often this reassessment should be conducted. The development and validation of a control tool that easily identifies patients who may require a step up/down in treatment and a closer follow-up has been successfully achieved in the management of asthma (2), but the concept of control has been elusive in COPD (3). The main reason for this is the identification of the concept of control with normality in fact, controlled asthma is associated with normal lung function and absence of symptoms (2). While it is obvious that these objectives are not attainable in COPD, it is also true that thanks to smoking cessation, vaccinations and appropriate pharmacological and non-pharmacological treatments many COPD patients may achieve a low level of symptoms and remain stable, without exacerbations and without significant worsening in lung function over prolonged periods of time (4). This status of low level of symptoms (or low impact) and stability is what has been defined as control in COPD (5,6).

A recent study demonstrated that the evaluation of control in COPD is more sensitive to clinical changes than changes in GOLD stage, in risk level or in phenotype (7). Moreover, a change in control status over 3 months is followed by significant changes in COPD assessment test (CAT) scores, thereby, making evaluation of COPD control a potential tool to reassess patients at each clinical visit (7).

However, the risk of poor outcomes (i.e. exacerbations of different severity) in the short term in association with the uncontrolled status or with the change in status from controlled to uncontrolled has not been previously reported. This is critical to understand the clinical value of the evaluation of control. If uncontrolled status or change to uncontrolled status is associated with

1
2
3
4 increased risk of exacerbations during the next months, this would justify investigating the cause
5
6 of the lack of control and an intervention to restore the control status and prevent undesirable
7
8 outcomes.
9

10
11 The current study is the first prospective, international, multicenter study designed with the
12
13 objective to ~~validate~~ ~~investigate~~ the concept of control of COPD. In a previous publication we
14
15 ~~validated~~ ~~described~~ control status as a marker of increased risk of poor outcomes during follow-
16
17 up (8). Here, we describe the changes in control status in patients over long-term follow-up, the
18
19 short-term risk associated with uncontrolled patient visits and the main characteristics of patients
20
21 that remained either controlled or uncontrolled during the study.
22
23
24
25
26
27

28 **METHOD**

29 **Design of the study**

30
31 This was a prospective international, multicenter study aimed at ~~validating~~ ~~developing~~ the concept
32
33 of control in COPD. The design of the study has been described in detail in previous publications
34
35 (8-10). Eligible patients were recruited to the study and underwent a screening visit (V-1)
36
37 involving full clinical assessment, including socioeconomic variables, evaluation of current
38
39 smoking status, current treatment, respiratory symptoms, presence of comorbidities, lung function
40
41 measured by spirometry and questionnaires. The BODEx (Body mass index (BMI), obstruction,
42
43 dyspnea and exacerbations) index was calculated (10). A baseline visit (V0) was scheduled after
44
45 3 months, in which the control status of the patients was assessed as indicated below. After
46
47 baseline, the patients were followed with 3 visits at 6-month intervals for a total of 18 months. At
48
49 each follow-up visit the control status was assessed as at baseline.
50
51
52
53
54
55
56

57
58 At each clinical visit, the CAT (11) and the modified Medical Research Council (mMRC) dyspnea
59
60 scale (12) were administered, comorbidities were assessed with the Charlson index (13) and
61
62
63
64
65

1
2
3
4 physical activity were quantified by self-declared minutes of walking per day (14).
5

6 A **moderate** COPD exacerbation was defined as an increase in respiratory symptoms that required
7 the use of systemic corticosteroids and/or a course of antibiotics; **when exacerbation required**
8 **hospital admission it was considered severe** (15).
9
10
11
12
13

14 The primary outcome of the study was the difference in rates of a composite endpoint during the
15 18-month follow-up between patients who were controlled versus uncontrolled at baseline. The
16 composite endpoint was defined as occurrence of an ambulatory exacerbation, an emergency
17 room attendance or hospital admission due to an exacerbation, or death. The results of the analysis
18 of the primary outcome have recently been published (8). The current manuscript describes the
19 changes of control status during follow-up and the short-term (6 months) predictive value of the
20 lack of control in patients with COPD.
21
22
23
24
25
26
27
28
29

30 The study was approved by the local Research and Ethics Committees of each participating
31 center, and all patients provided written informed consent. The data from the United Kingdom
32 (UK) was obtained from the Optimum Patient Care Research Database (OPCRD) and permission
33 to access and link UK data to anonymous electronic medical records was obtained from the Health
34 Research Authority for clinical research use (Anonymised Data Ethics & Protocol Transparency
35 (ADEPT) approval number ADEPT0115). This study was registered with the European Network
36 of Centres for Pharmacoepidemiology and Pharmacovigilance (ENCePP), Register Number
37 EUPAS10679.
38
39
40
41
42
43
44
45
46
47
48
49

50 **Population**

51 The inclusion criteria included: age over 40 years, spirometry-defined COPD (i.e. post-
52 bronchodilator forced expiratory volume in 1 second (FEV₁) / forced vital capacity (FVC) <0.7),
53 current or ex-smoker of at least 10 pack-years and being in a stable clinical state at the screening
54 visit. Patients were excluded if they: 1) had any chronic concomitant respiratory condition other
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 than asthma or bronchiectasis; 2) were unable to understand the instructions of the study or fill in
5
6 the questionnaires; 3) had severe comorbidity with a life expectancy shorter than 2 years; or 4)
7
8 were participating in another clinical study or clinical trial.
9

10 11 **Definition of control**

12
13
14 A patient was considered controlled when COPD had a low impact and was clinically stable, after
15
16 adjustment for the severity of the disease (7). Impact was classified as low or high according to
17
18 the information collected in sputum (presence and color), breathlessness (mMRC), daily physical
19
20 activity and use of rescue medication. Evaluation of impact was adjusted for disease severity
21
22 according to the FEV₁. Patients with a FEV₁(%) \geq 50% predicted were classified as
23
24 mild/moderate and those with FEV₁(%) $<$ 50% as severe. Stability was defined as the absence of
25
26 moderate or severe exacerbations in the previous 3 months (Table 1).
27
28
29
30

31 **Statistical analysis**

32
33 Absolute frequencies and percentages were used for comparisons of qualitative variables. The
34
35 description of quantitative variables was performed using the mean and standard deviation (SD).
36
37 The Kolmogorov-Smirnov test was used to assess the normality of distributions. In the case of
38
39 quantitative variables, comparison of the characteristics among groups was carried out using the
40
41 Student t-test (Mann-Whitney U-test if normality was not assumed). The Chi-squared test (Fisher
42
43 test for frequencies $<$ 5) was employed for the comparison of categorical variables.
44
45
46
47

48 Multivariate logistic regression analysis was performed with the independent variable being
49
50 persistent control status and dependent variables all other demographic and clinical variables,
51
52 excluding those already used to describe control status.
53
54

55 Odd ratios with 95% confidence intervals (CI) were used to assess the risk of an outcome in the
56
57 6 months after an uncontrolled visit compared to a controlled visit. For all the tests p -values $<$ 0.05
58
59
60
61
62
63
64
65

1
2
3
4 were considered statistically significant. The statistical package R Studio (V2.5.1) was used for
5
6 the statistical analyses.
7

8 **RESULTS**

9 **Population**

10
11 A total of 349 patients were screened, 303 (87%) of whom fulfilled all the inclusion and exclusion
12
13 criteria and could be evaluated for control status at the baseline visit (V0). The mean age was 68.6
14
15 years (SD= 8.7], 73.9% were male, the mean CAT score was 14.4 (8.6) and the mean FEV₁(%)
16
17 was 52.5% (18.1%). During the 18-month follow-up, 26 patients were lost, and 10 patients died.
18
19 The characteristics of the whole population and the flow chart of the study have been described
20
21 in detail in a previous publication (8).
22
23
24
25
26
27

28 **Control status at baseline and during follow-up**

29
30 At baseline, 197 (65%) patients were classified as controlled, 68.5% being mild/moderate and
31
32 59.3% severe COPD patients. Of them, 128 (65%) persisted controlled at 6 months, 106 (84% or
33
34 54% of the initial 197) at 12 months and 80 (75.4% or 40.6% of the initial 197) at 18 months.
35
36 From the 106 (35%) uncontrolled at baseline, 70 (66%) persisted uncontrolled at 6 months, 54
37
38 (77% or 51% of the initial 106) at 12 months and 43 (79.6% or 40.5% of the initial 106) at 18
39
40 months. Therefore, at the end of the 18-month follow-up, 80 (29.8%) patients remained
41
42 controlled, 43 (16%) persistently uncontrolled and the remaining 144 (53.7%) changed control
43
44 status during follow-up (Figure 1)
45
46
47
48
49

50 **Characteristics of patients and outcomes according to changes in control status during** 51 **follow-up**

52
53 In the univariate analysis, there were significant differences in several variables among patients
54
55 who were persistently controlled, had variable control status and were persistently uncontrolled
56
57 (Table 2). Persistently uncontrolled patients were more frequently women, with a higher BMI,
58
59
60
61
62
63
64
65

1
2
3
4 more bronchiectasis, worse dyspnea and lung function parameters, more moderate or severe
5
6 exacerbations in the previous year, few minutes walked per day and worse BODEx and CAT
7
8 scores.
9

10
11 The same trends were observed for outcomes during follow-up in persistently controlled patients
12
13 not presenting any exacerbation (by definition) and persistently uncontrolled patients presenting
14
15 a significantly higher number of all types of exacerbations compared with patients with variable
16
17 control status (Table 3).
18
19

20
21 Multivariate analysis showed that the characteristics independently and significantly associated
22
23 with persistent control status were male sex, a lower (not increased) BMI and a better FEV₁(%)
24
25 (table 4).
26
27

28 **Short-term outcomes after each visit according to control status**

29
30 Of the 874 clinical visits with a 6-month follow-up that were valid for analysis of short-term
31
32 outcomes, 511(58.4%) fulfilled the criteria of controlled and 363 (41.5%) were considered
33
34 uncontrolled. All the outcomes were more frequent during the 6 months after an uncontrolled
35
36 patient visit compared to the controlled patient visits (10.9% and 4%, respectively with an
37
38 emergency room visit and 40.4% and 16.6%, respectively with an ambulatory exacerbation;
39
40 p<0.001 for both) (Figure 2). There were 10 deaths, 3 (0.6%) after controlled patient and 7 (1.9%)
41
42 after uncontrolled patient visits.
43
44
45
46

47
48 Uncontrolled patient visits resulted in a highly significant increased risk of poor outcomes during
49
50 the next 6 months with an OR between 2.94 for emergency visits to 4.25 for hospitalizations due
51
52 to exacerbations compared with controlled patient visits (Table 5, Figure 3). We investigated
53
54 whether having an exacerbation in the previous 3 months (instability) could offer the same
55
56 predictive value as the control status. Odds ratios for the different outcomes were also significant
57
58 but of lower magnitude than those of the control status (data not shown). However, there were
59
60
61
62
63
64
65

1
2
3
4 119 visits (32.8% of those classified as uncontrolled) that did not have an exacerbation in the
5
6 previous 3 months and therefore would have been missed as uncontrolled if only the criteria of
7
8 previous exacerbation had been used
9

10 11 12 13 14 **DISCUSSION**

15
16 According to the previously defined control criteria, more than half of the population of our study
17
18 was classified as controlled at baseline; however, the control status changed frequently during the
19
20 18-month follow-up. Only 30% of patients remained controlled and 16% persistently
21
22 uncontrolled throughout the observation period, while the control status changed in 54% of
23
24 patients during follow-up.
25
26

27
28 Patients who remained controlled over the course of the study were more frequently men, with a
29
30 normal (not increased) BMI and with a more preserved FEV₁(%). Uncontrolled status had a great
31
32 predictive value for adverse outcomes with 3-fold higher risk of an emergency room visit for an
33
34 exacerbation, a 3.4-fold increased risk of an ambulatory exacerbation and a 4-fold increased risk
35
36 of hospitalization during the next 6 months. These results suggest that lack of control is a useful
37
38 warning sign that should prompt investigation of the causes and intervention in order to minimize
39
40 the short-term risk of exacerbations.
41
42
43
44

45
46 Control is a term frequently used to describe a status in which the disease has a minimal impact
47
48 on the health status of the patient and the prognosis is the best possible. Intuitively, when a chronic
49
50 disease is uncontrolled it should require more careful evaluation and action taking to improve the
51
52 symptoms and/or prognosis (17,18). This concept has been very well established in asthma (2)
53
54 but has been elusive in COPD for a variety of reasons, mainly because of a lack of control
55
56 standard. Since COPD is defined as a chronic, progressive, and non fully reversible airflow
57
58 limitation, control cannot be defined as the normalization of lung function and disappearance of
59
60
61
62
63
64
65

1
2
3
4 symptoms and exacerbations. Instead, we need to define a tool that includes some easy to obtain
5
6 clinical characteristics with meaningful thresholds adjusted for severity, that allow the
7
8 identification of “uncontrolled” patients and which has prognostic value for relevant outcomes.
9
10 Control in COPD was initially defined as including two dimensions: impact and stability (5), and
11
12 the criteria were refined and validated in subsequent prospective studies (6,8,19,20).
13
14

15
16 The criteria selected are easy to obtain at each clinical visit and basically consist in variables that
17
18 patients should be questioned about, such as the level of dyspnea, sputum production and
19
20 characteristics, the level of physical activity and the use of rescue medication, in addition to the
21
22 history of previous moderate or severe exacerbations (21). Using the thresholds derived from
23
24 previous studies (19), it has been demonstrated that control status is a good predictor of long-term
25
26 outcomes (8,19,20), and that control evaluation based on clinical data is a better predictor than
27
28 the definition of control based on CAT scores (19,20). Therefore, we used these clinical criteria
29
30 in the current analysis.
31
32
33

34
35 The last revision of the GOLD strategic document separated, for the first time, the
36
37 recommendations for initial treatment of COPD and changes in treatment based on a reassessment
38
39 of the patients in subsequent visits (1). In this context, the control evaluation can be a useful tool
40
41 to reassess the clinical status of the patients and their future risks. A previous study using the
42
43 same criteria of control demonstrated that the control tool is more sensitive to changes in the
44
45 clinical status of the patient than changes in phenotype, the level of risk according to the Spanish
46
47 COPD guidelines (22), or the GOLD A-D classification frequently used to guide therapy (7). In
48
49 addition, changes in control status were clinically relevant because they resulted in changes in
50
51 health status over a period of only 3 months (7). However, the importance of the evaluation of
52
53 control status at each clinical visit was not demonstrated. In this study, we observed that the
54
55 control status of the patients frequently changes in subsequent clinical visits separated by 6
56
57
58
59
60
61
62
63
64
65

1
2
3
4 months, and more importantly, after an uncontrolled patient visit, the risk of having a moderate
5 or severe exacerbation in the following 6 months is increased by 3- to 4-fold according to the type
6 of episode. This increased risk justifies the use of control evaluation as a warning sign to foster
7 more careful evaluation of the patients and the adoption of therapeutic measures according to the
8 results of these investigations. In contrast, if the patient fulfils the criteria of control during a
9 clinical visit, we may be quite confident that the risk of an adverse outcome during the next
10 months is low. Therefore, the control status could be used to reduce or extend the time period
11 between routine follow-up visits, with uncontrolled patients probably requiring more frequent
12 appointments compared to controlled patients.
13

14
15
16
17
18
19
20
21
22
23
24
25
26 Regarding the limitations of the present study, it is important to consider that control status does
27 not provide any specific diagnosis; a patient may be uncontrolled for a variety of reasons. The
28 uncontrolled status is only an alert to take action but does not tell us what action to take. However,
29 the control tool can be an excellent reminder to the clinician about the questions that COPD
30 patients must be asked at each clinical visit and provide a simple prognostic tool. Only one quarter
31 of the population were women, so extrapolation of the results to women must be made with
32 caution.
33

34
35
36
37
38
39
40
41
42
43 The results of the current study are in agreement with those of previous studies demonstrating the
44 predictive value of control status. The results obtained in our study highlight the significantly
45 increased risk of moderate or severe exacerbations during the following 6 months in patients
46 classified as uncontrolled. If these results are confirmed in other cohorts, control criteria should
47 be incorporated into clinical practice as a simple to use tool to help reassess COPD patients at
48 each follow-up visit.
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

REFERENCES

1. Singh D, Agusti A, Anzueto A, Barnes PJ, Bourbeau J, Celli BR, et al. Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Lung Disease: the GOLD science committee report 2019. *Eur Respir J*. 2019;53(5). pii: 1900164
2. Chapman KR, Boulet LP, Rea RM, Franssen E. Suboptimal asthma control: prevalence, detection and consequences in general practice. *Eur Respir J* 2008; 31(2):320-325.
3. Soler-Cataluña JJ, Alcázar B, Miravittles M. Clinical Control in COPD: A New Therapeutic Objective? *Arch Bronconeumol*. 2020; 56: 68-69.
4. de-Torres JP, Marín JM, Pinto-Plata V, Divo M, Sanchez-Salcedo P, Zagaceta J, et al. Is COPD a Progressive Disease? A Long Term Bode Cohort Observation. *PLoS One*. 2016; 11(4):e0151856.
5. Soler-Cataluña JJ, Alcazar-Navarrete B, Miravittles M. The concept of control in COPD: a new proposal for optimising therapy. *Eur Respir J* 2014; 44: 1072-1075.
6. Soler-Cataluña JJ, Alcazar B, Miravittles M. The concept of control of COPD in clinical practice. *Int J Chron ObstructPulmon Dis* 2014; 9: 1397-1405.
7. Soler-Cataluña JJ, Alcazar B, Marzo M, Pérez J, Miravittles M. Evaluation Of Changes In Control Status In COPD: An Opportunity For Early Intervention. *Chest*. 2019 Nov 21. pii: S0012-3692(19)34222-9. doi: 10.1016/j.chest.2019.11.004. [Epub ahead of print]
8. Miravittles M, Sliwinski P, Rhee CK, Costello RW, Carter V, Tan J, et al. Predictive value of control of COPD for risk of exacerbations: An international, prospective study. *Respirology* 2020 Apr 6. doi: 10.1111/resp.13811. [Epub ahead of print].

- 1
2
3
4 9. Miravittles M, Sliwinski P, Rhee CK, Costello RW, Carter V, Tan J, et al. Evaluation criteria for
5
6 clinical control in a prospective, international, multicenter study of patients with COPD. *Respir Med*
7
8 2018; 136: 8-14.
- 10
11 10. Kim KY, Miravittles M, Sliwinski P, Costello R, Carter V, Tan J, et al. Comparison of clinical
12
13 baseline characteristics between Asian and Western COPD patients in a prospective, international,
14
15 multicenter study. *Int J Chron Obstruct Pulmon Dis*. 2019; 14: 1595-1601.
- 16
17
18 11. Soler-Cataluña JJ, Martínez-García MA, Sánchez L, Perpiña M, Román P. Severe exacerbations
19
20 and BODE index: two independent risk factors for death in male COPD patients. *Respir Med* 2009;
21
22 103: 692-699.
- 23
24
25 12. Jones PW, Harding G, Berry P, Wiklund I, Chen WH, Kline Leidy N. Development and first
26
27 validation of the COPD Assessment Test. *Eur Respir J* 2009; 34: 648-654.
- 28
29
30 13. Bestall JC, Paul EA, Garrod R, Garnham R, Jones PW, Wedzicha JA. Usefulness of the
31
32 Medical Research Council (MRC) dyspnoea scale as a measure of disability in patients with
33
34 chronic obstructive pulmonary disease. *Thorax* 1999; 54: 581–586.
- 35
36
37
38 14. Charlson M, Szatrowski TP, Peterson J, Gold J. Validation of a combined comorbidity index.
39
40 *J Clin Epidemiol* 1994; 47: 1245–1251.
- 41
42
43
44 15. Ramon MA, Esquinas C, Barrecheguren M, Pleguezuelos E, Molina J, Quintano JA, et al.
45
46 Self-reported daily walking time in COPD: relationship with relevant clinical and functional
47
48 characteristics. *Int J Chron Obstruct Pulmon Dis* 2017; 12: 1173-1181.
- 49
50
51
52 16. Wedzicha JA, Miravittles M, Hurst JR, Calverley PMA, Albert RK, Anzueto A, et al.
53
54 Management of COPD exacerbations: An European Respiratory Society/American Thoracic Society
55
56 (ERS/ATS) guideline. *Eur Respir J* 2017; 49: pii 1600791.
- 57
58
59
60 17. Duarte Araújo A, Hespanhol V, Correia-de-Sousa J. Is COPD control a useful concept?
61
62
63
64
65

1
2
3
4 Assessing treatment success by evaluating COPD-related health status. Arch Bronconeumol. 2017;
5
6 53(9): 530-531.
7

8
9 18. Guimarães M, Bugalho A, Oliveira AS, Moita J, Marques A; GI DPOC-Grupo de Interesse na
10
11 Doença Pulmonar Obstrutiva Crónica. COPD control: Can a consensus be found? Rev Port Pneumol
12
13 2016; 22(3): 167-176.
14

15
16 19. Soler-Cataluña JJ, Marzo M, Catalán P, Miralles C, Alcazar B, Miravittles M. Validation of
17
18 clinical control in COPD as a new tool for optimizing treatment. Int J Chron Obstruct Pulmon Dis.
19
20 2018; 13: 3719-3731.
21

22
23 20. Barrecheguren M, Kostikas K, Mezzi K, Shen S, Alcazar B, Soler-Cataluña JJ, et al. COPD
24
25 clinical control as a predictor of future exacerbations: concept validation in the SPARK study
26
27 population. Thorax. 2020; 75: 351-353.
28

29
30 21. Calle Rubio M, Rodríguez Hermosa JL, Soler-Cataluña JJ, López-Campos JL, Alcazar Navarrete
31
32 B, Soriano JB, et al. Medical care according to risk level and adaptation to Spanish COPD Guidelines
33
34 (GesEPOC): The Epoconsul Study. Arch Bronconeumol 2018; 54: 270-279.
35

36
37 22. Erro Iribarren M, Alonso Pérez T, Soriano JB, Ancochea Bermúdez J. Adjusting the Level of
38
39 Intervention in Patients with Chronic Obstructive Pulmonary Disease According to the Risk
40
41 Stratification Proposed by the Spanish COPD Guidelines (GesEPOC) Version 2017. Arch
42
43 Bronconeumol 2020; 56(3): 183-185.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

FIGURE LEGENDS

Figure 1. Control status and changes in control status at baseline and follow-up visits

Figure 2. Percentages of outcomes during the following 6 months after each controlled (n=511) and uncontrolled (n=363) visits. Exacerbations, emergency visits and hospitalisations are caused by COPD

Figure 3. Odds ratios for each outcome during the following 6 months after uncontrolled (n=363) compared with controlled (n=511) visits. Exacerbations, emergency visits and hospitalisations are caused by COPD

Table 1: Control criteria with adjustment of severity according to FEV₁%.

CONTROL CRITERIA	
Low Clinical Impact (Must meet at least 3 of the 4 criteria)	
- Dyspnea	0-1 if FEV ₁ ≥ 50% 0-2 if FEV ₁ < 50%
- Rescue medication	≤ 3 times/week
- Sputum color	White or absent
- Physical activity	≥ 30 min/day
Clinical Stability	
- Moderate or severe exacerbations in the last 3 months.	None
Control	Low impact + Stability

Table 2. Demographic and clinical characteristics of patients according to control status at the different follow-up visits.

Baseline characteristics	All patients (N=267)	Always Controlled (N=80)	Variable Control (N=144)	Always Uncontrolled (N=43)	P value
Age, years	68.3 (8.6)	68.5 (8.7)	68.3 (8.7)	68.1 (8.3)	0.916
Men, n (%)	196 (73.4)	72 (90.0)	102 (70.8)	22 (51.2)	<0.001
Active smokers, n (%)	71 (26.6)	18 (22.5)	42 (29.2)	11 (25.6)	0.549
Pack-years	47.6 (30.2)	43.5 (24.3)	47.8 (31.3)	54.5 (35.5)	0.390
BMI (Kg/m ²)	26.7 (5.3)	25.3 (5.2)	26.9 (5.1)	28.7 (5.7)	0.001
Chronic bronchitis, n (%)	136 (59.9)	46 (60.5)	66 (57.9)	24 (64.9)	0.829
Emphysema, n (%)	175 (77.1)	61 (80.3)	88 (77.2)	26 (70.3)	0.494
ACO, n (%)	27 (10.1)	10 (12.5)	14 (9.7)	3 (7)	0.609
Bronchiectasis, n (%)	35 (13.1)	15 (18.8)	10 (6.9)	10 (23.3)	0.004
Charlson index	4.2 (1.6)	4.0 (1.6)	4.3 (1.5)	4.4 (1.8)	0.261
mMRC	1.5 (1)	0.9 (0.8)	1.1 (0.6)	1.9 (1)	<0.001
FVC, mL	2945 (892)	3303 (865)	2857 (894)	2581 (712)	<0.001
FVC (%)	67.5 (14.5)	71.6 (15.2)	66.5 (13.5)	63.8 (14.1)	<0.001
FEV1, mL	1512 (582)	1711 (607)	1466 (573)	1302 (459)	<0.001
FEV1 (%)	53.0 (17.9)	59.3 (20.1)	50.5 (16.5)	49.5 (15.1)	0.004
Exacerbations in the previous year	1.4 (2.5)	0.3 (0.8)	1.4 (2.2)	3.6 (3.9)	<0.001
BODEx index	2.3 (1.7)	1.5 (1.3)	2.4 (1.6)	3.2 (1.8)	<0.001
CAT score	14.2 (7.9)	10.8 (5.9)	13.7 (7.2)	22.3 (7.9)	<0.001
Minutes walked/day	88.0 (72.7)	104.5 (75.3)	87.5 (74.1)	58.8 (52.4)	<0.001

Footnote: BMI: Body mass index; ACO: Asthma-COPD overlap; mMRC: modified Medical Research council dyspnea scale; FVC: Forced vital capacity; FEV1: Forced expiratory volume in 1 second; BODEx: Body mass index, obstruction, dyspnea and exacerbations index; CAT: COPD Assessment Test.

Table 3. Outcomes over the 18-month follow-up of COPD patients according to control status at the different follow-up visits. **Exacerbations, emergency visits and hospitalisations are caused by COPD**

Outcomes during follow-up	All patients (N=271)	Always Controlled (N=80)	Variable Control (N=144)	Always Uncontrolled (N=43)	P value
Combined event, n (%)	152 (56.1)	0 (0)	109 (75.7)	40 (93.0)	<0.001
Combined event, mean (SD)	1.7 (2.6)	0 (0)	1.9 (2.3)	4.2 (3.6)	<0.001
Hospitalization, n (%)	48 (17.7)	0 (0)	28 (19.4)	19 (44.2)	<0.001
Hospitalization, mean (SD)	0.3 (0.9)	0 (0)	0.3 (0.9)	0.8 (1.1)	<0.001
Emergency, n (%)	41 (15.1)	0 (0)	26 (18.1)	14 (32.6)	<0.001
Emergency, mean (SD)	0.3 (0.8)	0 (0)	0.3 (0.8)	0.7 (1.2)	<0.001
Ambulatory exacerbations, n (%)	136 (50.2)	0 (0)	97 (67.4)	36 (83.7)	<0.001
Ambulatory exacerbations, mean (SD)	1.1 (1.6)	0 (0)	1.2 (1.3)	2.7 (2.6)	<0.001

Table 4. Univariate and multivariate logistic regression analyses for controlled patients during follow-up (I).

Variable	Univariate		Multivariate ^a	
	OR (95% CI)	P value	OR (95% CI)	P value
Age (+1 year)	1.00 (0.97 to 1.03)	0.836	-	-
Men	4.57 (2.07 to 10.08)	<0.001	3.20 (1.41 to 7.28)	0.005
Pack-years (+1)	0.99 (0.98 to 1.00)	0.150	-	-
BMI (+1 Kg/m ²)	0.93 (0.88 to 0.98)	0.005	0.93 (0.88 to 0.98)	0.010
Chronic bronchitis	1.04 (0.59 to 1.83)	0.893	-	-
Emphysema	1.32 (0.67 to 2.59)	0.421	-	-
ACO	1.43 (0.62 to 3.27)	0.399	-	-
Bronchiectasis	1.93 (0.93 to 3.99)	0.077	-	-
Charlson index (+1)	0.87 (0.72 to 1.05)	0.141	-	-
FEV ₁ (+100 mL)	1.09 (1.04 to 1.14)	<0.001	1.07 (1.02 to 1.12)	0.006

Footnote: OR: Odds ratio; CI: Confidence interval; BMI: Body mass index; ACO: Asthma-COPD overlap; FEV₁: Forced expiratory volume in 1 second.

^a P value for the Hosmer-Lemeshow test for goodness-of-fit: 0.18.

Table 5. Short-term outcomes at 6 months after each clinical visit according to control status at that visit. **Exacerbations, emergency visits and hospitalisations are caused by COPD.**

Number of visits with outcome during the next 6 months	All follow-up visits up to V2 (N=874)		OR (95%CI)	P value
	Controlled visits (N=511)	Uncontrolled visits (N=363)		
Combined event, n (%)	105 (20.5)	169 (46.6)	3.37 (2.50 to 4.54)	<0.001
Hospitalization, n (%)	20 (4)	51 (15)	4.25 (2.48 to 7.27)	<0.001
Emergency visits, n (%)	20 (4)	37 (10.9)	2.94 (1.68 to 5.16)	<0.001
Ambulatory exacerbations, n (%)	83 (16.6)	137 (40.4)	3.41 (2.47 to 4.69)	<0.001
Deaths (%)	3 (0.6)	7 (1.9)	3.33 (0.86 to 12.96)	0.083

Footnote: V2: Visit 2; OR: Odds ratio; CI: Confidence interval. Combined event was defined as occurrence of an ambulatory exacerbation, an emergency room attendance or hospital admission due to an exacerbation, or death. Emergency visits and hospitalizations were due to exacerbations of COPD.

FIGURES
Figure 1







Figure 2



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65


Figure 3





[Click here to access/download](#)

Otros documentos (Supplementary files)
control status REG for ABN R1 CLEAN.docx



Changes in control status of COPD over time and their consequences: A prospective international, study.

The current study is the first prospective, international, multicenter study designed with the objective to investigate the concept of control of COPD. In a previous publication we described control status as a marker of increased risk of poor outcomes during follow-up. Here, we describe the changes in control status in patients over long-term follow-up, the short-term risk associated with uncontrolled patient visits and the main characteristics of patients that remained either controlled or uncontrolled during the study.





Figure 3. Odds ratios for each outcome during the following 6 months after uncontrolled (n=363) compared with controlled (n=511) visits. Exacerbations, emergency visits and hospitalisations are caused by COPD

Figure 2. Percentages of outcomes during the following 6 months after each controlled (n=511) and uncontrolled (n=363) visits. Exacerbations, emergency visits and hospitalisations are caused by COPD

Conclusion: The results obtained in our study highlight the significantly increased risk of moderate or severe exacerbations during the following 6 months in patients classified as uncontrolled. If these results are confirmed in other cohorts, control criteria should be incorporated into clinical practice as a simple to use tool to help reassess COPD patients at each follow-up visit.

Ética de la publicación

1. ¿Su trabajo ha comportado experimentación en animales?:

No

2. ¿En su trabajo intervienen pacientes o sujetos humanos?:

Sí

- Si la respuesta es afirmativa, por favor, mencione el comité ético que aprobó la investigación.:

CEIC Hospital Vall d'Hebron, Barcelona

- Si la respuesta es afirmativa, por favor, confirme que los autores han cumplido las normas éticas relevantes para la publicación. :

Sí

- Si la respuesta es afirmativa, por favor, confirme que los autores cuentan con el consentimiento informado de los pacientes. :

Sí

3. ¿Su trabajo incluye un ensayo clínico?:

No

4. ¿Todos los datos mostrados en las figuras y tablas incluidas en el manuscrito se recogen en el apartado de resultados y las conclusiones?:

Sí