





















RESEARCH ARTICLE

REVISED Cohort profile for the STRatifying Resilience and Depression Longitudinally (STRADL) study: A depression-focused investigation of Generation Scotland, using detailed clinical, cognitive, and neuroimaging assessments [version 2; peer review: 2 approved]

Tina Habota¹, Anca-Larisa Sandu ¹, Gordon D. Waiter ¹, Christopher J. McNeil¹, J. Douglas Steele ², Jennifer A. Macfarlane ³, Heather C. Whalley ⁴, Ruth Valentine¹, Dawn Younie¹, Nichola Crouch¹, Emma L. Hawkins⁴, Yoriko Hirose ⁴, Liana Romaniuk ⁴, Keith Milburn ², Gordon Buchan¹, Tessa Coupar ², Mairi Stirling², Baljit Jagpal², Beverly MacLennan¹, Lucasz Priba², Mathew A. Harris⁴, Jonathan D. Hafferty⁴, Mark J. Adams ⁴, Archie I. Campbell ⁴, Donald J. MacIntyre ⁴, Alison Pattie⁴, Lee Murphy ⁴, Rebecca M. Reynolds ⁵, Rebecca Elliot⁶, Ian S. Penton-Voak⁷, Marcus R. Munafò ⁷, Kathryn L. Evans ⁴, Jonathan R. Seckl ⁴, Joanna M. Wardlaw ⁴, Stephen M. Lawrie ⁴, Christopher S. Haley ⁴, David J. Porteous ⁴, Ian J. Deary⁴, Alison D. Murray ^{1*}, Andrew M. McIntosh ^{4*}

¹University of Aberdeen, Aberdeen, UK

²University of Dundee, Dundee, UK

³Ninewells Hospital & Medical School, Dundee, UK

⁴University of Edinburgh, Edinburgh, UK

⁵Queen's Medical Research Institute, Edinburgh, UK

⁶University of Manchester, Manchester, UK

⁷University of Bristol, Bristol, UK

* Equal contributors

v2 **First published:** 25 Nov 2019, 4:185
<https://doi.org/10.12688/wellcomeopenres.15538.1>
Latest published: 16 Jul 2021, 4:185
<https://doi.org/10.12688/wellcomeopenres.15538.2>

Open Peer Review

Reviewer Status  

Abstract

STratifying Resilience and Depression Longitudinally (STRADL) is a population-based study built on the Generation Scotland: Scottish Family Health Study (GS:SFHS) resource. The aim of STRADL is to


subtype major depressive disorder (MDD) on the basis of its aetiology, using detailed clinical, cognitive, and brain imaging assessments. The GS:SFHS provides an important opportunity to study complex gene-environment interactions, incorporating linkage to existing datasets and inclusion of early-life variables for two longitudinal birth cohorts. Specifically, data collection in STRADL included: socio-economic and lifestyle variables; physical measures; questionnaire data that assesses resilience, early-life adversity, personality, psychological health, and lifetime history of mood disorder; laboratory samples; cognitive tests; and brain magnetic resonance imaging. Some of the questionnaire and cognitive data were first assessed at the GS:SFHS baseline assessment between 2006-2011, thus providing longitudinal measures relevant to the study of depression, psychological resilience, and cognition. In addition, routinely collected historic NHS data and early-life variables are linked to STRADL data, further providing opportunities for longitudinal analysis. Recruitment has been completed and we consented and tested 1,188 participants.

Keywords

Cognition, Depression, Generation Scotland, Longitudinal, Neuroimaging, Psychological resilience



This article is included in the [Generation Scotland gateway](#).

Invited Reviewers		
	1	2
version 2 (revision) 16 Jul 2021	✓ report	
	↑	
version 1 25 Nov 2019	✗ report	✓ report
<ol style="list-style-type: none"> Debbie A. Lawlor , University of Bristol, Bristol, UK Helen Lavretsky, University of California, Los Angeles (UCLA), Los Angeles, USA 		
Any reports and responses or comments on the article can be found at the end of the article.		

Corresponding author: Tina Habota (tina.habota@abdn.ac.uk)

Author roles: **Habota T:** Formal Analysis, Investigation, Methodology, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; **Sandu AL:** Formal Analysis, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Waiter GD:** Investigation, Methodology, Writing – Review & Editing; **McNeil CJ:** Investigation, Methodology, Writing – Review & Editing; **Steele JD:** Methodology, Supervision, Writing – Review & Editing; **Macfarlane JA:** Investigation, Methodology, Writing – Review & Editing; **Whalley HC:** Conceptualization, Methodology, Writing – Review & Editing; **Valentine R:** Investigation, Writing – Review & Editing; **Younie D:** Project Administration, Writing – Review & Editing; **Crouch N:** Investigation, Writing – Review & Editing; **Hawkins EL:** Data Curation, Formal Analysis, Project Administration, Writing – Review & Editing; **Hirose Y:** Investigation, Project Administration, Writing – Review & Editing; **Romaniuk L:** Investigation, Methodology, Writing – Review & Editing; **Milburn K:** Project Administration, Writing – Review & Editing; **Buchan G:** Investigation, Software, Writing – Review & Editing; **Coupar T:** Investigation, Writing – Review & Editing; **Stirling M:** Investigation, Writing – Review & Editing; **Jagpal B:** Investigation, Project Administration, Writing – Review & Editing; **MacLennan B:** Investigation, Project Administration, Writing – Review & Editing; **Priba L:** Investigation, Writing – Review & Editing; **Harris MA:** Formal Analysis, Investigation, Writing – Review & Editing; **Hafferty JD:** Investigation, Writing – Review & Editing; **Adams MJ:** Investigation, Writing – Review & Editing; **Campbell AI:** Data Curation, Project Administration, Writing – Review & Editing; **MacIntyre DJ:** Methodology, Writing – Review & Editing; **Pattie A:** Methodology, Writing – Review & Editing; **Murphy L:** Project Administration, Writing – Review & Editing; **Reynolds RM:** Methodology, Writing – Review & Editing; **Elliot R:** Conceptualization, Methodology, Writing – Review & Editing; **Penton-Voak IS:** Conceptualization, Methodology, Writing – Review & Editing; **Munafò MR:** Conceptualization, Methodology, Writing – Review & Editing; **Evans KL:** Conceptualization, Funding Acquisition, Methodology, Writing – Review & Editing; **Seckl JR:** Conceptualization, Funding Acquisition, Methodology, Writing – Review & Editing; **Wardlaw JM:** Conceptualization, Funding Acquisition, Methodology, Writing – Review & Editing; **Lawrie SM:** Conceptualization, Funding Acquisition, Writing – Review & Editing; **Haley CS:** Conceptualization, Funding Acquisition, Methodology, Writing – Review & Editing; **Porteous DJ:** Conceptualization, Funding Acquisition, Methodology, Writing – Review & Editing; **Deary IJ:** Conceptualization, Funding Acquisition, Methodology, Writing – Review & Editing; **Murray AD:** Conceptualization, Funding Acquisition, Methodology, Supervision, Writing – Review & Editing; **McIntosh AM:** Conceptualization, Funding Acquisition, Methodology, Supervision, Writing – Review & Editing

Competing interests: ISP-V and MRM are co-directors of Jericoe Ltd, a company that designs software for the assessment and modification of emotion recognition.

Grant information: STRADL is supported by the Wellcome Trust through a Strategic Award (104036/Z/14/Z). GS:SFHS received core support from the CSO of the Scottish Government Health Directorates (CZD/16/6) and the Scottish Funding Council (HR03006). ADM is supported by Innovate UK, the European Commission, the Scottish Funding Council via the Scottish Imaging Network SINAPSE, and the CSO. HCW is supported by a JMAS SIM Fellowship from the Royal College of Physicians of Edinburgh, by an ESAT College Fellowship from the University of Edinburgh, and has received previous funding from the Sackler Trust. LR has previously received financial support from Pfizer (formerly Wyeth) in relation to imaging studies of people with schizophrenia and bipolar disorder. JDH is supported by the MRC. DJM is an NRS Clinician, funded by the CSO. RMR is supported by the British Heart Foundation. ISP-V and MRM are supported by the NIHR Biomedical Research Centre at the University Hospitals Bristol NHS Foundation Trust and the University of Bristol. The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health; and MRM is also supported by the MRC (MC_UU_00011/7). JMW is supported by MRC UK Dementia Research Institute which receives funding from the UK Medical Research Council, Alzheimer's Society and Alzheimer's Research UK, and MRC Centre and project grants, EPSRC, Fondation Leducq, Stroke Association, British Heart Foundation, Alzheimer Society, and the European Union H2020 PHC-03-15 SVDs@Target grant agreement (666881). DJP is supported by Wellcome Trust Longitudinal Population Study funding (216767/Z/19/Z) the Eva Lester bequest to the University of Edinburgh. AMM is additionally supported by the MRC (MC_PC_17209, MC_PC_MR/R01910X/1, MR/S035818/1), The Wellcome Trust (216767/Z/19/Z), The Sackler Trust, and has previously received research funding from Pfizer, Eli Lilly, and Janssen. Both AMM and IJD are members of The University of Edinburgh Centre for Cognitive Ageing and Cognitive Epidemiology, part of the cross council Lifelong Health and Wellbeing Initiative (MR/K026992/1); funding from the BBSRC and MRC is gratefully acknowledged.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2021 Habota T *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Habota T, Sandu AL, Waiter GD *et al.* **Cohort profile for the STRatifying Resilience and Depression Longitudinally (STRADL) study: A depression-focused investigation of Generation Scotland, using detailed clinical, cognitive, and neuroimaging assessments [version 2; peer reviewed: 2 approved]** Wellcome Open Research 2021, 4:185 <https://doi.org/10.12688/wellcomeopenres.15538.2>

First published: 25 Nov 2019, 4:185 <https://doi.org/10.12688/wellcomeopenres.15538.1>

REVISED Amendments from Version 1

The revised manuscript includes amendments to the body of the paper in several areas, along with some changes to [Table 1](#) and [Figure 1](#). More specifically, the Introduction now describes in greater detail the focus of stratification of MDD and resilience, and we describe some specific initial hypotheses. The Introduction also now more clearly describes the overlap and distinction between the current wave of STRADL and studies on which the current project was built on – specifically, the wider Generation Scotland population and the first wave of STRADL. Related to this point, we include an additional table ([Table 1](#)), which shows a list of data linkages between the current study and existing datasets. We also amended [Figure 1](#) so that it now shows the recruitment and attrition for the first wave of STRADL, as well as the current wave of STRADL. The Methods now includes a clearer and more detailed overview of the wider Generation Scotland population, specifically the selection process and recruitment criteria. The Results includes further analysis of demographic (i.e., age) differences between the current study and existing datasets. In the Discussion we present a more detailed and clearer overview of the study limitations, particularly focusing on how the dataset for the current study may be influenced by cultural and societal norms and/or possible selection bias.

Any further responses from the reviewers can be found at the end of the article

Introduction**Why was the study set up?**

Major depressive disorder (MDD) affects approximately 13% of the population at least once in their lifetime¹, and remains a leading cause of economic burden and non-lethal global disability^{2,3} due to its recurrent or chronic nature. At present, MDD diagnosis is based on arbitrary and clinically heterogeneous criteria⁴. Consequently, and even with optimal management, much of the disability caused by MDD persists⁵ because of the absence of targeted disease-modifying treatments. The underlying pathophysiology of MDD is believed to be heterogeneous⁶, with genetic and environmental factors acting to influence disease expression. Thus, it is important for treatment to shift from the current “trial and error” approach, towards precision prevention and stratified medicine based on markedly different disease mechanisms. However, progress in this area has been severely restricted because the aetiology of MDD is complex, and remains poorly understood.

Stratifying Resilience and Depression Longitudinally (STRADL) aims to subtype MDD on the basis of its aetiology using detailed clinical, cognitive, and brain imaging assessments. STRADL will examine the interaction between genetic and environmental factors that increase risk and occurrence of different MDD subtypes, and assess common and distinct mechanisms and clinical trajectories of MDD phenotypes. Additionally, STRADL aims to assess individual resilience, or the ability to adapt positively and ‘avoid’ psychopathology despite exposure to known risk factors such as stress, early-life adversity, and family history⁷. Stratification of MDD will be based on several variables to address its underlying causal

and clinical heterogeneity, including: age of onset of MDD; single episode or recurrent depression; obstetric trauma; and developmental factors such as childhood maltreatment, early socioeconomic adversity, and stressful life events. Our key initial predictions are that depression can be stratified on the basis of age of onset into early-onset forms that show a stronger phenotypic and genetic relationship with schizophrenia and other severe mental disorders, and later onsets that show stronger associations with cardiovascular disease and dementia.

STRADL was built on the Generation Scotland: Scottish Family Health Study resource (GS:SFHS)⁸, which undertook its first major baseline assessments between 2006 and 2011. GS:SFHS is a population-based study of genetic health and complex disease in a cohort of 24,096 individuals, who have been extensively phenotyped for MDD and related traits. This cohort provides an important opportunity to study gene-environment interactions, and remains one of the richest sources of data available, incorporating linkage of existing phenotypic and genomic data, detailed lifestyle and socioeconomic characterisation, extensive eHealth Record linkage⁹, and the inclusion of two longitudinal birth cohorts – the Walker birth cohort¹⁰, and Aberdeen Children of the 1950s (ACONF)¹¹. [Table 1](#) shows data linkages between the current study and existing datasets.

The first wave of STRADL included depression-focused follow-up assessment of GS:SFHS, which involved remote questionnaires that specifically assessed aspects of psychological resilience, coping style, and response to psychological distress; study protocol and cohort characteristics are described elsewhere¹². Here, we describe the second wave of STRADL, a depression-focused deep phenotyping face-to-face assessment, using detailed clinical and cognitive tests, and neuroimaging. The results describe the cohort profile and baseline questionnaire and cognitive data, and we provide a summary of key demographic data from the current wave of STRADL, compared to STRADL remote follow-up and wider GS:SFHS baseline assessment. A summary of all data available and the proportion of valid and useable data is also provided.

Methods**Who is in the cohort?**

We aim to study people both with and without depression, and therefore our recruitment targeted the whole GS:SFHS population, not merely people with a depression history. GS:SFHS included participants aged 35–65 years who were identified at random from collaborating medical practices across Scotland, with some family members further afield. Initially, only Glasgow and Tayside areas were involved, but the study was extended in 2010 to include Ayrshire, Arran and Northeast Scotland, with the age range also broadened (to 18–65 years). Participants were included if: they met the age criteria; had capacity to give informed consent; and could identify at least one first-degree relative who would also participate. Follow-up of participants was done through the NHS Scotland Community Health Index (CHI): 7% of the original cohort could not be matched; no participants withdrew; and ~1,200 had died. Those who

Table 1. STRADL face-to-face linkage to existing studies and permanently linked datasets.

Database	Description	Data acquired
<i>GS:SFHS</i>	GS:SFHS baseline assessment	2006 – 2011
<i>STRADL</i>	STRADL remote questionnaire follow-up	2015 – 2017
<i>Birth cohorts</i>		
ACONF	Aberdeen Children of the 1950s	1950 – 1956
The Walker Cohort	The Walker Project	1952 – 1966
<i>Routine linkage</i>		
Outpatient Attendance (SMR00)	New and follow-up outpatient attendance	1996 – present
General/Acute Inpatient and Day Case (SMR01)	Hospital inpatients discharged from non-obstetric and non-psychiatric specialties	1981 – present
Maternity Inpatient and Day Case (SMR02)	Hospital inpatient discharges from obstetrics specialties	1975 – present
Mental Health Inpatient and Day Case (SMR04)	Psychiatric hospital discharges and diagnostic information	1981 – present
Scottish Cancer Registry (SMR06)	Personal, demographic, and diagnostic information on all new cases of cancer	1958 – present
Neonatal (SMR11)	Neonatal discharges submitted for babies who are sick or have congenital anomalies	1957 – 2002
National Records of Scotland – Deaths	Death registrations	1974 – present
Scottish Drug Misuse Database	Problem drug use	1990 – present
Prescribing Information System	Prescriptions prescribed, dispensed and reimbursed with the community setting	1993 – present

participated in GS:SFHS were invited to take part in the STRADL remote follow-up based on the following eligibility criteria : they had given consent for re-contact; were living in Scotland; and had a CHI number. 9,618 GS:SFHS responded to the invitation, including 2,460 unrelated individuals and 2,460 families (7,158 individuals) of between 2 and 18 family members.

Participants in the Tayside and Grampian areas who had already taken part in GS:SFHS between 2006–2011, and who were eligible for re-contact, were sent a postal invitation by the University of Dundee Health Informatics Centre (HIC). Included in the invitation was a reply slip to indicate whether the participant would be willing to undergo face-to-face assessment and brain magnetic resonance imaging (MRI), described here. Those who replied positively were contacted by telephone by a researcher at the most local recruitment centre. In Dundee (Tayside) recruitment targeted members of the Walker cohort, and in Aberdeen (Grampian) recruitment initially targeted members of ACONF, due to the rich early-life data already available for these cohorts.

In total, 5,649 potential participants were invited to take part in the study; 576 (10.2%) were members of ACONF; 1,103

(19.5%) were members of the Walker cohort; and 3,970 (70.3%) were members of the wider GS:SFHS population. Out of these potential participants, 646 (11.4%) people declined participation at first point of contact with HIC, and we received no reply from 3,358 (59.4%) people, even after sending up to three reminders. Initially, 1,645 (29.1%) people responded positively; however, a further 170 (3.0%) declined once they were contacted by our research team or withdrew before consenting. Recruitment ended in May 2019 and we consented and tested 1,188 (72.2%) of positive respondents across Aberdeen ($n = 582$) and Dundee ($n = 606$) sites. This meant that we tested 21% of the $n = 5,649$ who were initially invited to participate. [Figure 1](#) shows the recruitment process and attrition.

What has been measured and when?

[Table 2](#) shows all variables collected in STRADL face-to-face assessments, and those that were repetitions of the GS:SFHS baseline assessment and STRADL remote questionnaire follow-up. Before any new data were collected, participants signed a consent form permitting data and samples to be shared with other researchers through a secure data management system, and provided permission to be re-contacted in the future for additional research. Consent for linkage of participant data and samples to routine NHS records was previously obtained as




Figure 1. STRADL recruitment flowchart.

Table 2. List of novel and repeated variables collected in STRADL face-to-face assessment, compared to STRADL remote follow-up and GS:SFHS baseline assessment.

Measures	New data in STRADL face-to-face assessment (current)	Repeat from STRADL remote follow-up (2015)	Repeat from GS:SFHS baseline assessment (2006–2011)
<i>Basic demographics</i>		✓	✓
<i>Health and lifestyle</i>			
Alcohol history		✓	✓
Smoking history		✓	✓
Fatigue	✓		
Loneliness	✓		
Medical and mental health history		✓	✓
List of medications			✓
Current infection(s)	✓		
Cardiovascular health		✓	✓

Measures	New data in STRADL face-to-face assessment (current)	Repeat from STRADL remote follow-up (2015)	Repeat from GS:SFHS baseline assessment (2006–2011)
<i>Physical measures</i>			
Height			✓
Weight			✓
Blood pressure			✓
Grip strength	✓		
<i>Laboratory samples</i>			
Full Blood Count	✓		
C-Reactive Protein	✓		
Blood – DNA			✓
Saliva – DNA			✓
RNA	✓		
Plasma biomarkers (EDTA)			✓
Plasma biomarkers (LiH)			✓
Hair sample	✓		
<i>Psychiatric assessment & questionnaire data</i>			
Structured Clinical Interview for DSM Disorders			✓
Brief Resilience Scale		✓	
Cannabis use	✓		
Hospital Anxiety and Depression Scale	✓		
Mood Disorder Questionnaire			✓
Quick Inventory of Depressive Symptomology	✓		
General Health Questionnaire		✓	✓
Eysenck Personality Questionnaire – Revised			✓
International Personality Item Pool	✓		
General Causality Orientations Scale	✓		
Childhood Trauma Questionnaire	✓		
<i>Cognitive tests</i>			
Logical Memory test (immediate and delayed)			✓
Digit Symbol Coding			✓
Verbal fluency			✓
Mill Hill Vocabulary test			✓
Matrix Reasoning test	✓		
Bristol Emotion Recognition test	✓		

Measures	New data in STRADL face-to-face assessment (current)	Repeat from STRADL remote follow-up (2015)	Repeat from GS:SFHS baseline assessment (2006–2011)
Affective Go/No-Go task	✓		
Cambridge Gambling task	✓		
<i>Brain MRI</i>			
T1-weighted	✓		
NimStim fearful faces task	✓		
Reward task	✓		
Resting state	✓		
Fluid attenuation inversion recover (FLAIR)	✓		
Diffusion tensor imaging (DTI)	✓		
T2-weighted	✓		
Susceptibility weighted imaging (SWI)	✓		

Abbreviations: DSM, Diagnostic and Statistical Manual of Mental Disorders; DNA, Deoxyribonucleic acid; EDTA, Ethylenediaminetetraacetic acid; LiH, Lithium Heparin; RNA, Ribonucleic acid.

part of the original GS:SFHS (05/S1401/89). All subsequent procedures were conducted following an independent, but linked, ethics application (14/SS/0039).

At each site participants attended three testing ‘stations’, which involved i) collection of clinical and questionnaire data, and biological samples ii) cognitive assessment, and iii) neuroimaging, the order of which varied at random between participants. Data from the clinical station were collected without a set order; however, cognitive tests were administered in the same order for each participant, and the MRI sequences also remained the same – except for one fMRI task, which was counterbalanced (details described in section *Brain magnetic resonance imaging*). All measures were administered in accordance with rigorous standard operating procedures based on best practice.

Clinical assessment

Medical history was updated from previous GS:SFHS baseline assessment and any new diagnoses or medical episodes were recorded. General health and lifestyle data were also collected, as were the physical measurements of height, weight, two automated measures of blood pressure, and left- and right-hand grip strength (using a Patterson Medical Jamar hand dynamometer). We collected laboratory blood samples for genetic and additional genomic analyses, including the study of DNA methylation, transcription (RNA) and protein expression. Additionally, detailed questionnaire data were collected that will be used to test the structure of depressive symptoms and their association with each measure.

Laboratory samples. A small sample of hair was collected from the posterior vertex region for longitudinal cumulative cortisol. Cortisol assays from hair samples provide a more stable marker of chronic cortisol exposure compared to cross-sectional blood or urine samples, which show considerable diurnal variation¹³. Other available assays include cortisone, testosterone, progesterone, and dehydroepiandrosterone. Venepuncture was carried out using a butterfly needle kit. Blood was extracted into the following vacutainers types (analyses in parentheses): 1) EDTA (Full blood count; FBC); 2) clot activator gel for serum separation (C-reactive protein CRP); 3) EDTA (DNA extraction); 4) 2 x Tempus RNA (RNA extraction); 5) EDTA (plasma biomarkers); 6) Lithium Heparin (plasma biomarkers). FBC and CRP samples were taken and sent to NHS laboratories for screening of clinically significant markers of anaemia and inflammation. When blood samples could not be collected, a saliva DNA collection kit (Oragene or GeneFiX) was used instead. These laboratory samples were temporarily stored at each collection site. RNA and blood DNA samples were stored at -80°C, and all others at -20°C, before being sent to the Edinburgh Clinical Research Facility at the University of Edinburgh for analysis and long-term storage. A summary of the completeness of these clinical data is shown in [Table 3](#) and [Table 4](#). FBC and CRP analysis are complete, and other blood and hair samples are in the process of being analysed.

Clinical interview and questionnaire data. All participants were assessed for a lifetime history of MDD. We used a research version of the Structured Clinical Interview for DSM-IV disorders (SCID)¹⁴ to assess symptoms of mood disorder

Table 3. Summary of demographic and background data, and proportion valid data (n = 1,188).

Measurement		%	
<i>Demographics</i>	Age	100	
	Gender	100	
	Highest education attained	99.9	
	Occupation	99.8	
	Father's occupation	98.8	
	Marital status	99.9	
<i>Health and lifestyle</i>	Alcohol history	99.7	
	Smoking history	99.7	
	Fatigue (preceding 3 months)	86.7	
	Loneliness (preceding 1 week)	99.9	
	Medical or mental health history	99.9	
	List of medications	100	
	Current infection(s)	96.6	
	<i>Cardiovascular health</i>	Stroke or transient ischaemic attack	99.9
		Myocardial infarction or angina	99.9
		Other heart disease	99.9
Peripheral arterial disease		99.9	
Jaw claudication		96.5	
Diabetes		99.9	
Hypertension		100	
Hypercholesterolaemia		99.9	

Table 4. Summary of deep phenotype data available, and proportion valid data (n = 1,188).

Measurement		%
<i>Physical measures</i>	Height	99.9
	Weight	99.9
	Blood pressure – first recording	99.7
	Blood pressure – second recording	97.1
	Grip strength	98.3
<i>Laboratory samples</i>	Full Blood Count	96.5
	C-Reactive Protein	97.1
	DNA	99.2
	RNA	96.0

Measurement		%	
	Plasma biomarkers (EDTA)	97.0	
	Plasma biomarkers (LiH)	96.6	
	Hair sample	91.0	
<i>Psychiatric assessment</i>	Structured Clinical Interview for DSM Disorders	100	
<i>Questionnaire data</i>	Brief Resilience Scale	100	
	Cannabis use	99.9	
	Hospital Anxiety and Depression Scale	100	
	Mood Disorder Questionnaire	100	
	Quick Inventory of Depressive Symptomatology	99.9	
	General Health Questionnaire	99.9	
	Eysenck Personality Questionnaire – Revised	99.9	
	International Personality Item Pool	100	
	General Causality Orientations Scale	98.5	
	Childhood Trauma Questionnaire	100	
<i>Cognitive tests</i>	Logical Memory test (immediate and delayed)	99.9	
	Digit Symbol Coding	99.2	
	Verbal fluency	99.9	
	Mill Hill Vocabulary test	99.9	
	Matrix Reasoning test	99.9	
	Bristol Emotion Recognition test	98.0	
	Affective Go/No-Go task	97.7	
	Cambridge Gambling task	98.6	
	<i>Brain MRI data</i>	T1-weighted	91.2
		NimStim fearful faces task	89.1
Reward task		89.0	
Resting state		90.2	
Fluid attenuation inversion recover (FLAIR)		90.6	
Diffusion tensor imaging (DTI)		89.3	
	T2-weighted	89.4	
	Susceptibility weighted imaging (SWI)	90.6	

Note. 96.5% of DNA was extracted via blood sample and 2.7% via saliva sampling.

Abbreviations: DSM, Diagnostic and Statistical Manual of Mental Disorders; DNA, Deoxyribonucleic acid; EDTA, Ethylenediaminetetraacetic acid; LiH, Lithium Heparin; RNA, Ribonucleic acid.

(including MDD and episodes of mania and hypomania), repeating the GS:SFHS baseline assessment. Diagnostic criteria were based on the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV-TR). For participants who met full criteria for MDD, we assessed if any episode had a post-partum onset, and if criteria for melancholic or atypical MDD subtypes were met. The research version of the SCID was designed to allow assessors to systematically evaluate individuals against the key DSM-IV-TR criteria for unipolar depression and bipolar disorder. The SCID has good reliability, and it is considered the “gold standard” in determining clinical diagnoses and their accuracy¹⁵.

Participants also completed a series of short questionnaires that assessed resilience, psychological well-being and mild psychiatric problems, and personality, some of which were repeated after first being completed for GS:SFHS (see Table 2). The Brief Resilience Scale (BRS)¹⁶ is a six-item questionnaire used as a measure of psychological resilience, or the ability to ‘bounce back’ from stress. Participants were assessed for a life history of cannabis use using the Drug Use questionnaire developed for UK Biobank¹⁷. Those who used cannabis more than once were asked follow-up questions about the frequency and functional impact of their use. Three mood questionnaires were administered: the Mood Disorder Questionnaire (MDQ)¹⁸, which is a sensitive screen for bipolar spectrum disorders; the Quick Inventory of Depressive Symptomatology (QIDS)¹⁹, which is a 16-item inventory designed to assess the severity of depressive symptoms; and the Hospital Anxiety and Depression Scale (HADS)²⁰ anxiety subscale (seven items) was used to screen for symptoms of anxiety. In addition, the General Health Questionnaire (GHQ)²¹, a 28-item test, was used to assess general psychological well-being on four scales: somatic symptoms; anxiety; social dysfunction; and depression. We used a Likert scoring system for the GHQ to calculate scores for each scale separately, as well as a total score.

We administered two measures that assess core personality traits: we used the neuroticism and extraversion scales from the Eysenck Personality Questionnaire – Revised Short Form (EPQ-R)²², each of which has 12 items; and the International Personality Item Pool (IPIP), Five-Factor Personality Inventory²³, which is a 50-item questionnaire that assesses the following core personality traits: extraversion; agreeableness; conscientiousness; emotional stability (the reverse of neuroticism); and imagination/intellect (similar to openness). Additionally, the General Causality Orientations Scale²⁴, which consists of 12 vignettes describing scenarios to determine each person’s orientation of causality²⁵, was used to assess one’s inclination towards being motivated autonomously, externally, or passively.

Finally, we assessed early-life adversity (childhood or adolescent abuse or neglect) using the Childhood Trauma Questionnaire (CTQ)²⁶. This is a 28-item retrospective inventory that assesses three areas of abuse (emotional, physical, and sexual) and two areas of neglect (emotional and physical). The CTQ also includes a minimisation/denial scale that identifies potential underreporting of maltreatment. A mean score was

calculated for each measure by totalling the item responses, with appropriate reverse scoring (e.g., GHQ, BRS). Higher scores represent higher levels of psychological distress, personality trait, or childhood trauma, except for the BRS where higher scores indicate greater resilience. Scoring and interpretation of data were based on the administration manual of each test.

Cognitive testing

The cognitive tests that were applied will be used to assess the cognitive phenotype of depression, and whether genetic risk variants are related to impairment in specific cognitive domains. As with the questionnaire data, some cognitive tests were also repetitions of the GS:SFHS baseline assessment (Table 2). We included “cold” (emotion-independent) and “hot” (emotion-laden) cognitive tests, given growing evidence for distinct and interacting relationships between depression and measures of hot and cold cognition²⁷.

The cold cognitive test battery included validated and widely used cognitive tests that measure crystallised- and fluid-type cognitive tasks. The Mill Hill Vocabulary test²⁸ was used as a measure of acquired verbal intelligence, and is an estimate of ‘crystallised intelligence’ and peak cognitive ability. The Controlled Oral Word Association task²⁹ was used as a measure of phonemic verbal fluency using three letters (C, F, and L). The Digit Symbol Coding subtest from the Wechsler Adult Intelligence Scale-III³⁰ was used to measure information processing speed. A United Kingdom version of the Logical Memory subtest from the Wechsler Memory Scale-III³¹ was used to assess verbal memory and provided a measure of immediate and delayed verbal declarative recall. Total scores were created for each cognitive test by adding the number of correct responses; higher scores indicate better performance. The Matrix Reasoning test, a paper adaptation of the computerised version from the COGNITO psychometric examination³², was used to measure perceptual organisation and visuospatial logic. A summary of all mood, personality, and cognitive data and their completeness is shown in Table 4.

Three ‘hot’ cognitive measures were administered on a touch-screen laptop computer. The first task – the Bristol Emotion Recognition Test – consisted of 96 trials (16 of each emotion) that assessed recognition of six basic human facial emotions (happiness, anger, sadness, disgust, surprise, and fear), and biases in the attribution of emotion. The Affective Go/No-Go task comprised 120 trials that assessed behavioural inhibition using facial emotional stimuli (happy, sad, and neutral expressions). Finally, given evidence for impairments in reward processing in depression³³, we also included a modified version of the Cambridge Gambling Task, which assesses decision-making, risk-taking behaviour, and reward processing (30 trials). These three tests are described in detail elsewhere^{34,35}.

Brain magnetic resonance imaging

The neuroimaging protocol will allow analysis of potential risk factor relationships with brain structure and function, and test neurobiological mechanisms that are associated with depressive symptoms and resilience. In Aberdeen, participants were imaged on a 3T Philips Achieva TX-series MRI system (Philips

Healthcare, Best, Netherlands) with a 32 channel phased-array head coil and a back facing mirror (software version 5.1.7; gradients with maximum amplitude 80 mT/m and maximum slew rate 100 T/m/s). A projector and “Presentation” (Neurobehavioural Systems Inc, Berkeley, CA, USA) version 18.1 were used for the presentation of task-based fMRI. In Dundee, participants were scanned using a Siemens 3T Prisma-FIT (Siemens, Erlangen, Germany) with 20 channel head and neck phased array coil and a back facing mirror (Syngo E11, gradient with max amplitude 80 mT/m and maximum slew rate 200 T/m/s). A magnetic resonance compatible LCD screen was used to display fMRI (NordicNeuroLab, Bergen, Norway) task stimuli using “Presentation” version 20.0.

Both centres used the same protocol including structural and functional sequences. The structural sequences collected were

as follows: 3D T1-weighted fast gradient echo with magnetisation preparation; 3D T2-weighted fast spin echo; 3D Fluid Attenuation Inversion Recovery (FLAIR); Diffusion Tensor Imaging (DTI); and Susceptibility Weighted Imaging (SWI) or T2*-weighted gradient echo. The functional sequences comprised of two task-based fMRI tasks and a resting state fMRI sequence. The sequence parameters, as well as the order of acquisitions, are presented in [Table 5](#).

T1-weighted images of the brain were used to assess brain regional volumes, cortical thickness, gyrification index, voxel-based morphometry analysis, certain lesions such as lacunes, cortical and larger subcortical infarcts, and will also serve as the basis for co-registration with other sequences. A 3D T2-weighted sequence was used to detect lacunes, perivascular spaces, cortical and subcortical infarcts, and other morphological

Table 5. Magnetic resonance imaging (MRI) sequences and their characteristics acquired in Aberdeen and Dundee.

MR sequences ^a	Aberdeen	Dundee
3D T1-weighted (1)	160 sagittal slices TR = 8.2 ms* TE = 3.8 ms TI = 1031 ms FA = 8° FOV = 240 mm matrix size = 240 × 240 voxel size = 1.0 × 1.0 × 1.0 mm ³ acquisition time = 5 min 38 s	208 sagittal slices TR = 1740 ms TE = 2.62 ms TI = 900 ms FA = 8° FOV = 256 mm matrix size = 256 × 256 voxel size = 1.0 × 1.0 × 1.0 mm ³ acquisition time = 4 min 3 s
3D T2-weighted (8)	360 sagittal slices TR = 2500 ms TE = 314 ms FA = 90° FOV = 250 mm matrix size = 252 × 250 voxel size = 0.5 × 0.5 × 0.5 mm ³ acquisition time = 7 min 17 s	320 sagittal slices TR = 3200 ms TE = 408 ms FA = variable FOV = 256 mm matrix size = 256 × 256 voxel size = 0.5 × 0.5 × 0.5 mm ³ acquisition time = 7 min 10 s
3D FLAIR (5)	160 sagittal slices TR = 8000 ms TE = 349 ms TI = 2400 ms FOV = 240 mm matrix size = 240 × 238 voxel size 0.94 × 0.94 × 1.00 mm ³ acquisition time = 4 min 32 s	160 sagittal slices TR = 5000 ms TE = 386 ms TI = 1800 ms FOV = 256 mm matrix size = 256 × 256 voxel size 1.00 × 1.00 × 1.00 mm ³ acquisition time = 4 min 37 s
DTI (7)	60 axial slices TR = 7010 ms TE = 90 ms FA = 90° FOV = 220 mm matrix size = 96 × 94 voxel size = 2.3 × 2.3 × 2.3 mm ³ 64 non-collinear gradient directions (b = 1200 s/mm ²) eight unweighted (b = 0) acquisition time = 9 min 28s	60 axial slices TR = 7100 ms TE = 87 ms FA = 90° FOV = 220 mm matrix size = 96 × 94 voxel size = 2.3 × 2.3 × 2.3 mm ³ 64 non-collinear gradient directions (b = 1200 s/mm ²) eight unweighted (b = 0) acquisition time = 8 min 54 s

MR sequences ^a	Aberdeen	Dundee
T2* weighted or SWI (6)	130 axial slices TR = 31 TE = 7.2/13.4/19.6/25.8 ms FA = 17° FOV = 230 mm matrix size = 384 × 316 voxel size = 0.3 × 0.3 × 1 mm ³ acquisition time = 4 min 29 s	144 axial slices TR = 28 ms TE = 20 ms FA = 17° FOV = 230 mm matrix size = 320 × 160 voxel size = 0.4 × 0.4 × 1 mm ³ acquisition time = 5 min 17 s
fMRI sequences (2, 3, 4)	32 axial slices TR = 1560 ms TE = 26 ms FA = 70° FOV = 217 mm matrix size = 64 × 64 voxel size = 3.4 × 3.4 × 4.5 mm ³ total acquisition time = 24 min 13 s	32 axial slices TR = 1560 ms TE = 22 ms FA = 70° FOV = 217 mm matrix size = 64 × 64 voxel size = 3.4 × 3.4 × 4.5 mm ³ total acquisition time = 24 min 13 s

^aOrder of acquisition is presented in parentheses.

*8.2 ms × 240 phase encoding steps = 1968 ms.

Abbreviations: TR, repetition time; TE, echo time; TI, inversion time; FA, flip angle; FOV, field of view; DTI, diffusion weighted tensor images; FLAIR, fluid attenuation inversion recovery; SWI, susceptibility weighted imaging.

measurements, such as hippocampal subfield extraction. A 3D FLAIR was used to detect white matter hyperintensities. SWI data, for the determination of brain microbleeds, basal ganglia mineralisation, and cortical superficial siderosis, were acquired using a 3D multi-echo gradient-echo sequence in Aberdeen and a single-echo protocol in Dundee. Phase and magnitude data were saved for the calculation of T2* relaxation. All vascular lesions listed above are defined in the Standards for Reporting Vascular changes on Neuroimaging standards³⁶. All structural images were reviewed by a neuroradiologist for visual analysis of vascular changes and incidental findings. Whole-brain DTI were recorded to allow assessment of micro-structural integrity of white matter including fibre direction and structural connectivity. This protocol reflects established neuroimaging approaches as used in several large cohort studies of ageing and of cerebrovascular diseases^{37,38}.

There were two task-based fMRI sequences: an implicit emotional processing task (fearful versus neutral faces), and a modified version of an instrumental reward task with an additional choice value component. Both of these, as well as resting state fMRI, were acquired at 30 degrees away from the anterior commissure–posterior commissure (AC-PC), towards the coronal plane. The fearful faces from NimStim³⁹ facial stimuli set assessed emotional-limbic circuitry through a block fMRI design, and measures the brain's neural responses to the viewing of fearful faces in the absence of learning. In order to avoid a gender bias of the images, two versions of the tasks were used, counterbalanced across participants. The Reward task measured reward-related brain activity using an event-related fMRI design in a reinforcement learning

context. The resting state fMRI was used to investigate functional connectivity and brain networks.

Results

What are the key findings?

Here, we report findings for the complete data set including 1,188 participants. Table 6 shows some demographic similarities and differences between the current STRADL cohort and existing samples. More specifically, the median age of the STRADL sample was 62 years, which is older compared to both STRADL remote follow-up and wider GS:SFHS populations. Analysis of group differences for age showed that participants who are part of ACONF were generally older ($M = 62.32$; $SD = 1.55$) compared to the Walker cohort and wider GS:SFHS participants ($ts \geq 6.28$; $ps < .001$). However, age did not differ ($t = 1.83$; $p = .068$) between participants in the Walker cohort ($M = 59.48$; $SD = 6.76$) and those in GS:SFHS ($M = 58.28$; $SD = 12.63$). Gender distributions were comparable to existing data, with 59% being female, and our sample had higher levels of education (university-level education = 40%), compared to existing data. Furthermore, based on SCID interviews, a higher proportion of STRADL participants were diagnosed with a lifetime history of mood disorder (30.7%), compared to GS:SFHS (13.2%). Out of the total sample, 28.8% received a diagnosis of MDD, and a further 1.9% of bipolar disorder. Recurrent mood disorder was present in 72.7%, and melancholic features (56%) were dominant in the group. Of those with a diagnosis, 71% were female. Overall, however, the cohort was of good psychological health at the time of assessment, as indicated by mean scores on the GHQ, HADS, and QIDS (Table 7), which fell below the thresholds for the presence of

Table 6. Comparison of demographic data between STRADL face-to-face assessment, STRADL remote follow-up, and wider GS:SFHS baseline assessment.

Variable	STRADL face-to-face assessment (n = 1,188)	STRADL remote follow-up ¹² (n = 9,618)	GS:SFHS baseline assessment ⁸ (n = 21,525)
Median age (years)			
Male	62	54	47
Female	61	52	48
Gender (%) (female)	59	62	59
Employment (%) (those aged up to 75 years)			
Unemployed	3	4	2
Retired	32	18	15
Employed	65	71	63
Education (%)			
University-level degree	40	37	33
No qualification	25	7	5

Note. Data were extracted from time of each assessment point.

psychological distress. Cognitive scores across all measures were normally distributed. Psychological health, personality, and cognitive scores are presented in [Table 7](#).

Discussion

Strengths and limitations

STRADL data have been robustly collected on a wide range of key phenotypes that allow epidemiological study of depression and resilience in a population-based cohort. The MRI and detailed depression phenotyping protocol described here was cross-sectional; however, STRADL can provide longitudinal measures of cognition, personality, and psychological health. This is because many of the cognitive tests applied in STRADL are deliberately the same as those used at the GS:SFHS baseline assessment, as well as some personality and mood measures, as shown in [Table 2](#). The availability of repeated cognitive and questionnaire testing allows us to assess potential determinants of change in cognition and psychological health. Similarly, routine NHS data, and ACONF and Walker cohorts' early-life variables, are linked to STRADL data, providing further opportunities for longitudinal predictors on depression and resilience across the full life-course. However, a limitation of these data is that, as the present study is based on ages of different cohorts including people born in the 1950s onwards, early life variables are likely to have been influenced by cultural and societal norms of that time. For

example, people born in the 1950s and 1960s may not have considered housing circumstances such as crowdedness as an 'adverse experience', contrary to more contemporary perspectives. These possible confounding variables will be considered in downstream analyses, but it may be difficult to completely mitigate their effects.

A further limitation of this study includes possible selection bias as we tested only 21% of the total pool of eligible and invited participants ($n = 5,649$). As with many longitudinal population studies, participants in this cohort were more likely to be of good health, and come from more advantaged backgrounds such as higher education and better socioeconomic circumstances than the population in general – findings that are similar to GS:SFHS and STRADL remote follow-up cohort profiles^{8,12}, and UK Biobank¹⁷. Further, because our cohort was of good psychological health at the time of testing, it seems possible that we did not capture as many people with poorer psychological health as what might be available in the wider cohort. Notably, however, participants from a range of health and demographic backgrounds were represented in this group.

Ethical approval and consent

Ethical approval for the study was obtained from the Scotland A Research Ethics Committee (REC reference number

Table 7. Baseline questionnaire and cognitive measures (n = 1,188).

Variable	Maximum score	Mean (SD)	Range
Questionnaire data			
BRS	30	21.46 (4.89)	0–30
HADS – anxiety	21	4.15 (3.52)	0–20
QIDS – total	27	4.64 (3.65)	0–22
General Health Questionnaire			
GHQ – somatic symptoms	21	3.95 (3.23)	0–21
GHQ – anxiety	21	3.24 (3.32)	0–21
GHQ – social dysfunction	21	7.18 (2.01)	0–21
GHQ – depression	21	0.81 (2.39)	0–21
GHQ – total	84	15.15 (8.56)	2–75
EPQ-R – neuroticism	12	3.40 (3.19)	0–12
EPQ-R – extraversion	12	7.05 (3.80)	0–12
International Personality Item Pool			
IPIP – extraversion	50	31.17 (7.45)	11–50
IPIP – agreeableness	50	41.50 (5.34)	20–50
IPIP – conscientiousness	50	37.62 (6.04)	11–50
IPIP – emotional stability	50	33.58 (7.85)	10–50
IPIP – intellect	50	34.78 (5.58)	11–50
Childhood Trauma Questionnaire			
CTQ – physical abuse	25	6.11 (2.41)	0–25
CTQ – sexual abuse	25	6.04 (3.57)	0–25
CTQ – emotional abuse	25	7.00 (3.47)	0–25
CTQ – physical neglect	25	6.35 (2.36)	0–21
CTQ – emotional neglect	25	8.56 (4.21)	0–25
Cognitive data			
Logical Memory test – immediate	25	16.34 (3.63)	4–24
Logical Memory test – delayed	25	15.40 (3.91)	3–24
Logical Memory test – combined	50	31.74 (7.29)	9–48
Digit Symbol Coding	133	68.15 (15.24)	24–116
Verbal fluency test – combined	-	42.98 (12.02)	12–88
Mill Hill Vocabulary test	44	31.56 (4.09)	16–44
Matrix Reasoning test	15	8.26 (2.39)	1–15

Abbreviations: BRS, Brief Resilience Scale; CTQ, Childhood Trauma Questionnaire; EPQ-R, Eysenck Personality Questionnaire – Revised; GHQ, General Health Questionnaire; HADS, Hospital Anxiety and Depression Scale; IPIP, International Personality Item Pool; QIDS, Quick Inventory of Depressive Symptomology.

14/55/0039) and the local Research and Development offices. All participants provided written informed consent prior to the collection of any data or samples.

Data availability

All data underlying the results are available as part of the article and no additional source data are required.

A phenotype data dictionary is [available](#) and open access genome-wide association study summary statistics can be [downloaded](#). Non-identifiable information from the GS:SFHS cohort is available to researchers in the UK and to international collaborators through application to the Generation Scotland Access Committee (access@generationscotland.org) and through the Edinburgh Data Vault (<https://doi.org/10.7488/8f68f1ae-0329-4b73-b189-c7288ea844d7>). Generation Scotland

operates a managed data access process including an [online application form](#), and proposals are reviewed by the Generation Scotland Access Committee. The data and samples collected by the STRADL study have been incorporated in the main Generation Scotland dataset and governance process. Summary information to help researchers assess the feasibility and statistical power of a proposed project is available on request by contacting resources@generationscotland.org.

Acknowledgements

We are grateful to the GS:SFHS participants, and the ACONF and Walker subset of GS:SFHS who have taken part in the STRADL study. We also thank the families of the participants, and the wider GS:SFHS and STRADL project teams.

References

- Fernandez-Pujals AM, Adams MJ, Thomson P, *et al.*: **Epidemiology and heritability of major depressive disorder, stratified by age of onset, sex, and illness course in Generation Scotland: Scottish Family Health Study (GS:SFHS)**. *PLoS One*. 2015; **10**(11): e0142197. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gustavsson A, Svensson M, Jacobi F, *et al.*: **Cost of disorders of the brain in Europe 2010**. *Eur Neuropsychopharmacol*. 2011; **21**(10): 718–779. [PubMed Abstract](#) | [Publisher Full Text](#)
- Üstün TB, Ayuso-Mateos JL, Chatterji S, *et al.*: **Global burden of depressive disorders in the year 2000**. *Br J Psychiatry*. 2004; **184**: 386–392. [PubMed Abstract](#) | [Publisher Full Text](#)
- Major Depressive Disorder Working Group of the Psychiatric GWAS Consortium: **A mega-analysis of genome-wide association studies for major depressive disorder**. *Mol Psychiatry*. 2013; **18**(4): 497–511. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Von Korff M, Ormel J, Katon W, *et al.*: **Disability and depression among high utilizers of health care: a longitudinal analysis**. *Arch Gen Psychiatry*. 1992; **49**(2): 91–100. [PubMed Abstract](#) | [Publisher Full Text](#)
- Howard DM, Clarke TK, Adams MJ, *et al.*: **The stratification of major depressive disorder into genetic subgroups**. *bioRxiv*. 2017; 134601. [Publisher Full Text](#)
- Luthar SS, Cicchetti D, Becker B: **The construct of resilience: a critical evaluation and guidelines for future work**. *Child Dev*. 2000; **71**(3): 543–562. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Smith BH, Campbell A, Linksted P, *et al.*: **Cohort profile: Generation Scotland: Scottish Family Health Study (GS:SFHS). The study, its participants and their potential for genetic research on health and illness**. *Int J Epidemiol*. 2013; **42**(3): 689–700. [PubMed Abstract](#) | [Publisher Full Text](#)
- Kerr SM, Campbell A, Marten J, *et al.*: **Electronic health record and genome-wide genetic data in Generation Scotland participants**. *bioRxiv*. 2017; 154609: 85. [Publisher Full Text](#)
- Libby G, Smith A, McEwan NF, *et al.*: **The Walker Project: a longitudinal study of 48,000 children born 1952–1966 (aged 36–50 years in 2002) and their families**. *Paediatr Perinat Epidemiol*. 2004; **18**(4): 302–312. [PubMed Abstract](#) | [Publisher Full Text](#)
- Batty GD, Morton SM, Campbell D, *et al.*: **The Aberdeen Children of the 1950s cohort study: background, methods and follow-up information on a new resource for the study of life course and intergenerational influences on health**. *Paediatr Perinat Epidemiol*. 2004; **18**(3): 221–239. [PubMed Abstract](#) | [Publisher Full Text](#)
- Navrady L, Wolters M, MacIntyre D, *et al.*: **Cohort Profile: Stratifying Resilience and Depression Longitudinally (STRADL): a questionnaire follow-up of Generation Scotland: Scottish Family Health Study (GS:SFHS)**. *Int J Epidemiol*. 2017; **47**(1): 13–14g. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Meyer JS, Novak MA: **Minireview: hair cortisol: a novel biomarker of hypothalamic-pituitary-adrenocortical activity**. *Endocrinology*. 2012; **153**(9): 4120–4127. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- First M, Gibbon M, Spitzer R, *et al.*: **Structured Clinical Interview for DSM-IV-TR Axis I Disorders (Research Version)**. New York, NY: New York State Psychiatric Institute; 2002. [Reference Source](#)
- Lobbstaël J, Leurgans M, Arntz A: **Inter-rater reliability of the Structured Clinical Interview for DSM-IV Axis I disorders (SCID I) and Axis II disorders (SCID II)**. *Clin Psychol Psychother*. 2011; **18**(1): 75–79. [PubMed Abstract](#) | [Publisher Full Text](#)
- Smith BW, Dalen J, Wiggins K, *et al.*: **The brief resilience scale: assessing the ability to bounce back**. *Int J Behav Med*. 2008; **15**(3): 194–200. [PubMed Abstract](#) | [Publisher Full Text](#)
- Sudlow C, Gallacher J, Allen N, *et al.*: **UK Biobank: An open access resource for identifying the causes of a wide range of complex diseases of middle and old age**. *PLoS Med*. 2015; **12**(3): e1001779. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hirschfeld RM, Williams JB, Spitzer RL, *et al.*: **Development and validation of a screening instrument for bipolar spectrum disorder: the Mood Disorder Questionnaire**. *Am J Psychiatry*. 2000; **157**(11): 1873–1875. [PubMed Abstract](#) | [Publisher Full Text](#)
- Rush AJ, Giles DE, Schlessler MA, *et al.*: **The Inventory for Depressive Symptomatology (IDS): preliminary findings**. *Psychiatry Res*. 1986; **18**(1): 65–87. [PubMed Abstract](#) | [Publisher Full Text](#)
- Zigmond AS, Snaith RP: **The Hospital Anxiety and Depression Scale**. *Acta Psychiatr Scand*. 1983; **67**(6): 361–370. [PubMed Abstract](#) | [Publisher Full Text](#)
- Goldberg DP: **Manual of the General Health Questionnaire**. Windsor: NFER-Nelson; 1978. [Reference Source](#)
- Eysenck HJ, Eysenck SBG: **Manual of the Eysenck Personality Questionnaire: (EPQ-R adult)**. San Diego, Calif.: EdITS/Educational and Industrial Testing Service; 1994. [Reference Source](#)
- Goldberg LR: **The development of markers for the big-five factor structure**. *Psychol Assess*. 1992; **4**(1): 26–42. [Publisher Full Text](#)
- Deci EL, Ryan RM: **The General Causality Orientations Scale: self-determination in personality**. *J Res Pers*. 1985; **19**(2): 109–134. [Publisher Full Text](#)
- Ryan RM, Deci EL: **Self-determination theory: Basic psychological needs in motivation, development, and wellness**. Guilford Publications. 2017. [Reference Source](#)
- Bernstein DP, Fink L: **Childhood Trauma Questionnaire: a retrospective**

- self-report.** Harcourt Brace & Company. 1998.
[Reference Source](#)
27. Roiser JP, Sahakian BJ: **Hot and cold cognition in depression.** *CNS Spectr.* 2013; **18**(3): 139–149.
[PubMed Abstract](#) | [Publisher Full Text](#)
 28. Raven JC: **Raven's progressive matrices and vocabulary scales.** Oxford, England: Oxford Psychologists Press Ltd. 1989.
 29. Benton A, Hamsher K, Sivan A: **Multilingual aphasia examination.** 3rd ed. San Antonio, TX: Psychological Corporation; 1994.
[Reference Source](#)
 30. Wechsler D: **Wechsler Adult Intelligence Scale-Third Edition (WAIS-III).** San Antonio: Harcourt Assessment Inc. 1997.
[Reference Source](#)
 31. Wechsler D: **Wechsler Memory Scale-Third Edition (WMS-III).** San Antonio: Harcourt Assessment Inc. 1997.
 32. Ritchie K, Roquefeuil G de, Ritchie C, *et al.*: **COGNITO: computerized assessment of information processing.** *J Psychol Psychother.* 2014; **4**(2): 1000136.
[Publisher Full Text](#)
 33. Whitton AE, Treadway MT, Pizzagalli DA: **Reward processing dysfunction in major depression, bipolar disorder and schizophrenia.** *Curr Opin Psychiatry.* 2015; **28**(1): 7–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 34. Attwood AS, Easey KE, Dalili MN, *et al.*: **State anxiety and emotional face recognition in healthy volunteers.** *R Soc Open Sci.* 2017; **4**(5): 160855.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 35. Bland AR, Roiser JP, Mehta MA, *et al.*: **EMOTICOM: A Neuropsychological Test Battery to Evaluate Emotion, Motivation, Impulsivity, and Social Cognition.** *Front Behav Neurosci.* 2016; **10**: 25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 36. Wardlaw JM, Smith EE, Biessels GJ, *et al.*: **Neuroimaging standards for research into small vessel disease and its contribution to ageing and neurodegeneration.** *Lancet Neurol.* 2013; **12**(8): 822–838.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 37. Wardlaw JM, Bastin ME, Valdés Hernández MC, *et al.*: **Brain aging, cognition in youth and old age and vascular disease in the Lothian Birth Cohort 1936: rationale, design and methodology of the imaging protocol.** *Int J Stroke.* 2011; **6**(6): 547–559.
[PubMed Abstract](#) | [Publisher Full Text](#)
 38. Valdés Hernández Mdel C, Armitage PA, Thrippleton MJ, *et al.*: **Rationale, design and methodology of the image analysis protocol for studies of patients with cerebral small vessel disease and mild stroke.** *Brain Behav.* 2015; **5**(12): e00415.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 39. Tottenham N, Tanaka JW, Leon AC, *et al.*: **The NimStim set of facial expressions: judgments from untrained research participants.** *Psychiatry Res.* 2009; **168**(3): 242–249.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:  

Version 2

Reviewer Report 23 July 2021

<https://doi.org/10.21956/wellcomeopenres.18737.r45009>

© 2021 Lawlor D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Debbie A. Lawlor 

Medical Research Council Integrative Epidemiology Unit, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

The authors have responded to my comments and I consider this a useful cohort profile

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Epidemiology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 07 April 2020

<https://doi.org/10.21956/wellcomeopenres.17011.r38261>

© 2020 Lavretsky H. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Helen Lavretsky

Department of Psychiatry, University of California, Los Angeles (UCLA), Los Angeles, CA, USA

The reported study, STRatifying Resilience and Depression Longitudinally (STRADL) is a population-based study built on the Generation Scotland: Scottish Family Health Study (GS:SFHS) resource. The aim of STRADL is to subtype major depressive disorder (MDD) on the basis of its etiology,

using detailed clinical, cognitive, and brain imaging assessments. Recruitment has been completed and we consented and tested 1,188 participants.

The manuscript describes the methods of data collection in STRADL included: socio-economic and lifestyle variables; physical measures; questionnaire data that assesses resilience, early-life adversity, personality, psychological health, and lifetime history of mood disorder; laboratory samples; cognitive tests; and brain magnetic resonance imaging. Some of the questionnaire and cognitive data were first assessed at the GS:SFHS baseline assessment between 2006-2011, thus providing longitudinal measures of depression and resilience. The presented results are purely description of the sample. Overall, the study has a lot of potential in providing links to the biomarkers and predictors of depression and resilience, including personality traits, early trauma exposure, and genetic vulnerability. However, it would help to provide clarification about study initial hypotheses and the analytical plan.

The title of the study implies that stratification can be made between depression and resilience longitudinally. Nothing in the description explains what will be done about subtyping MDD (only unipolar and bipolar depression diagnoses are mentioned), or what would it take to stratify by resilience and depression.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

No

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Resilience, late life depression, biomarkers

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 08 Jun 2021

Tina Habota, University of Aberdeen, UK

Response:

Thank you for your comments above. We have amended our manuscript to include further clarification about the study hypotheses and focus of stratification of MDD under the section "Why was the study set up?":

"Stratification of MDD will be based on several variables to address its underlying causal and clinical heterogeneity, including: age of onset of MDD; single episode or recurrent depression; obstetric trauma; and developmental factors such as childhood maltreatment, early socioeconomic adversity, and stressful life events. Our key initial predictions are that depression can be stratified on the basis of age of onset into early-onset forms that show a stronger phenotypic and genetic relationship with schizophrenia and other severe mental disorders, and later onsets that show stronger associations with cardiovascular disease and dementia".

Competing Interests: No competing interests were disclosed.

Reviewer Report 24 February 2020

<https://doi.org/10.21956/wellcomeopenres.17011.r37159>

© 2020 Lawlor D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Debbie A. Lawlor 

Medical Research Council Integrative Epidemiology Unit, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

This profile provides details of the second (face-to-face) assessment of the STRADL (STratifying Resilience and Depression Longitudinally) study. STRADL is itself a sub-study of the Generation Scotland – Scottish Family Health Study (GS-SFHS). The GS-SFHS is a truly impressive resource for the scientific community and this profile is a very valuable paper updating the community on some detailed and novel assessments within a subgroup of that resource.

My main comments are related to trying to improve the value to the wider scientific community and are in the context of someone who does not work predominantly in mental health, but does work on cohort studies and recognises that for those studies we work on we often forget how much is 'inside our heads' and may be assumed to much of readers who are nothing like as familiar as 'we are' (in this case the authors of this paper who know GS-SFHS and STRADL in considerable detail). There were places where I felt a bit lost and could have done with more detail that publication on WOR supports.

Please note I felt obliged to tick 'not approved' as that was the only option that states I am suggesting revisions that I feel have to be undertaken. I think this is an important paper that should be indexed and widely read. I do think that some of my key concerns to require that

changes are made to the paper.

Specifically:

1. Some brief description of the selection criteria for GS-SFHS and its initial aims would be really useful. Did it cover the whole of Scotland? Was it family/house based inclusion criteria? What were the initial inclusion and exclusion criteria? How many in that cohort as a whole have been lost to follow-up? A paragraph providing this background context (rather than relying on the reader knowing GS-SFHS or going and reading that cohort profile) would be really useful.
2. Similarly, some detail on STRADL – i.e. the first (remote) assessment I think is essential – what were the eligibility criteria at the start of STRADL? What was the response to that first remote STRADL assessment? What data were collected in that remote STRADL? And were the aims of that remote STRADL assessment the same as those for this more detailed face-to-face follow-up? This to me is essential both to understand the value of this recent face-to-face assessment and to understand some of the demographic differences presented and the potential for selection bias in STRADL as a whole and also this face-to-face assessment
3. Related to point 2 above I think that Figure 1 needs to be amended so that STRADL remote is also included – this way a clearer picture of the relation between that and the current face-to-face is clear.
4. The overall response of the 5649 invited to take part in the face-to-face presented here is 21% (N = 1188) that fact is never mentioned and it really has to be – it should be stated in the abstract; clearly shown in Figure 1 and stated in the text related to that figure and needs to be discussed in more detail in the discussion limitations (again it is not mentioned as a limitation in that section).
5. It is stated that STRADL recruitment and data collection ended in May 2019 but I could not find any statement showing when it started. As this is a study about longitudinal analyses that to me seems essential to report. As noted below, I found the chronology and what was truly available for repeat longitudinal assessment (i.e. of change in an exposure or an outcome) or for prospective analyse (i.e. of an exposure at one time point with an outcome at a later time point) confusing. I think a figure that showed the chronology on a time scale of the baseline GS-SFHS (2006-2011), STRADL-remote (initial formation of STRADL; 2015? month of start and finish) and STRADL face-to-face (?? To May 2019) – on a horizontal date metre with an indication of median and full range of time between these – would be really useful. It would also be useful to include in that for the Walker and the ACONF the baseline collection of data in those (e.g. could be an arrow back to 1956 for ACONF) and also the date ranges of different linkages (I think for some hospital records may be only back to 90s and may differ between 'psychiatric' diagnoses and 'physical' diagnoses).
6. I found the content, including column headings of Table 1 confusing, in particular as the content often seemed to be contradicted by the text that referred to that table.

Specifically:

1. The chronology and repeats seemed erroneous. How can the GS-SFHS be a repeat when that is the baseline measure .. yet the heading calls it 'repeat'. Similarly, the

STRADL remote is repeat but that was also before the current face-to-face. As above no specific date is given for STRADL face-to-face it simply says 'current', which isn't quite true if recruitment and data collection finished in May 2019 – what is needed is a start and end date for each of these column headings, and I would order chronologically and take out 'repeat' from any of the column headings as it is clear if something has been measured more than once from the entries (or at least should be).

2. In the text it states that cognitive function measures assessed at STRADL face-to-face are the same as those done at GS-SFHS in several places (including in the discussion as a strength that these were "deliberately the same"), but in Table 1 under cognitive measurements there is no overlap in the ticks for measures from GS-SFHS and STRADL face-to-face. If some were deliberately repeated there should be a tick for both GS-SFHS and STRADL face-to-face.
3. The text also implies that DNA was taken again at the face-to-face but there is only a tick for that at GS-SFHS & the same applies for some of the plasma biomarkers.
7. Could the authors clarify when the demographic data presented in Table 4 were taken from. I assume that they were all taken from GS-SFHS baseline assessment, which would be the appropriate comparison (i.e. suggesting that those who agree to participate in the initial STRADL study (the remote one) were older at that baseline than those who did not respond/take part in that and those who took part in STRADL face-to-face). Wherever they are taken from it is important this is stated in the text describing the table and title / footnote to the table; at the moment I could not see it anywhere.
8. It would be useful in Table 4 and/or in text to have some indication of demographic differences in the two birth cohorts (Walker and ACONF) at each of the assessments. This seems important to me as the older age in STRADL face-to-face (and possibly STRADL remote but not possible to tell without further information on that) could be due to oversampling from those birth cohorts. The Walker cohort were born 1952 to 1966 and the ACONF in 1956 – if that makes them generally older than the GS-SFHS to me that has different implications for selection than if they are similarly aged to the rest of GS-SFHS.
9. For a cohort profile – which in my book is about telling the international academic community what is available and what it might be used for and where there are limitations and caution required – I was very surprised at the lack of any discussion of limitation. To me the following all need detailed discussion:
 1. Sample size. The aim of STRADL and in particular the face-to-face follow-up described here is said to be to "examine the interaction between genetic and environmental factors that increase risk and occurrence of different MDD subtypes, and assess common and distinct mechanisms and clinical trajectories of MDD phenotypes. Additionally, STRADL aims to assess individual resilience, or the ability to adapt positively and 'avoid' psychopathology despite exposure to known risk factors such as stress, early-life adversity, and family history" 'Interactions' and 'subtypes' (i.e. with the two together subgroups within subgroups) require very large sample sizes to give meaningful results and it is not clear to me how this will be possible with just over 1000 participants. Some examples of clinically meaningful results for stratified medicine / precision prevention

(the focus of the introduction and justification for the STRADL sub study) would be really valuable.

2. Selection bias As noted previously the 21% of those eligible and invited to the STRADL face-to-face study that is the focus of this profile seems to be ignored. There have been a very large number of publications recently that have described issues related to this -including when exploring genetic associations (and particularly in relation to gene-environment interactions) and a clear discussion of that in relation to this study is necessary.
3. Availability of data for the aims Related to both points above it is unclear from what is written how much missing data will impact on the research that this study aims to particularly address. For example, of the 1188 recruited 544 (~47%) will have prospectively collected birth / early life data (i.e. those in the Walker and ACONF birth cohorts) but the remaining 53% will not. And whilst Table 2 & 3 provide some reassurance they do not cover the early life data and even with small amounts of missing data (as low as 2-3%) for single variables once one tries to combine these the numbers with complete data for all models/analyses can be considerably smaller than 100%. Related to this could the authors clarify where data in Tables 2 & 3 come from. Are they from one particular assessment (of the three that the 1188 are likely to have participated in) or from any – e.g. could weight and height be available on 99.9% because they are coded 'available' if they had at least one measure from any of the three assessments). Again this information is essential to understanding the potential for repeat measures and prospective analyses that the study aims to address.
4. Relevance of data for contemporary populations I appreciate that all of our studies are somewhat restricted in terms of their target populations, but as this study (based on ages of different groups) were born in the 1950s and 60s predominantly some discussion of how the data collected for early life exposures will have been influenced by norms and cultures at the time, including how people now in their 50s and 60s might remember or be willing to retrospectively report what would now-adays be considered 'adversity' but may not have been at the time. How will possible residual confounding be dealt with for analyses of the early life exposures when these may not have been collected. For example, smoking during pregnancy was not recorded in ACONF)

As well as discussing these limitations I think one actual analysis for this paper relevant to the aims of STRADL would be valuable to give the authors and readers a sense of what is truly possible. For example, one could try to explore whether there is an interaction between childhood adversity and adult alcohol consumption and one of the face-to-face MRI or cognitive function continuously measured outcomes. That would provide some evidence of power in relation to precision of any effect estimates an idea of missing data when trying to ensure all potential confounders are accounted for and would mean having to discuss any results in the context of possible selection bias.

Minor comments

- The following sentences in the abstract "Some of the questionnaire and cognitive data were first assessed at the GS:SFHS baseline assessment between 2006-2011, **thus** providing longitudinal measures of depression and resilience. **Similarly**, routine NHS data and early-

life variables are linked to STRADL data, further providing opportunities for longitudinal analysis" (my highlight) read a little oddly to me. As the date for the STRADL study describe here is not prevented it is not clear how / why these provide longitudinal measures. Also it feels like depression and resilience are being used as interchangeable with cognitive function to some extent in the first of these two sentences.

- In the introduction "Thus, it is important for treatment to shift from the current "trial and error" approach, towards **personalised** and preventative forms of treatment **for individuals** with markedly different disease mechanisms." As we cannot make inference to individuals or personalise (i.e. to an individual) treatments from cohort or RCT studies I would change this to the terms that are more widely accepted – 'stratified medicine' and 'precision prevention'
- It is worth clarifying (for a wide international audience) that Aberdeen is in Grampian (as you do for Dundee and Tayside).
- Under the summary of biological samples, you mention repeat genetic analyses. Might be worth changing to repeat DNA Methylation as genetic variants will not change over time. Also I would clarify in that section when you refer to 'epigenomic status' you mean DNA methylation and transcription (you refer to RNA later). Also, worth clarifying that DNA methylation will be in white blood cells only (I think that is correct – if not then state with other tissues). From the text I understood that DNA would be extracted from samples again at the STRADLE face-to-face but there is not tick in Table 1 to confirm that?
- In Table 1 it appears as if on all participants DNA from blood and saliva are available whereas that is not the case. I would suggest where you first mention extracting DNA from saliva when blood cells are not available you give the N & % in parentheses where that had to be done & in Table 1 just have one line 'Extracted DNA' and a footnote stating that for 97.5% that was from white blood cells with the remaining 2.5% being from saliva

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Not applicable

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Epidemiology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to state that I do not consider it to be of an acceptable scientific standard, for reasons outlined above.

Author Response 08 Jun 2021

Tina Habota, University of Aberdeen, UK

- Some brief description of the selection criteria for GS-SFHS and its initial aims would be really useful. Did it cover the whole of Scotland? Was it family/house-based inclusion criteria? What were the initial inclusion and exclusion criteria? How many in that cohort as a whole have been lost to follow-up? A paragraph providing this background context (rather than relying on the reader knowing GS-SFHS or going and reading that cohort profile) would be really useful.

Response:

The premise and aims of GS:SFHS were described briefly, but we have included further details that describes the selection process/criteria addressing the reviewer's comment:

"GS:SFHS included participants aged 35-65 years who were identified at random from collaborating medical practices across Scotland, with some family members further afield. Initially, only Glasgow and Tayside areas were involved, but the study was extended in 2010 to include Ayrshire, Arran and Northeast Scotland, with the age range also broadened (to 18-65 years). Participants were included if: they met the age criteria; had capacity to give informed consent; and could identify at least one first-degree relative who would also participate. Follow-up of participants was done through the NHS Scotland Community Health Index (CHI): 7% of the original cohort could not be matched; no participants withdrew; and ~1,200 had died".

- Similarly, some detail on STRADL – i.e. the first (remote) assessment I think is essential – what were the eligibility criteria at the start of STRADL? What was the response to that first remote STRADL assessment? What data were collected in that remote STRADL? And were the aims of that remote STRADL assessment the same as those for this more detailed face-to-face follow-up? This to me is essential both to understand the value of this recent face-to-face assessment and to understand some of the demographic differences presented and the potential for selection bias in STRADL as a whole and also this face-to-face assessment

Response:

Thank you for this comment and we agree that this could be clarified further in the manuscript. We have clarified the differences between the two STRADL waves by elaborating and clarifying the aims of STRADL remote follow-up:

"The first wave of STRADL included depression-focused follow-up assessment of GS:SFHS, which involved remote questionnaires that specifically assessed aspects of psychological resilience, coping style, and response to psychological distress; study protocol and cohort characteristics are

described elsewhere [12]".

We have also included a brief description of STRADL remote study:

"Those who participated in GS:SFHS were invited to take part in the STRADL remote follow-up based on the following eligibility criteria if: they had given consent for re-contact; were living in Scotland; and had a CHI number. 9,618 GS:SFHS responded to the invitation, including 2,460 unrelated individuals and 2,460 families (7,158 individuals) of between 2 and 18 family members".

- Related to point 2 above I think that Figure 1 needs to be amended so that STRADL remote is also included – this way a clearer picture of the relation between that and the current face-to-face is clear.

Response:

We agree with the reviewers and have added this additional information to Figure.

- The overall response of the 5649 invited to take part in the face-to-face presented here is 21% (N = 1188) that fact is never mentioned and it really has to be – it should be stated in the abstract; clearly shown in Figure 1 and stated in the text related to that figure and needs to be discussed in more detail in the discussion limitations (again it is not mentioned as a limitation in that section).

Response:

We agree that this was not sufficiently clear in our original submission and we thank the reviewer for drawing our attention to this issue. We have now clarified the participation rate which states:

"...we consented and tested 1,188 (72.2%) of positive respondents across Aberdeen (n = 582) and Dundee (n = 606) sites. This meant that we tested 21% of the n = 5,649 who were initially invited to participate".

We did not include this statistic in Figure 1 because we only report sample sizes and not proportions, but the sample size is consistent with the quoted proportion.

- It is stated that STRADL recruitment and data collection ended in May 2019 but I could not find any statement showing when it started. As this is a study about longitudinal analyses that to me seems essential to report. As noted below, I found the chronology and what was truly available for repeat longitudinal assessment (i.e. of change in an exposure or an outcome) or for prospective analyse (i.e. of an exposure at one time point with an outcome at a later time point) confusing. I think a figure that showed the chronology on a time scale of the baseline GS-SFHS (2006-2011), STRADL-remote (initial formation of STRADL; 2015? month of start and finish) and STRADL face-to-face (?? To May 2019) – on a horizontal date metre with an indication of median and full range of time between these – would be really useful. It would also be useful to include in that for the Walker and the ACONF the baseline collection of data in those (e.g. could be an arrow back to 1956 for ACONF) and also the date ranges of different linkages (I think for some hospital records may be only back to 90s and may differ between 'psychiatric' diagnoses and 'physical' diagnoses).

Response:

Thank you for your comments above. We have now included an additional table that shows all existing data sets that are linked to the STRADL face-to-face study. We have now introduced Table 1 and state: "Table 1 shows the linkage between the current study and existing datasets".

- I found the content, including column headings of Table 1 confusing, in particular as the content often seemed to be contradicted by the text that referred to that table. Specifically:
-
- The chronology and repeats seemed erroneous. How can the GS-SFHS be a repeat when that is the baseline measure .. yet the heading calls it 'repeat'. Similarly, the STRADL remote is repeat but that was also before the current face-to-face. As above no specific date is given for STRADL face-to-face it simply says 'current', which isn't quite true if recruitment and data collection finished in May 2019 – what is needed is a start and end date for each of these column headings, and I would order chronologically and take out 'repeat' from any of the column headings as it is clear if something has been measured more than once from the entries (or at least should be).

Response:

Table 2 (previously Table 1) lists all of the variables collected in STRADL face to face assessment compared to STRADL remote follow-up and GS, and specifically we have indicated which variables are "new" (column 1). Then we have also indicated which were "repeated" from STRADL remote study and GS baseline assessment (columns 2 and 3).

- In the text it states that cognitive function measures assessed at STRADL face-to-face are the same as those done at GS-SFHS in several places (including in the discussion as a strength that these were "deliberately the same"), but in Table 1 under cognitive measurements there is no overlap in the ticks for measures from GS-SFHS and STRADL face-to-face. If some were deliberately repeated there should be a tick for both GS-SFHS and STRADL face-to-face.

Response:

Table 2 (previously Table 1) lists all of the variables collected in STRADL face to face assessment, which included eight cognitive tasks. Four cognitive tests (Logical Memory; Digit Symbol Coding; Verbal Fluency; Mill Hill Vocabulary) were repeated from GS baseline assessment; and four (Matrix Reasoning; Bristol Emotion Recognition test; Affective Go/No-Go task; Cambridge Gambling task) were new tests only administered in face-to-face assessment.

- The text also implies that DNA was taken again at the face-to-face but there is only a tick for that at GS-SFHS & the same applies for some of the plasma biomarkers.

Response:

DNA was obtained from whole blood at the STRADL face-to-face assessment. Table 2 shows all of the variables that were a part of the STRADL face to face assessments – therefore we have now clearly indicated that this is a repeat from the GS baseline assessment. Table 4 shows the proportion of DNA obtained via blood sample and saliva sampling.

- Could the authors clarify when the demographic data presented in Table 4 were taken from. I assume that they were all taken from GS-SFHS baseline assessment, which would be the appropriate comparison (i.e. suggesting that those who agree to participate in the initial STRADL study (the remote one) were older at that baseline than those who did not respond/take part in that and those who took part in STRADL face-to-face). Wherever they are taken from it is important this is stated in the text describing the table and title / footnote to the table; at the moment I could not see it anywhere.

Response:

The demographic data from Table 6 (previously Table 5) was taken at time of assessment at each of the study points (GS, STRADL remote follow-up, and STRADL face-to-face assessment). We have noted this in a footnote to the table on page 23:

“Data were extracted from time of each assessment point”.

- It would be useful in Table 4 and/or in text to have some indication of demographic differences in the two birth cohorts (Walker and ACONF) at each of the assessments. This seems important to me as the older age in STRADL face-to-face (and possibly STRADL remote but not possible to tell without further information on that) could be due to oversampling from those birth cohorts. The Walker cohort were born 1952 to 1966 and the ACONF in 1956 – if that makes them generally older than the GS-SFHS to me that has different implications for selection than if they are similarly aged to the rest of GS:SFHS.

Response:

Under the section “what are the key findings”, we have elaborated on the age differences between groups at STRADL face-to-face assessment, as below:

“Analysis of group differences for age showed that participants who are part of ACONF are generally older ($M = 62.32$; $SD = 1.55$) compared to participants in the Walker cohort and wider GS:SFHS participants ($t = 6.28$; $p < .001$). However, age did not differ ($t = 1.83$; $p = .068$) between participants in the Walker cohort ($M = 59.48$; $SD = 6.76$) and those in GS:SFHS ($M = 58.28$; $SD = 12.63$)”.

- For a cohort profile – which in my book is about telling the international academic community what is available and what it might be used for and where there are limitations and caution required – I was very surprised at the lack of any discussion of limitation. To me the following all need detailed discussion:
 - Sample size. The aim of STRADL and in particular the face-to-face follow-up described here is said to be to “*examine the interaction between genetic and environmental factors that increase risk and occurrence of different MDD subtypes, and assess common and distinct mechanisms and clinical trajectories of MDD phenotypes. Additionally, STRADL aims to assess individual resilience, or the ability to adapt positively and ‘avoid’ psychopathology despite exposure to known risk factors such as stress, early-life adversity, and family history*” ‘Interactions’ and ‘subtypes’ (i.e. with the two together subgroups within subgroups) require very large sample sizes to give meaningful results and it is

not clear to me how this will be possible with just over 1000 participants. Some examples of clinically meaningful results for stratified medicine / precision prevention (the focus of the introduction and justification for the STRADL sub study) would be really valuable.

Response:

No sample size calculation was provided for the recruiting participants in the current investigation. A larger sample of individuals was originally planned, which was reduced in light of UK Biobank's stated plans to recruit upwards of 10,000 participants for neuroimaging. The study plan was changed to reduce the number of participants included for imaging purposes – but representing at least a 4-fold increase on the current average sample size of a depression imaging cohort study. We estimated that this sample would have sufficient power to detect previously reported true differences from studies of approximately 50 cases and 50 controls, and detect differences explaining approximately 50% of the variance previously reported by earlier studies. However, this power calculation was not conducted prospectively and we have not therefore included this in the text of the manuscript, although we agree that this is a limitation.

- Selection bias As noted previously the 21% of those eligible and invited to the STRADL face-to-face study that is the focus of this profile seems to be ignored. There have been a very large number of publications recently that have described issues related to this -including when exploring genetic associations (and particularly in relation to gene-environment interactions) and a clear discussion of that in relation to this study is necessary.

Response:

Under the “Strengths and Limitations” section, we have elaborated on the above point and state:

“A limitation of this study includes possible selection bias as we tested only 21% of the total pool of eligible and invited participants (n = 5,649). As with many longitudinal population studies, participants in this cohort were more likely to be of good health, and come from more advantaged backgrounds such as higher education and better socioeconomic circumstances than the population in general – findings that are similar to GS:SFHS and STRADL remote follow-up cohort profiles [8, 12], and UK Biobank [17]. Further, because our cohort was of good psychological health at the time of testing, it seems possible that we did not capture as many people with poorer psychological health as what might be available in the wider cohort”.

- Availability of data for the aims Related to both points above it is unclear from what is written how much missing data will impact on the research that this study aims to particularly address. For example, of the 1188 recruited 544 (~47%) will have prospectively collected birth / early life data (i.e. those in the Walker and ACONF) birth cohorts) but the remaining 53% will not. And whilst Table 2 & 3 provide some reassurance they do not cover the early life data and even with small amounts of missing data (as low as 2-3%) for single variables once one tries to combine these the numbers with complete data for all models/analyses can be considerably smaller than 100%. Related to this could the authors clarify where data in Tables 2 & 3 come from. Are they from one particular assessment (of the three that the 1188 are likely to have participated in) or from any – e.g. could weight and height be available on 99.9%

because they are coded 'available' if they had at least one measure from any of the three assessments). Again this information is essential to understanding the potential for repeat measures and prospective analyses that the study aims to address.

Response:

Linkage to early-life variables will be carried out through data available from the ACONF and Walker birth cohorts, but also through routine NHS data. Whilst we have provided the proportion of available data in Tables 3 and 4 for the current study, we do not have the missing values for the NHS datasets as these are continuously updated and missing values differ greatly on an item-by-item basis. The impact of any missing data will, however, need to be addressed in individual analyses and papers.

- Relevance of data for contemporary populations I appreciate that all of our studies are somewhat restricted in terms of their target populations, but as this study (based on ages of different groups) were born in the 1950s and 60s predominantly some discussion of how the data collected for early life exposures will have been influenced by norms and cultures at the time, including how people now in their 50s and 60s might remember or be willing to retrospectively report what would nowadays be considered 'adversity' but may not have been at the time. How will possible residual confounding be dealt with for analyses of the early life exposures when these may not have been collected. For example, smoking during pregnancy was not recorded in ACONF).

Response:

Thank you for your comments above, with which we agree – particularly on the issue of retrospective reporting of adversity. We have included a discussion under "Strengths and Limitations" that addresses this point:

"As the present study is based on ages of different cohorts including people born in the 1950s onwards, early life variables are likely to have been influenced by cultural and societal norms of that time. For example, people born in the 1950s and 1960s may not have considered housing circumstances such as crowdedness as an 'adverse experience', contrary to more contemporary perspectives. These possible confounding variables will be considered in downstream analyses, but it may be difficult to completely mitigate their effects".

- As well as discussing these limitations I think one actual analysis for this paper relevant to the aims of STRADL would be valuable to give the authors and readers a sense of what is truly possible. For example, one could try to explore whether there is an interaction between childhood adversity and adult alcohol consumption and one of the face-to-face MRI or cognitive function continuously measured outcomes. That would provide some evidence of power in relation to precision of any effect estimates an idea of missing data when trying to ensure all potential confounders are accounted for and would mean having to discuss any results in the context of possible selection bias.

Response:

We agree that this would be an interesting analysis, but we would not wish to add it to this paper. Our aims for this paper were to focus on describing the baseline cohort results without analyses that address specific study aims – these are currently being undertaken in

a series of separate analyses and papers. The additional methods required to address this suggestion would also complicate and extend a paper that is already fairly complex.

Minor comments

- The following sentences in the abstract “Some of the questionnaire and cognitive data were first assessed at the GS:SFHS baseline assessment between 2006-2011, **thus** providing longitudinal measures of depression and resilience. **Similarly**, routine NHS data and early-life variables are linked to STRADL data, further providing opportunities for longitudinal analysis” (my highlight) read a little oddly to me. As the date for the STRADL study describe here is not presented it is not clear how / why these provide longitudinal measures. Also it feels like depression and resilience are being used as interchangeable with cognitive function to some extent in the first of these two sentences.

Response:

Thank you for raising these points, which we now clarify in the Abstract:

“Some of the questionnaire and cognitive data were first assessed at the GS:SFHS baseline assessment between 2006-2011, thus providing longitudinal measures relevant to the study of depression, psychological resilience, and cognition. In addition, routinely collected historic NHS data and early-life variables are linked to STRADL data, further providing opportunities for longitudinal analysis. Recruitment has been completed and we consented and tested 1,188 participants.”

- In the introduction “Thus, it is important for treatment to shift from the current “trial and error” approach, towards **personalised** and preventative forms of treatment **for individuals** with markedly different disease mechanisms.” As we cannot make inference to individuals or personalise (i.e. to an individual) treatments from cohort or RCT studies I would change this to the terms that are more widely accepted – ‘stratified medicine’ and ‘precision prevention’

Response:

We agree and have changed the text as follows:

“Thus, it is important for treatment to shift from the current “trial and error” approach, towards precision prevention and stratified medicine based on markedly different disease mechanisms”.

- It is worth clarifying (for a wide international audience) that Aberdeen is in Grampian (as you do for Dundee and Tayside).

Response:

We have now clarified this in the manuscript.

- Under the summary of biological samples, you mention repeat genetic analyses. Might be worth changing to repeat DNA Methylation as genetic variants will not change over time. Also I would clarify in that section when you refer to ‘epigenomic status’ you mean DNA rillmethylation and transcription (you refer to RNA later). Also, worth clarifying that DNA methylation will be in white blood cells only (I think that is correct – if not then state which other tissues). From the text I understood that DNA

would be extracted from samples again at the STRADLE face-to-face but there is no tick in Table 1 to confirm that?

Response:

Thank you for these comments. We have clarified this point and state:

“We collected laboratory blood samples for genetic and additional genomic analyses, including the study of DNA methylation, transcription (RNA) and protein expression.”

DNA samples were first collected in GS baseline assessment thus Table 2 (previously Table 1) has a tick to indicate this, and there is no tick under the STRADL face to face assessment because it is not a “new” variable.

- In Table 1 it appears as if on all participants DNA from blood and saliva are available whereas that is not the case. I would suggest where you first mention extracting DNA from saliva when blood cells are not available you give the N & % in parentheses where that had to be done & in Table 1 just have one line ‘Extracted DNA’ and a footnote stating that for 97.5% that was from white blood cells with the remaining 2.5% being from saliva

Response:

Thank you for this comment. Table 2 (previously Table 1) shows all of the variables that were collected overall in the study, whereas Table 4 (previously Table 3) shows the specific proportion of available data – we have amended Table 4 so that it states the total proportion of DNA available (99.2), but we include a footnote that states:

“96.5% of DNA was extracted via blood sample and 2.7% via saliva sampling.”

Competing Interests: No competing interests were disclosed.