

The feasibility and effects of eye movement training for visual field loss after stroke: a mixed methods study

Christine Hazelton¹ , Alex Pollock², Diane Dixon³, Anne Taylor⁴, Bridget Davis⁵, Glyn Walsh⁶, Marian C Brady⁷

British Journal of Occupational Therapy
2021, Vol. 84(5) 278–288

© The Author(s) 2020



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/0308022620936052

journals.sagepub.com/home/bjot



Abstract

Introduction: Visual field loss affects around 20% of stroke survivors, reducing quality of life. Eye movement training is a promising rehabilitation method, and several different interventions are used by occupational therapists. This study aimed to explore the feasibility and effects of four eye movement training interventions for stroke survivors with visual field loss.

Method: A mixed methods study – quantitative n-of-1 with qualitative interviews. The participants were 11 home-dwelling stroke survivors with visual field loss. The interventions used were MyHappyNeuron, NVT, Rainbow Readers and VISIOcoach, delivered in a randomised order. Visual search, reading speed, activities of daily living and quality of life were assessed three times before intervention use, then immediately after each intervention; these were analysed visually. A final semi-structured interview was then analysed using framework methods.

Results: Evidence of effect was divergent. Quantitatively there was no measured effect, but qualitatively participants reported benefits in visual skills, daily life skills and emotions, which varied by intervention. Median training time was 3–4 hours (range 0.5–6.5) for NVT, Rainbow Readers and MyHappyNeuron, and 9.5 hours (range 2.3–16.8) for VISIOcoach.

Conclusion: Eye movement training interventions were feasible for stroke survivors at home. Qualitative evidence suggests that variations in the eye movements trained and delivery modality underlie variations in perceived effect.

Keywords

Stroke, visual field loss, rehabilitation, eye movement training, compensation, mixed methods

Received: 9 April 2020; accepted: 1 June 2020

Introduction

Stroke causes lasting visual impairment in around one fifth of stroke survivors (Ali et al., 2013). Visual field loss, produced by damage to the visual pathway, typically causes loss of one half of the visual field on the same side in both eyes, effectively making the person blind to one side of space. Subsequent eye movement changes, involving smaller, repetitive movements, increase the time taken to view an entire scene (Zihl, 1995). The result is limited mobility and navigation (de Haan et al., 2015a), causing social isolation (Hazelton et al., 2019a). Reading and driving difficulties impact on leisure, occupations and return to work (Hepworth and Rowe, 2016), and psychological consequences include fear and lack of confidence (Hazelton et al., 2019a). Stroke survivors with visual field loss report limitations in activities of daily living (ADL) (Warren, 2009), poorer quality of life (QoL) (Chen et al., 2009), with restricted engagement in rehabilitation and poorer functional outcome (Ali et al., 2013). Occupational therapists have a key role in providing rehabilitation interventions for visual field loss in stroke survivors (Warren, 1993).

Identifying effective interventions for visual loss is a top ten research priority for stroke survivors, carers and clinicians (Pollock et al., 2012). Systematic review evidence suggests that training compensatory eye movements towards the side of visual field loss may improve QoL (Pollock et al., 2019). A range of approaches to training compensatory eye movements exist, including:

¹Nursing, Midwifery and Allied Health Professions (NMAHP) Research Unit, Glasgow Caledonian University, Glasgow, UK

²NMAHP Research Unit, Glasgow Caledonian University, Glasgow, UK

³Institute of Applied Health Sciences, University of Aberdeen, Aberdeen, UK

⁴Faculty of Health Sciences and Sport, University of Stirling, Stirling, UK

⁵Department of Nursing and Community Health, Glasgow Caledonian University, Glasgow, UK

⁶Vision Sciences, Glasgow Caledonian University, Glasgow, UK

⁷NMAHP Research Unit, Glasgow Caledonian University, Glasgow, UK

Corresponding author:

Christine Hazelton, Nursing Midwifery and Allied Health Professions Research Unit, A600, Govan Mbeki Building, Glasgow Caledonian University, Cowcaddens Road, Glasgow G4 0BA, UK.

Email: christine.hazelton@gcu.ac.uk

(a) scanning training, which teaches broad horizontal eye (and head) movements to enable a full view of a scene; (b) search training, which stimulates a stroke survivor's natural pattern of eye movements; and (c) reading training, which aims to improve the small left–right eye movements required to read across a line of text. Overall, eye movement training may lead to larger, less repetitive and more frequent eye movements (Passamonti et al., 2009; Schuett, 2009), improve visual search (Aimola et al., 2014) and mobility (Aimola et al., 2014; Hayes et al., 2012). However, high quality adequately powered studies providing evidence of benefits to stroke survivors' activities and participation are needed (Pollock et al., 2011).

Eye movement training is utilised by occupational therapists and other professions (Rowe, 2013). Our earlier work identified therapist-supported specialist devices, paper-based tools, online training and computer programs in use to train eye movements (Hazelton et al., 2019b). These interventions vary in relation to the training aims, area of visual field covered, inclusion of head movements, cost and number, duration and frequency of training sessions (Hazelton et al., 2019b). We lack studies which directly compare these different eye movement training devices, despite their routine clinical use. Evidence on effectiveness and feasibility is required to inform clinical decisions and support optimal rehabilitation for stroke survivors with visual field loss.

The aim of this study was therefore to use mixed methods to explore and compare the feasibility of use and the effects of eye movement training interventions. We aimed to directly compare four eye movement interventions, selected for maximum variation, from those known to be used in clinical practice: MyHappyNeuron, NVT (NeuroVision Technology), Rainbow Readers, and VISIOcoach.

Methods

Our mixed methods approach combined a quantitative n-of-1 design with qualitative semi-structured interviews (Borglin, 2015). In an n-of-1 design each participant acts as their own control: quantitative outcome measures were conducted three times before the introduction of any interventions, spaced 2 weeks apart, in order to establish a baseline (Supplementary material 1). Participants were then provided with each of the four eye movement training interventions, delivered sequentially in a randomised order. After completing training with an intervention, the outcomes measures were repeated; these repeated quantitative measures were used to explore within-person effects associated with the use of each intervention. Qualitative interviews were conducted after all interventions had been used, to enable participants to compare their experiences with each. Baseline data were collected at a clinic-based assessment; all other study activities took place in participants' own homes. Ethical approval was granted by the relevant ethical committees.

Recruitment

Participants were recruited from the waiting lists of two Scottish vision rehabilitation centres. Stroke-specialist low vision rehabilitation officers identified potential participants from their caseloads, by reviewing their notes in order to apply the inclusion criteria. The inclusion criteria were: clinical diagnosis of stroke at least 6 months earlier, hemianopic visual field loss, age 18 years or over, medically stable, living in the community and no prior community visual training. Visual field loss was assessed using confrontation, with any pattern of binocular field loss in the same vertical hemifield accepted. Exclusion criteria were: unable to provide informed consent, non-stroke visual impairment and involvement in another rehabilitation study. Low vision rehabilitation officers explained the study to eligible clients and provided large print/audio information and consent forms; participation was based on return of written consent forms.

Interventions

We tested MyHappyNeuron, NVT, Rainbow Readers and VISIOcoach. Table 1 provides a description of each intervention, derived in a previous study (Hazelton et al., 2019b) (for full TIDieR description see Supplementary material 2).

Data collection

Baseline demographic data, including data on their stroke and vision were collected by an experienced optometrist (Supplementary material 1).

Quantitative measures collected at each of the seven timepoints were: visual search speed and accuracy – computer-based conjunction visual search (Treisman and Gelade, 1980); reading speed – International Reading Speed Texts (Trauzettel-Klosinski et al., 2012); self-reported abilities in ADL – Veteran's Affairs Low Vision Visual Function Questionnaire (Stelmack et al., 2004); self-reported quality of life – Visual Core Measure 1 (Lamoreaux et al., 2008).

Participants used diaries to record adherence to the advised intervention regimen for each intervention. For NVT the diary was completed by the low vision rehabilitation officer delivering training.

Qualitative semi-structured interviews were conducted with each participant after using all four interventions, to explore and compare how well they could use each of the interventions, and any possible effects each had on them. A topic guide (Supplementary material 3) was used to ensure core questions were included and minimise interviewer biases; this was refined to explore unexpected responses and emergent themes (Rubin and Rubin, 2005).

Data analysis

Outcome measure data for each participant were plotted in line graphs, and analysed using visual inspection

Table 1. Intervention description.

Name	Modality	Description	Training aims		Overall	Schedule	Total dosage (mins)	Delivery
			Eye movements	Advised use				
MyHappy Neuron	Computer – online ^a	'Brain-training' with many detailed, engaging exercises, onscreen instructions and feedback on performance. The difficulty level progresses with improvement. Access: happy-neuron.com	Not specified	Cognitive skills	8 × 35 mins, over 2 weeks	280	Self-delivered. Initial instruction and demonstration session with researcher, who setup a direct desktop link to website. Written guidance given.	
NVT scanning device (NeuroVision Technology)	Specialist equipment	A 'lightbox' around 1.5 m wide, with two rows of coloured lights. Used by a therapist to assess and then train a systematic scan, typically from unseen to seen visual field. Lightbox training followed by 'dynamic' training in real life environments and tasks. Access: via trained rehabilitation officers	Broad horizontal scan, also moving the head	Mobility and activities of daily living	8 × 45 mins, over 5 weeks ^b	360	Low vision rehabilitation worker delivered (face-to-face). Equipment brought by therapist for each session.	
Rainbow Readers	Paper-based	A booklet of exercises, made up of simple symbols arranged in broken lines across the page. The aim is to search for, and circle, a specific symbol within a line of other (distractor) symbols. Exercises increase in difficulty. Access: nadia.northway@gcu.ac.uk	Left-to-right scan	Reading	10 mins/day, 6 days/week, for 2 weeks	120	Self-delivered. Booklet given at initial instruction and demonstration session by researcher. Written guidance given.	
VISIOcoach	Computer software ^a	On-screen exercises, requiring the user to find the indicated number/letter, hidden in an irregular pattern of numbers/letters on a white background. There are three difficulty levels. Access: visiocoach.de/	Visual search, with no set pattern/instructions given	Mobility and activities of daily living	2 × 30 mins/day, 5 days/week, for 2 weeks	600	Self-delivered. Initial instruction and demonstration session with researcher, who uploaded software and created desktop icon for access. Written guidance given.	

^aParticipant's own computer and internet access were used if available, if not a laptop with dongle for internet access was provided.

^bNVT was provided over a longer timeframe in order to fit with low vision rehabilitation officer's existing caseload and schedule.




Figure 1. Participant involvement flowchart.

(Smith, 2012), with input from a methodological specialist (DD), and following a systematic process with clear guidelines to minimise bias and inconsistency (Kazdin, 2011). This involved assessment of the trend or stability of each phase and comparison of the intervention and baseline phases. A process of considered judgement (Kazdin, 2011) recorded the intervention effect as either possible effect (beneficial/detrimental), no effect or unclear. Diary data were analysed using descriptive statistics.

Transcribed interview audio files were analysed inductively with a five-stage thematic framework method (Ritchie and Spencer, 2002), using NVivo v10 software. This involved: (a) analysts developed a ‘feel’ for the data, by reading the transcripts; (b) line-by-line coding of the first four interviews to create a thematic framework of higher-order ‘categories’; (c) applying the framework to remaining transcripts; (d) creating charts, with intervention data displayed in rows and categories forming the columns; and (e) interpreting the data, looking for emergent connections, patterns or ‘themes’.

The first two analysis stages involved two authors (CH and BD), with one (CH) conducting the subsequent stages, with input throughout from a topic expert (AP) and methodological expert (AT). A framework of 52 codes in nine categories was developed, which covered intervention effect, intervention feasibility and opinions of the study (Supplementary material 4).

Results

Seventeen eligible stroke survivors were identified and 12 were enrolled, but one dropped out before intervention delivery due to changed family circumstances (Figure 1). Demographic data are given in Table 2. Service changes meant low vision rehabilitation officers did not deliver NVT to the final two participants. Quantitative outcome measure completion rate was 10/11 for ADL and QoL scales, with one participant not completing each due to fatigue, 7/11 for reading speed, due to language-based reading problems, and 9/11 for visual search, due to poor comprehension of the test.

The Results section presents the feasibility data first followed by those on effects, to reflect the study aim, and quantitative data followed by qualitative data, to reflect the order of their use within the study.

Feasibility

Quantitative results. The training times recorded are given in Table 3.

Qualitative data. When asked about their ability to use interventions, participants typically identified the difficulties they experienced, relating to ease of use (the most important factor), ease of initial set-up and level of carer support required.

MyHappyNeuron: Most reported difficulties with set-up or use were related to MyHappyNeuron. A

Table 2. Baseline data for participants.

Characteristic	Participants (n = 11)
Age (years)	Median: 56 Min-max range: 42-80
Gender	6 Women (54.5%)
Visual field loss	
Side	
Right	6 (54.5%)
Left	5 (45.5%)
Size	
Half	5 (45.5%)
Quarter	5 (45.5%)
Other (sectoral)	1 (9.0%)
Visual neglect	n = 1 (9.0%)
Time since stroke (months)	Median: 9 Min-max range: 6-24
Stroke severity (Stroke Impact Scale) ^a	Median (min-max range)
Physical	78.9 (35.2-100)
Communication	82.1 (42.6-100)
Memory	53.6 (32.1-100)
Emotion	58.3 (25-97.2)
Participation/role	40.6 (15.6-87.5)
Living situation	
Alone	3 (27.3%)
With spouse	4 (36.4%)
With larger family group	4 (36.4%)
Access to computer with internet	
Yes	10 (90.9%)
No	1 (9.1%)

^aStroke impact scores: higher score indicates better ability, or less severe impact.

Table 3. Intervention feasibility: training time recorded.

	Number accessing intervention (n = 11)	Total time recorded	
		Minutes median	% of recommended time median (range)
MyHappyNeuron	8 (73%)	260	92.9% (10.7-114.3%)
NVT	9 (82%)	180 ^a	50.0% (16.7-77.8%)
Rainbow Readers	10 (91%)	190	158.3% (29.2-325.0)
VISIOcoach	11 (100%)	570 ^b	95.0% (23.3-168.3%)

^aBased on five participants: low vision rehabilitation officers did not complete all diaries.

^bBased on 10 participants: one forgot to complete his diary.

clear split was apparent between those who found MyHappyNeuron usable and enjoyable and found it too complicated. The latter group reported struggling to read and understand the instructions, remember the 'sequences' within exercises and comprehend their aim. This was exacerbated by the task speed and short time limits for exercises.

we tried it and we tried it, and . . . it was too quick, if it gave you more time. . . . (Participant 10, right half-field loss)

Support was essential for several participants, but family and carers were not always able to provide the aid needed for MyHappyNeuron use.

NVT was set up and delivered by low vision rehabilitation officers, therefore as participants were less involved in accessing the intervention, they spoke relatively little about its feasibility. The ease of training varied from 'very simple'

(participant 10) to 'really difficult at times' (participant 5). For some, the 'dynamic' part of training, involving practising scanning in daily life tasks was limited by stroke-related lower-limb, pain or balance problems. Dynamic exercises were then tailored by low vision rehabilitation officers to the participant's specific limitations.

Rainbow Readers was seen as the easiest intervention to set up, as the process of opening the book and selecting the page to start on was quick and straightforward:

That's not too taxing, not any set-up (Participant 5, right half-field loss)

Even though the difficulty level increased, it was always achievable, and few required support. The lack of feedback provided meant that some participants sought confirmation they were doing exercises correctly. Carer feedback thus appeared to provide emotional support, encouraging their continued training:

I'd like to do that for myself. But the thing is, the problem being is I don't know if I'm going wrong [...] because if I was doing anything wrong I liked somebody to say 'That's alright' (Participant 1, left quadrant loss)

VISIOcoach was perceived as easy to set up compared to MyHappyNeuron, the other computer-based tool:

I just switched the computer on and it was there ... there wasn't lots of sequences to go through (Participant 1, left quadrant loss)

Participants reported that VISIOcoach's very clear exercise aims and the relatively straightforward tasks made it achievable. Support was often needed initially, but as training progressed individuals reported learning how to use the intervention independently.

I would sit at the kitchen table and do it ... and it would only be if I got stuck that I would say [to his wife to help him] (Participant 6, left half-field loss)

Effects

Quantitative results. The effect of the four interventions across the group is summarised in Table 4. Visual plots were created for each outcome measured for each participant; an example using data from the visual search test is given in Supplementary material 5. There was no evidence of effect arising from the use of MyHappyNeuron or VISIOcoach, with very limited evidence that NVT had a potentially beneficial effect on ADL and QoL and Rainbow Redears potentially improving reading speed.

Qualitative results. Participants described that using each of the four interventions had different effects:

Well there was a different outcome from ... (each) obviously (Participant 7, right quadrant loss)

Reported effects were in four interacting areas: visual skills, cognitive skills, emotions and tasks and activities. Intervention use produced effects in all four categories, but the type and number in each varied (Supplementary material 6). Participants noted that the effects on emotions and tasks and activities were most important to them.

MyHappyNeuron was reported to have little effect on visual skills. Instead, participants felt it provided mental stimulation and improved general knowledge, memory, problem-solving and overall mental agility.

Improved confidence was a key emotional effect of using all four interventions, including MyHappyNeuron. This was both general self-confidence or confidence related to specific tasks. It was founded on successful exercise completion.

(MyHappyNeuron) was building up my confidence 'I can do this, I can do this!' (Participant 5, right half-field loss)

MyHappyNeuron's similarity to pre-stroke activities meant people felt they were returning to normality. Conversely, those who struggled with MyHappyNeuron saw this as a personal 'failure' (participant 9), a reminder of post-stroke disabilities, and their confidence was reduced.

Yes I think I ... [laughs] how stupid am I? I do think it's made me aware of what I can't do, not what I can do and that's sadly so (Participant 3, left quadrant loss)

NVT led to participants reporting the use of much broader scanning movements. Some identified their use of both head and eye movements (taught by low vision rehabilitation officers) as much more natural. This combined movement was a key step in improving their scanning:

Definitely NVT (was most effective) because I hadn't realised until then that if I turned my head to the left [...] but I hadn't realised that ... It hadn't clicked, hadn't clicked. (Participant 1, left quadrant loss)

In contrast to the other interventions, the task and activity effects of NVT included many based outdoors. The key improvement reported was the ability to navigate and walk safely and independently, and participants stated they were not experiencing the same number of collisions, trips or falls. Importantly, participants described reduced fear when considering walking outside:

it helped me with going outside and not walking into things (Participant 9, right half-field loss)

It was appreciable that the perceived improvement in confidence gained from NVT was greater than with other interventions. Through performing outdoor tasks supported by their low vision rehabilitation officer, participants developed awareness of the number and scope of their own abilities and experienced a real sense of achievement in tasks that related directly to their daily life.

It helped to make me more confident ... It let me get out and see what I could do, what I could see [...] and I realised I can do this myself without anybody (Participant 10, right half-field loss)

Rainbow Readers was perceived by participants to lead to enhanced visual attention, plus the use of broader eye movements. The effect on daily activities was not always clear, but participants suggested it improved reading:

Table 4. Summary of outcome measure results for each intervention.

Intervention	Visual search ^a				Reading speed ^a	Activities of daily living ^a	Quality of life ^a	Summary
	Accuracy	Response time	Response time	Response time				
MyHappy Neuron	No changes	Changes identified in two participants: one with improvement in RT (Pt 1) and one with worsening of RT (Pt 3)	No changes	No changes	Worsening of reading speed in one participant (Pt 5)	Changes identified in two participants: in one ADL scores deteriorated (Pt 1) and in the other scores improved (Pt 7)	Changes identified in one participant (Pt 7), suggesting improvement in QoL	No evidence of effect
NVT	No changes	Change identified in one participant, which suggested worsening performance (Pt 4)	Worsening of reading speed in one participant (Pt 5)	Worsening of reading speed in one participant (Pt 5)	Improvement noted for three participants. In two this affected the majority of subscales (Pts 1 and 3), in one it primarily affected the mobility subscale (Pt 7). Decrease in reading score suggested a worsening of reading ability in one participant (Pt 4)	Improvement noted in four participants (Pts 1, 3, 4 and 5); magnitude of change 0.3–0.8	Limited evidence suggests possible beneficial effect on ADL and QoL	
Rainbow Readers	No changes	One change of RT suggestive of worsening of performance (Pt 11)	Improved reading speed in four participants (Pts 1, 3, 4 and 12) in 3 this was at clinically significant levels (Pts 1, 3 and 12)	Improved reading speed in four participants (Pts 1, 3, 4 and 12) in 3 this was at clinically significant levels (Pts 1, 3 and 12)	Changes identified in one participant (Pt 5), with worsening in mobility score	Change was noted in one participant (Pt 5), suggesting deterioration in QoL	Limited evidence suggests possible beneficial effect on reading speed	
VISIO coach	No changes	Large RT change of 3.2 seconds in one participant suggestive of improvement (Pt 1)	Notable change of score in three participants: two with improved speed (Pts 3, 12), one with worsening speed (Pt 7)	Notable change of score in three participants: two with improved speed (Pts 3, 12), one with worsening speed (Pt 7)	Changes identified in two participants: in one the scores suggested improved ADL performance (Pt 3) and in the other suggested deterioration (Pt 11)	Changes identified in three participants: in two the score improved (Pts 2, 7) and in the other it worsened (Pt 11)	No evidence of effect	

Pt: participant; RT: response time; ADL: activities of daily living; QoL: quality of life.

^aMeasurement tools were: visual search – computer-based conjunction visual search (Treisman and Gelade, 1980); reading – International Reading Speed Texts (Trauzettel-Klosinski et al., 2012); activities of daily living – Veteran's Affairs Low Vision Visual Function Questionnaire (Stelmack et al., 2004); quality of life – Visual Core Measure 1 (Lamoreaux et al., 2008).

Rainbow Readers kind of teaches you to read properly along the lines (Participant 1, left quadrant loss)

VISIOcoach was reported to encourage users to consider the entire visual scene:

I found it good in as much as it wasn't just teaching you to scan from left to right, it was teaching you to scan the whole page (Participant 5, right half-field loss)

Uniquely, two participants described that VISIOcoach's lack of instructions on how to find the targets allowed them to determine their own way of making better eye movements. For one the 'natural thing to do' (participant 5) was to scan the task quarter by quarter. Another discovered that vertical, rather than horizontal eye movements worked better:

Something just clicked in my brain that, you know, do it that way and it's better. ... I think it probably did make me quicker as well by doing it up and down. (Participant 12, left quadrant loss)

Participants felt VISIOcoach improved their computer use: some reported a return to using laptops and tablets, and in activities including gaming, surfing the internet and sending emails, which improved their contact with friends.

Participants were asked if there was a preferred intervention, or combination: they clearly appreciated the use of a range of tools. It was felt the affects of each was different, but 'complementary' (participant 12). Learning different patterns of eye (and head) movements meant participants felt they had a range of methods they could potentially use to approach a visual task:

It just expanded your visual possibilities ... it was giving you the flexibility to do it different ways, which again I found good (Participant 5, left half-field loss)

Discussion

This study explored the feasibility and effect of eye movement training interventions for 11 home-based stroke survivors with visual field loss. The effect of each intervention on four outcome measures was explored in a series of experimental n-of-1 studies, with diaries used to gather usage data, and qualitative interviews conducted.

Feasibility

This study has demonstrated that eye movement training was generally feasible for home-dwelling stroke survivors with visual field loss. The median training time was 3–4 hours for NVT, Rainbow Readers and

MyHappyNeuron, and around 9.5 hours for VISIOcoach. For the self-delivered interventions, participants typically achieved at least 93% of the recommended training time. For NVT, the low vision rehabilitation officers reported tailoring the training time to individual's needs and abilities, resulting in an average training time of 50% of that expected. Few studies have explored this aspect of eye movement training: Aimola et al. (2014) found that only three (of 28) participants achieved the training goal of 35 hours of therapy over 5 weeks, (equivalent to around 1 hour training per day), in an unsupervised computer-based training. However, in our study 11 participants using computer-based VISIOcoach (which also recommended 1 hour training per day) achieved 95% of this training time, suggesting acceptable adherence to the intervention.

The qualitative data from this study help explain some of the variation seen in the quantitative results. Stroke-related impairments, primarily in cognition but also language, motor or visual skill, appeared to contribute to barriers to access, limiting training time or necessitating carer support. Low vision rehabilitation officers delivering NVT reported modifying training sessions according to participants' ability, to ensure it was accessible. The level of intervention challenge relative to participant ability, and the ability either to reduce intervention difficulty or provide stroke survivors with support, appears critical to the feasibility of intervention access.

Effect

Evidence from the quantitative and qualitative methods was essentially divergent, with no, or very limited, measured effect of eye movement training on visual search, reading, ADL and QoL, but participants reported a range of perceived beneficial effects from all four interventions. The reported effects were on visual and cognitive skills, functional abilities, including using the internet, walking and using public transport, and emotions. Increased confidence was a key effect, particularly in response to NVT training. Thus, the lack of quantitative evidence of effect cannot be interpreted as evidence of no effect, as this is contradicted by the qualitative data.

Previous quantitative research suggests VISIOcoach increases visual search results (Roth et al., 2009), and NVT can improve visual search, peripheral vision and mobility (Hayes et al., 2012). Possible explanations for the difference between our results and previous research may relate to this study design: there was a relatively short training period of 2 weeks, so the full training effect may not have been established and therefore was not captured by quantitative outcomes. A key issue is that our qualitative results may have overstated training impact; with participants aware they are receiving 'additional' intervention they may have sought to report positive experiences (Patton, 2002). This is a central factor in contradictory qualitative and quantitative results

(Campbell et al., 2003). However, the value of mixed methods research lies partly in the fact that the two methodologies explore different aspects of the same phenomenon (De Lisle, 2011). For participants, one of the key effects was emotional, improving their sense of confidence, motivation and hope, which have not been considered in detail in prior work. No quantitative outcome measure that directly addressed emotional changes was included; the interviews allowed the participants to report their experience more freely, enabling identification of this broader range of potential effects.

Our qualitative data helped explore and explain several variations noted between interventions. One important factor for participants was the use of head movements in combination with eye movements. This combination was taught by NVT, with Rainbow Readers and VISIOcoach minimising use of the head and aiming to improve eye movements only. The use of scanning head movements has been criticised as slower and potentially detrimental to functional ability (Kerkhoff et al., 1992); however, our findings contradict this. Participants clearly felt their use was beneficial, and for some was key in achieving improvements in ADL (such as walking outside and navigating safely). A clear variation was also identified in the task and activity improvements participants reported. These suggested that improvement in functional abilities reflected the intervention modality, with limited transfer to non-trained tasks reported; paper-based Rainbow Readers was perceived to improve reading, and computerised VISIOcoach training was perceived to improve computing and reading skills, with no reported impact on wider mobility or ADL skills, even though it aimed to do so (Table 1 and Supplementary material 6). Within the literature there are two opposing theories relating to transfer: (a) that eye movements learnt in one task will carry over into other activities, as the new eye movement patterns will be internalised and applied automatically in new tasks (Pambakian et al., 2004); and (b) that scanning training is non-transferrable, and therapy must include training of eye movements during the activities of interest, such as mobility and ADL (Hayes et al., 2012; Pommerenke et al., 1989). These findings give some support to the latter theory, with only NVT, which included task-specific training, showing improvements in these skills within the timeframe of this study.

Data from this study suggest that participants favoured using a combination of interventions. Participants felt the effects and visual skills acquired differed between interventions (Supplementary material 6) and were complementary, giving greater versatility in the skills available to use. Studies of stroke survivors with visual field loss have suggested that different eye movements are used for different tasks (Hardiess et al., 2010), so training that encourages a broad visual scanning movement is unhelpful in tasks that require visual search (de Haan et al., 2015b). In addition, reading-specific training which encourages small left-to-right horizontal eye movements does not improve visual

search and vice versa (Schuett et al., 2012). Our results therefore support the need for training across several strategies (using a range of tools) to give participants the flexibility of visual skills required to tackle the broad variety of activities seen in daily life, and the different visual demands they present to a stroke survivor with field loss.

Strengths and weaknesses

A key strength of this study was that our n-of-1 method allowed the inclusion of four interventions, which represented key differences (modality, mechanisms, area of visual field covered and training schedule) in current clinical practice and enabled the identification of variations in their feasibility and hours of training completed. Our qualitative data (rarely considered in this field) has allowed stroke survivor experiences and perceptions of each intervention to be explored, providing valuable insight into the reasons for variation in feasibility and range of effects of each training method. Our comprehensive semi-structured interviews and rigorous analysis increased the insights provided by the participants' accounts. Combining and comparing the quantitative and qualitative results has allowed further interpretation of study findings, especially in relation to the conflicting data of the intervention effect.

A number of limitations of this study must be noted. The degree to which our findings are applicable to the wider population is uncertain given our small sample size, use of convenience sampling strategy and recruitment from the vision-specific centres, which may only see the more severe cases. The number of participants using each intervention was also not equal, which may limit the ability to draw consistent conclusions. The use of the n-of-1 study design means that placebo effects cannot be accounted for, and factors other than the interventions may have created the effects reported. For practicality, training duration for each intervention lasted around 2 weeks; although eye movement training has been shown to have an effect after just one session (Jacquin-Courtois et al., 2013), this short duration may have reduced training effect. Training dosage was recorded by participants and therapists – in some cases diaries were not fully completed and the method is open to errors of recording or recall so data may not be accurate. Some participants had memory difficulties (Table 1), which may have affected their ability to recount their experiences, a difficulty inherent in interviewing individuals post-stroke. As noted previously, the qualitative aspect is open to bias on the part of the interviewer, analyst and participant, and this may have acted to over-state the perceived effect of the interventions.

Implications for occupational therapy

Several eye movement training interventions were feasible, and could be provided to home-dwelling stroke survivors with visual field loss. Individuals' abilities

(especially cognition and language) may impact on the accessibility, so occupational therapists should consider these and the ability to tailor the level of challenge an intervention presents when developing a care plan. Our study suggests that interventions may have different, but complementary effects, and that these appear modality specific. Consequently, clinicians should consider providing a range of interventions, using diverse modalities (paper based, computer based and specialist equipment) to maximise the functional impact for stroke survivors. Furthermore, the choice of intervention should be related to an individual's occupational performance needs and goals. Our qualitative evidence suggests that for a reading goal, Rainbow Readers would be the first choice of intervention, for a goal of returning to computer use VISIOcoach or MyHappyNeuron may be appropriate, with NVT most relevant for someone with navigation and community access goals. Given the value placed on learning new head movements alongside eye movements, clinicians should aim to teach both aspects to improve compensatory abilities.

Our study suggests further investigation of the effectiveness of NVT, Rainbow Readers and VISIOcoach eye movement training interventions (potentially in combination) may be warranted. Given the disagreement surrounding the eye movements required to compensate for visual field loss, the rationale for targeting specific eye/head movements should be stated alongside exploration of how variations in eye movement may impact on functional ability. Researchers should also consider the use of qualitative methods and their choice of quantitative outcome measures used to capture the effectiveness of treatment. Stroke survivors stated that tasks and activities plus the emotional effects of training were most important to them; however, our results suggest that current vision-based ADL and QoL scales may be insensitive to change and may inadequately capture the emotional impact of eye movement training.

Conclusion

Eye movement training interventions were essentially feasible for home-based rehabilitation. Feasibility was limited in tools which used fast, complex exercises, as they posed too high a linguistic or cognitive challenge. Quantitative data gave no clear evidence of intervention effect from, but there were perceived effects on visual skills, cognitive skills, emotions and functional abilities, which varied by intervention. Variations were linked to intervention modality, with a lack transfer to untrained tasks suggested, and to the training of head movements in combination with eye movements, which was perceived as beneficial.

Key findings

- Eye movement training is feasible for home-dwelling stroke survivors, and may improve ADL.

- Effects may vary by intervention and modality: using several interventions may maximise effects.

What the study has added

This study suggests eye movement training is feasible, with a range of perceived benefits. By exploring the factors affecting intervention success, it enables occupational therapists to choose interventions relevant to individuals' rehabilitation needs.

Acknowledgements

The author(s) would like to thank the designers of all included scanning training tools for providing free access during this study. They wish to note that MyHappyNeuron is designed for a general population, and a version specifically for healthcare use (HappyNeuron Pro) is also available. We would also like to thank the low vision centres and rehabilitation officers involved in this study.

Research ethics

Ethical approval was obtained from the NHS West of Scotland Research Ethics Service (reference 13/WS/0171).

Declaration of conflicting interests

The author(s) confirm that there are no conflicts of interest.

Funding

This study was funded by the Stroke Association (UK) by way of a Junior Research and Training Fellowship held by the lead author (TSA JRTF 2011/02). MCB, AP and the NMAHP Research Unit are funded by the Scottish Government Health and Social Care Directorates. The views expressed here are those of the authors and not necessarily those of the funders.


Consent

All participants provided written informed consent to participate in the study.

Contributorship

All authors contributed to the design of the project and data analysis plan. Data was collected by CH, quantitative analysis was conducted by CH, AP, DD and qualitative analysis by CH, BD, AT, and AP with further interpretation involving GW and MB. CH wrote the first draft of the article, all authors commented on and/or amended this and all authors read and approved the final manuscript.

ORCID iD

Christine Hazelton  <https://orcid.org/0000-0002-9554-4750>

References

- Aimola L, Lane AR, Smith DT, et al. (2014) Efficacy and feasibility of home-based training for individuals with homonymous visual field defects. *Neurorehabilitation and Neural Repair* 28(3): 207–218. DOI: 10.1177/1545968313503219.
- Ali M, Hazelton C, Lyden P, et al. (2013) Recovery from post-stroke visual impairment: evidence from a clinical trials resource. *Neurorehabilitation and Neural Repair* 27(2): 133–141.

- Borglin G (2015) The value of mixed methods for researching complex intervention. In: Richards DA and Hallberg IR (eds) *Complex Interventions in Health: An Overview of Research Methods*. New York: Routledge, pp. 29–45.
- Campbell R, Quilty B and Dieppe P (2003) Discrepancies between patients' assessments of outcome: qualitative study nested within a randomised controlled trial. *BMJ* 326(7383): 252–253. DOI: 10.1136/bmj.326.7383.252.
- Chen CS, Lee AW, Clarke G, et al. (2009) Vision-related quality of life in patients with complete homonymous hemianopia post stroke. *Topics in Stroke Rehabilitation* 16(6): 445–453.
- de Haan GA, Heutink J, Melis-Dankers BJM, et al. (2015a) Difficulties in daily life reported by patients with homonymous visual field defects. *Journal of Neuro-Ophthalmology* 35(3): 259–264.
- de Haan GA, Melis-Dankers BJM, Brouwer WH, et al. (2015b) The effects of compensatory scanning training on mobility in patients with homonymous visual field defects: a randomized controlled trial. *PLoS One* 10(8): 1–29. DOI: 10.1371/journal.pone.0134459.
- De Lisle J (2011) The benefits and challenges of mixing methods and methodologies: lessons learnt from implementing qualitatively led mixed methods research designs in Trinidad and Tobago. *Caribbean Curriculum* 18: 87–120.
- Hardiess G, Papageorgiou E, Schieferb U, et al. (2010) Functional compensation of visual field deficits in hemianopic patients under the influence of different task demands. *Visual Research* 50(12): 1158–1172.
- Hayes A, Chen CS, Clarke G, et al. (2012) Functional improvements following the use of the NVT vision rehabilitation program for patients with hemianopia following stroke. *NeuroRehabilitation* 31(1): 19–30.
- Hazelton C, Pollock A, Taylor A, et al. (2019a) A qualitative exploration of the effect of visual field loss on daily life in home-dwelling stroke survivors. *Clinical Rehabilitation* 33(7): 1264–1273. DOI: 10.1177/0269215519837580.
- Hazelton C, Pollock A, Walsh G, et al. (2019b) Scanning training for rehabilitation of visual field loss due to stroke: identifying and exploring training tools in use. *British Journal of Occupational Therapy* 82(8): 502–511. DOI: 10.1177/0308022618809900.
- Hepworth L and Rowe F (2016) Visual impairment following stroke – the impact on quality of life: a systematic review. *Ophthalmology Research: An International Journal* 5(2): 1–15. DOI: 10.9734/OR/2016/23272.
- Jacquin-Courtois S, Bays PM, Salemm R, et al. (2013) Rapid compensation of visual search strategy in patients with chronic visual field defects. *Cortex* 49(4): 994–1000. DOI: 10.1016/j.cortex.2012.03.025.
- Kazdin AE (2011) *Single-Case Research Designs: Methods for Clinical and Applied Settings*, 2nd ed. New York: Oxford University Press, pp. 287–293.
- Kerckhoff G, Munssinger U, Haaf E, et al. (1992) Rehabilitation of homonymous scotomata in patients with postgeniculate damage of the visual system: Saccadic compensation training. *Restorative Neurology & Neuroscience* 4(4): 245–254.
- Lamoreaux EL, Pesudovs K, Pallant JF, et al. (2008) An evaluation of the 10-item Vision Core Measure 1 (VCM1) scale (the core module of the vision related quality of life scale) using Rasch analysis. *Ophthalmic Epidemiology* 15(4): 224–233.
- Pambakian ALM, Mannan SK, Hodgson TL, et al. (2004) Saccadic visual search training: a treatment for patients with homonymous hemianopia. *Journal of Neurology, Neurosurgery and Psychiatry* 75(10): 1443–1448.
- Passamonti C, Bertini C and Ládavas E (2009) Audio-visual stimulation improves oculomotor patterns in patients with hemianopia. *Neuropsychologia* 47(2): 546–555. DOI: 10.1016/j.neuropsychologia.2008.10.008.
- Patton MQ (2002) *Qualitative Research and Evaluation Methods*, 3rd ed. Saint Paul: Sage Publications.
- Pollock A, George BS, Fenton M, et al. (2012) Top ten research priorities relating to life after stroke. *The Lancet Neurology* 11(3): 209.
- Pollock A, Hazelton C, Rowe FJ, et al. (2019) Interventions for visual field defects in people with stroke. *Cochrane Database of Systematic Reviews* (5) DOI: 10.1002/14651858.CD008388.pub3
- Pommerenke K, Markowitsch HJ, Pornmerenke K, et al. (1989) Rehabilitation training of homonymous visual field defects in patients with postgeniculate damage of the visual system. *Restorative Neurology & Neuroscience* 1(1): 47–63.
- Ritchie J and Spencer L (2002) Qualitative data analysis for applied policy research. In: Huberman AM and Miles MB (eds) *The Qualitative Researcher's Companion*, 1st ed. Thousand Oaks: Sage Publications, pp. 305–329.
- Roth T, Sokolov AN, Messias A, et al. (2009) Comparing explorative saccade and flicker training in hemianopia: a randomized controlled study. *Neurology* 72(4): 324–331.
- Rowe F (2013) *Care provision and unmet need for post stroke visual impairment*. Final report. Available at: https://www.stroke.org.uk/sites/default/files/final_report_unmet_need_2013.pdf (accessed 16 November 2016).
- Rubin HJ and Rubin IS (2005) *Qualitative Interviewing*, 2nd ed. Thousand Oaks: Sage Publications, pp. 108–129.
- Schuett S (2009) Eye movements in hemianopia and the rehabilitation of hemianopic dyslexia. Doctoral Thesis, Durham University, Durham, UK. Available at: <http://etheses.dur.ac.uk/2090/>.
- Schuett S, Heywood CA, Kentridge RW, et al. (2012) Rehabilitation of reading and visual exploration in visual field disorders: transfer or specificity? *Brain* 135(3): 912–921. DOI: 10.1093/brain/awr356.
- Smith JD (2012) Single-case experimental designs: a systematic review of published research and current standards. *Psychological Methods* 17(4): 510–550.
- Stelmack JA, Szlyk JP, Stelmack TR, et al. (2004) Psychometric properties of the Veterans Affairs low-vision visual functioning questionnaire. *Investigative Ophthalmology & Visual Science* 45(11): 3919–3928.
- Trauzettel-Klosinski S and Dietz K; and the IReST Study Group. (2012) Standardized assessment of reading performance: the new International Reading Speed Texts IReST. *Investigative Ophthalmology & Visual Science* 53(9): 5452–5461.
- Treisman A and Gelade G (1980) A feature-integration theory of attention. *Cognitive Psychology* 12(1): 97–136.
- Warren M (1993) A hierarchical model for evaluation and treatment of visual perceptual dysfunction in adult acquired brain injury, part 1. *American Journal of Occupational Therapy* 47(1): 42–54.
- Warren M (2009) Pilot study on activities of daily living limitations in adults with hemianopsia. *American Journal of Occupational Therapy* 63(5): 626–633.
- Zihl J (1995) Visual scanning behavior in patients with homonymous hemianopia. *Neuropsychologia* 33: 287–303.