

COMPLETE REDUCIBILITY FOR LIE SUBALGEBRAS AND SEMISIMPLIFICATION

MICHAEL BATE, SÖREN BÖHM, BENJAMIN MARTIN, GERHARD RÖHRLE,
AND LAURA VOGGESBERGER

To the memory of Irina Suprunenko

ABSTRACT. Let G be a connected reductive linear algebraic group over a field k . Using ideas from geometric invariant theory, we study the notion of G -complete reducibility over k for a Lie subalgebra \mathfrak{h} of the Lie algebra $\mathfrak{g} = \mathrm{Lie}(G)$ of G and prove some results when \mathfrak{h} is solvable or $\mathrm{char}(k) = 0$. We introduce the concept of a k -semisimplification \mathfrak{h}' of \mathfrak{h} ; \mathfrak{h}' is a Lie subalgebra of \mathfrak{g} associated to \mathfrak{h} which is G -completely reducible over k . This is the Lie algebra counterpart of the analogous notion for subgroups studied earlier by the first, third and fourth authors. As in the subgroup case, we show that \mathfrak{h}' is unique up to $\mathrm{Ad}(G(k))$ -conjugacy in \mathfrak{g} . Moreover, we prove that the two concepts are compatible: for H a closed subgroup of G and H' a k -semisimplification of H , the Lie algebra $\mathrm{Lie}(H')$ is a k -semisimplification of $\mathrm{Lie}(H)$.

1. INTRODUCTION

Let G be a connected reductive linear algebraic group over an arbitrary field k . We revisit the notion of G -complete reducibility over k for a Lie subalgebra \mathfrak{h} of the Lie algebra $\mathfrak{g} := \mathrm{Lie}(G)$ of G from [5, Def. 5.3]. A central theme of the present paper is an extension to the setting of Lie algebras of the construction of a k -semisimplification of a subgroup H of G from [10]. For motivation and an overview of the latter and further references, we refer to the introduction of [10]. The concept of the semisimplification of a module for a group or algebra is a well-known construction in representation theory. Building on this idea, for H a subgroup of $G(k)$, Serre introduced the concept of a “ G -analogue” of semisimplification from representation theory in [36, §3.2.4]. As in the subgroup case, one can think of the idea of a k -semisimplification of a subalgebra of \mathfrak{g} as an analogue of the Jordan-Hölder Theorem.

The definition of a k -semisimplification of a Lie subalgebra of \mathfrak{g} (Definition 5.4) for arbitrary k is new and generalizes the one for subgroups of G (Definition 5.1). We prove that the k -semisimplification of a subalgebra \mathfrak{h} of \mathfrak{g} is unique up to $\mathrm{Ad}(G(k))$ -conjugacy (Theorem 5.8). Moreover, for H a closed subgroup of G and H' a k -semisimplification of H , we show that $\mathrm{Lie}(H')$ is a k -semisimplification of $\mathrm{Lie}(H)$ (Theorem 5.9).

To prove Theorem 5.8 we use the theory of G -complete reducibility for subalgebras of \mathfrak{g} and some ideas from geometric invariant theory. In Theorem 5.14 we show that when $\mathrm{char}(k)$ is large enough, an ideal of a G -completely reducible subalgebra \mathfrak{h} is G -completely reducible and that the process of k -semisimplification behaves well under passing to ideals of \mathfrak{h} .

2010 *Mathematics Subject Classification.* 20G15 (14L24).

Key words and phrases. Semisimplification, G -complete reducibility, geometric invariant theory, rationality, cocharacter-closed orbits, degeneration of G -orbits.

We also prove some results of independent interest about G -complete reducibility for subalgebras, including Propositions 4.3 and 7.10. We consider solvable subalgebras in Section 6. Theorem 7.3 gives a necessary condition for a subalgebra to be G -completely reducible when $\text{char}(k) = p > 0$ is sufficiently large, and yields a characterisation of G -completely reducible subalgebras when $\text{char}(k) = 0$. We discuss some related results of Richardson, who pioneered the application of geometric invariant theory to the study of G -complete reducibility.

The notion of G -complete reducibility for subalgebras of \mathfrak{g} was first defined by McNinch [29] for algebraically closed k and was developed further in [12, §3] by the first, third and fourth authors; the non-algebraically closed case was first studied in [5, §5]. The approach via geometric invariant theory stems from work of Richardson [34], who studied subgroups of G and subalgebras of $\text{Lie}(G)$ — mainly for algebraically closed k in characteristic 0. Some of our arguments involve extending constructions from *op. cit.* to positive characteristic. We also use more recent techniques from [7] and [10] including some deep results from the theory of spherical buildings.

The theory of G -complete reducibility for Lie subalgebras closely parallels the theory of G -complete reducibility for subgroups [7], [10]. Some arguments are easier in the subalgebra case, as we need not worry about non-connected groups. There are, however, some extra problems for subalgebras, which can be traced back to the failure of certain normalisers to be smooth. It is to avoid these difficulties that we need assumptions on the characteristic of k : see Section 2.3.

2. PRELIMINARIES

2.1. Basic notation. Let k be a field with $\text{char}(k) = p \geq 0$. Following [15], [13], and [6], we regard an affine variety over a field k as a variety X over the algebraic closure \bar{k} together with a choice of k -structure. We write $X(k)$ for the set of k -points of X and $X(\bar{k})$ (or just X) for the set of \bar{k} -points of X . By a subvariety of X we mean a closed \bar{k} -subvariety of X ; a k -subvariety is a subvariety that is defined over k .

Throughout, G denotes a connected reductive linear algebraic k -group. Algebraic groups and their subgroups are assumed to be smooth (although we consider certain non-smooth subgroup schemes in Section 2.3). A k -defined affine G -variety X is an affine variety over k equipped with a k -defined morphic action of G . Two important examples of affine G -varieties in this paper are the group G itself, with G acting by inner automorphisms, and the Lie algebra $\mathfrak{g} = \text{Lie}(G)$ of G , with G acting by the adjoint action (that \mathfrak{g} admits a k -structure is [37, (12.2.3), (4.4.8)]). Recall that these two actions are closely related: if $g \in G$, then we have $\text{Inn}(g) : G \rightarrow G, x \mapsto gxg^{-1}$, and then $\text{Ad}(g) := d(\text{Inn}(g)) : \mathfrak{g} \rightarrow \mathfrak{g}$ is the differential (see below). To simplify notation we denote both of these actions by a dot: that is, if $g \in G$, then we let $g \cdot x = gxg^{-1}$ for each $x \in G$, and $g \cdot x = \text{Ad}(g)(x)$ for each $x \in \mathfrak{g}$. We define ad to be the usual adjoint action of \mathfrak{g} on \mathfrak{g} : so $\text{ad}(x)(y) = [x, y]$ for $x, y \in \mathfrak{g}$. If necessary we write Ad_G instead of Ad and ad_G instead of ad . Given any $m \in \mathbb{N}$, we may extend these actions to actions of G on $X = G^m$ or \mathfrak{g}^m , the m -fold Cartesian product of G or \mathfrak{g} , by setting

$$g \cdot (x_1, \dots, x_m) := (g \cdot x_1, \dots, g \cdot x_m)$$

for each $g \in G$ and $(x_1, \dots, x_m) \in X$. We refer to this action informally in both cases as the action *by simultaneous conjugation*.

When $\text{char}(k) = p > 0$, the Lie algebra \mathfrak{g} is *restricted* with a p -operation $\mathfrak{g} \rightarrow \mathfrak{g}, x \mapsto x^{[p]}$, [37, (4.4.3)]. By a subgroup of G we mean a closed \bar{k} -subgroup and by a k -subgroup we mean a subgroup that is defined over k . For a subgroup H of G we write H^0 for the identity component of H and $\text{Lie}(H)$ for its Lie algebra (which is a subalgebra of \mathfrak{g}). If H is k -defined, then $\text{Lie}(H)$ also admits a k -structure, [37, (12.2.3), (4.4.8)]. Subalgebras of \mathfrak{g} of the form $\text{Lie}(H)$ as above are called *algebraic*.

If $f: G_1 \rightarrow G_2$ is a homomorphism of algebraic groups then we denote the induced map from $\text{Lie}(G_1)$ to $\text{Lie}(G_2)$ by df .

2.2. Cocharacters and parabolic subgroups. We recall some basic notation and facts concerning cocharacters and parabolic subgroups in connected reductive groups G from [13], [34] and [37]. Recall that a cocharacter of G is a homomorphism of algebraic groups $\lambda: \mathbb{G}_m \rightarrow G$, where \mathbb{G}_m is the multiplicative group. We define $Y_k(G)$ to be the set of k -defined cocharacters of G and $Y(G) := Y_{\bar{k}}(G)$ to be the set of all cocharacters of G . Given $\lambda \in Y(G)$, we define

$$P_\lambda = \{g \in G \mid \lim_{a \rightarrow 0} \lambda(a)g\lambda(a)^{-1} \text{ exists}\}$$

and $L_\lambda = C_G(\text{Im}(\lambda)) \subseteq P_\lambda$ (for the definition of a limit, see [37, §3.2.13]). We have $P_\lambda = L_\lambda = G$ if and only if $\text{Im}(\lambda)$ is contained in the centre of G . Any parabolic subgroup (resp., any Levi subgroup of a parabolic subgroup of G) is of the form P_λ (resp., L_λ) [37, Prop. 8.4.5]. In fact, this also holds for k -defined parabolic and Levi subgroups of G , by the following, which is [37, Lem. 15.1.2(ii)]:

Lemma 2.1. *Every pair (P, L) consisting of a parabolic k -subgroup P of G and a Levi k -subgroup L of P is of the form $(P, L) = (P_\lambda, L_\lambda)$ for some $\lambda \in Y_k(G)$, and vice versa.*

For ease of reference, we record without proof some basic facts about these subgroups. The unipotent radical of a parabolic subgroup P of G is denoted by $R_u(P)$. The following is [34, 2.3].

Lemma 2.2. *If P is a k -defined parabolic subgroup then $R_u(P)$ is k -defined.*

The following is [13, Lem. 2.5 (i)+(iii)].

Lemma 2.3. *Let P be a parabolic subgroup of G and L a Levi subgroup of P .*

- (i) *We have $P \cong L \rtimes R_u(P)$, and this is a k -isomorphism if P and L are k -defined.*
- (ii) *Let T be a maximal torus of P . Then there is a unique Levi subgroup L of P such that $T \subseteq L$. If P and T are k -defined then L is k -defined.*
- (iii) *Any two Levi k -subgroups of an parabolic k -subgroup P are $R_u(P)(k)$ -conjugate.*

We denote the canonical projection from P to L by c_L ; this is k -defined if P and L are. Given any cocharacter λ of G such that $P = P_\lambda$ and $L = L_\lambda$ then we often write c_λ instead of c_L . We have $c_\lambda(g) = \lim_{a \rightarrow 0} \lambda(a)g\lambda(a)^{-1}$ for $g \in P_\lambda$; the kernel of c_λ is the unipotent radical $R_u(P_\lambda)$ and the set of fixed points of c_λ is L_λ .

All of this transfers over to the Lie algebra \mathfrak{g} through the adjoint action: since G acts on \mathfrak{g} by the adjoint action, so does any subgroup of G and, in particular, the image of any cocharacter of G . For $\lambda \in Y(G)$, we define and use throughout $\mathfrak{p}_\lambda := \text{Lie}(P_\lambda)$ and $\mathfrak{l}_\lambda := \text{Lie}(L_\lambda)$. We require some standard facts concerning Lie algebras of parabolic and Levi subgroups of G (cf. [34, §2.1]).

Lemma 2.4. *Let $x \in \mathfrak{g}$. Then with the notation from above we have*

- (i) $x \in \mathfrak{p}_\lambda$ if and only if $\lim_{a \rightarrow 0} \lambda(a) \cdot x$ exists;
- (ii) $x \in \mathfrak{l}_\lambda$ if and only if $\lim_{a \rightarrow 0} \lambda(a) \cdot x = x$ if and only if the image of λ centralizes x ;
- (iii) $x \in \text{Lie}(R_u(P_\lambda))$ if and only if $\lim_{a \rightarrow 0} \lambda(a) \cdot x$ exists and equals 0.

The map $c_\lambda := c_{\mathfrak{l}_\lambda} : \mathfrak{p}_\lambda \rightarrow \mathfrak{l}_\lambda$ given by $x \mapsto \lim_{a \rightarrow 0} \lambda(a) \cdot x$, where the action is given by adjoint action, coincides with the usual projection of \mathfrak{p}_λ onto \mathfrak{l}_λ . The kernel of c_λ is $\text{Lie}(R_u(P_\lambda))$, and the set of fixed points of c_λ is \mathfrak{l}_λ .

Let $m \in \mathbb{N}$ and recall that G acts on \mathfrak{g}^m by simultaneous conjugation. Given $\lambda \in Y(G)$, we have a map $\mathfrak{p}_\lambda^m \rightarrow \mathfrak{l}_\lambda^m$ given by $\mathbf{x} \mapsto \lim_{a \rightarrow 0} \lambda(a) \cdot \mathbf{x}$; we abuse notation slightly and also call this map c_λ . For any $\mathbf{x} \in \mathfrak{p}_\lambda^m$, there exists a Levi k -subgroup L of P_λ with $\mathbf{x} \in \mathfrak{l}^m$ if and only if $c_\lambda(\mathbf{x}) = u \cdot \mathbf{x}$ for some $u \in R_u(P_\lambda)(k)$, by [6, Prop. 2.11].

2.3. Smoothness of centralisers and normalisers. We are concerned with linear algebraic groups G over k : these are affine group schemes of finite type over k (assumed to be smooth). Note that smoothness is automatic if $\text{char}(k) = 0$, by a result of Cartier [19, II, §6, 1.1]. We do need to consider certain non-smooth subgroup schemes of G , as the lack of smoothness has implications for G -complete reducibility. If \mathfrak{h} is a subalgebra of \mathfrak{g} then we define $N_G(\mathfrak{h})$ to be the *scheme-theoretic* normaliser of \mathfrak{h} in G , and $C_G(\mathfrak{h})$ to be the *scheme-theoretic* centraliser of \mathfrak{h} in G . Define

$$\mathfrak{n}_{\mathfrak{g}}(\mathfrak{h}) := \{x \in \mathfrak{g} \mid [x, \mathfrak{h}] \subseteq \mathfrak{h}\}$$

and

$$\mathfrak{c}_{\mathfrak{g}}(\mathfrak{h}) := \{x \in \mathfrak{g} \mid [x, y] = 0 \text{ for all } y \in \mathfrak{h}\}.$$

It can be shown that $N_G(\mathfrak{h})$ is smooth if and only if $\mathfrak{n}_{\mathfrak{g}}(\mathfrak{h}) = \text{Lie}(N_G(\mathfrak{h}))$ and $C_G(\mathfrak{h})$ is smooth if and only if $\mathfrak{c}_{\mathfrak{g}}(\mathfrak{h}) = \text{Lie}(C_G(\mathfrak{h}))$. For further discussion, see [20] and [21]. For exact conditions for $C_G(\mathfrak{h})$ to be smooth, see [11, Thm. 1.2] and [20]; for explicit lower bounds on $\text{char}(k)$ for $N_G(\mathfrak{h})$ to be smooth, see [21, Thm. A].

Definition 2.5. Define $\text{char}(k) = p > 0$ to be *fabulous* for G if the following hold:

- (a) Centralisers and normalisers of subgroups of G and subalgebras of \mathfrak{g} are smooth.
- (b) If \mathfrak{h} is a subalgebra of \mathfrak{g} consisting of nilpotent elements then $\mathfrak{h} \subseteq \text{Lie}(B)$ for some Borel subgroup B of G .

Remarks 2.6. (i). If $x \in \mathfrak{g}$ is nilpotent and B is a Borel subgroup of G then $x \in \text{Lie}(B)$ if and only if $x \in \text{Lie}(R_u(B))$. So we can reformulate condition (b) as follows: if \mathfrak{h} is a subalgebra of \mathfrak{g} consisting of nilpotent elements then $\mathfrak{h} \subseteq \text{Lie}(U)$ for some maximal unipotent subgroup U of G .

(ii). If $\text{char}(k) = 0$, then both conditions above are satisfied, see [19, II, §6, 1.1], [17, Ch. VIII, §10, Cor. 2].

(iii). Suppose $k = \bar{k}$. Owing to [20, Thm. 1.1], centralisers of subgroups of G and centralisers of subalgebras of \mathfrak{g} are smooth if and only if $\text{char}(k)$ is 0 or a “pretty good prime” p for G , ([20, Def. 2.11]). The latter implies that p is not a torsion prime for the root system of G . It then follows from [27, Thm. 2.2 and Rems.(a)] that condition (b) holds. Hence condition (a) implies condition (b) in this instance. See also [21, Thm. 10.1].

(iv). Suppose $k = \bar{k}$. If \mathfrak{h} consists of nilpotent elements and $\dim(\mathfrak{h}) = 1$ then $\mathfrak{h} \subseteq \text{Lie}(B)$ for some Borel subgroup B of G : this follows from [15, IV.14.25 Prop.]. This can fail if $k \neq \bar{k}$. For example, let $G = \text{PGL}_2(k)$. For $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(k)$, we denote the image of A in

G by $\overline{\begin{pmatrix} a & b \\ c & d \end{pmatrix}}$, and likewise for elements of $\mathfrak{gl}_2(k)$. If $\text{char}(k) = 2$ and we choose $a \in k^{\frac{1}{2}}$

such that $a \notin k$ then the subspace of $\text{Lie}(\text{PGL}_2)$ spanned by $\overline{\begin{pmatrix} 0 & 1 \\ a^2 & 0 \end{pmatrix}}$ consists of nilpotent elements but is not contained in $\text{Lie}(B)$ for any Borel subgroup B of PGL_2 (compare [13, Rem. 5.10]). This is the Lie algebra counterpart of the phenomenon of “non-plongabilité” for unipotent elements of the group, cf. [39].

(v). If p is fabulous for G , then $Z(G) = C_G(G)$ is smooth, so $\mathfrak{z}(\mathfrak{g}) = \text{Lie}(Z(G))$; in particular, $\mathfrak{z}(\mathfrak{g}) = 0$ if G is semisimple.

Example 2.7. Suppose k is algebraically closed of characteristic $p = 2$ and let $G = \text{PGL}_2(k)$.

Let \mathfrak{h} be the abelian subalgebra of \mathfrak{g} spanned by $\overline{\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}}$ and $\overline{\begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}}$. It is easy to check that \mathfrak{h} is not contained in $\text{Lie}(B)$ for any Borel subgroup B of G . Similar examples are produced in [27] for any G such that p is a torsion prime for G .

Now let \mathfrak{m} be the subalgebra of \mathfrak{g} spanned by $\overline{\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}}$. Then $C_G(\mathfrak{m})(k)$ is (the image of) the group of upper unitriangular matrices and $N_G(\mathfrak{m})(k)$ is (the image of) the group of upper triangular matrices, but $\mathfrak{c}_{\mathfrak{g}}(\mathfrak{m}) = \mathfrak{h}$ and $\mathfrak{n}_{\mathfrak{g}}(\mathfrak{m}) = \mathfrak{g}$. Hence neither $C_G(\mathfrak{m})$ nor $N_G(\mathfrak{m})$ is smooth. A similar calculation shows that neither $C_G(\mathfrak{h})$ nor $N_G(\mathfrak{h})$ is smooth.

3. COCHARACTER-CLOSED ORBITS AND G -COMPLETE REDUCIBILITY

We introduce the notion of complete reducibility for subalgebras of \mathfrak{g} and explain the link with geometric invariant theory (GIT). At the end of the section, we extend to arbitrary k the main results from [29]. As in [10], our main tool from GIT is the notion of cocharacter-closure, introduced in [13] and [6].

Definition 3.1. Let X be a k -defined affine G -variety and let $x \in X$ (we do not require x to be a k -point). We say that the orbit $G(k) \cdot x$ is *cocharacter-closed over k* if for all $\lambda \in Y_k(G)$ such that $x' := \lim_{a \rightarrow 0} \lambda(a) \cdot x$ exists, x' belongs to $G(k) \cdot x$.

If $k = \bar{k}$ then it follows from the Hilbert-Mumford Theorem [25, Thm. 1.4] that $G(k) \cdot x$ is cocharacter-closed over k if and only if $G(k) \cdot x$ is closed.

The following is [6, Thm. 1.3].

Theorem 3.2 (Rational Hilbert-Mumford Theorem). *Let G, X, x be as above. Then there is a unique $G(k)$ -orbit \mathcal{O} such that (a) \mathcal{O} is cocharacter-closed over k , and (b) there exists $\lambda \in Y_k(G)$ such that $\lim_{a \rightarrow 0} \lambda(a) \cdot x$ belongs to \mathcal{O} .*

Next recall the notion of G -complete reducibility for subgroups of G .

Definition 3.3. Let H be a subgroup of G . We say that H is *G -completely reducible over k* (G -cr over k) if for any parabolic k -subgroup P of G such that P contains H , there is a

Levi k -subgroup L of P such that L contains H . We say that H is G -irreducible over k (G -ir over k) if H is not contained in any proper R -parabolic k -subgroup of G at all.

We say that H is G -completely reducible (G -cr) if H is G -completely reducible over \bar{k} .

For more on G -complete reducibility for subgroups of G , see [35], [36], [7]; our main focus in this paper is the analogue for Lie algebras. We first recall the definition of G -complete reducibility for Lie algebras and then also the link between this concept and GIT using generating tuples, due to Richardson. Most of what follows was originally written down in [5, §5].

Definition 3.4 ([5, Def. 5.3]). A Lie subalgebra \mathfrak{h} of \mathfrak{g} is G -completely reducible over k (G -cr over k) if for any parabolic k -subgroup P of G such that $\mathfrak{h} \subseteq \text{Lie}(P)$, there is a Levi k -subgroup L of P such that $\mathfrak{h} \subseteq \text{Lie}(L)$. We say that \mathfrak{h} is G -irreducible over k (G -ir over k) if \mathfrak{h} is not contained in $\text{Lie}(P)$ for any proper parabolic k -subgroup of G at all. We say that \mathfrak{h} is G -indecomposable over k (G -ind over k) if \mathfrak{h} is not contained in $\text{Lie}(L)$ for any proper Levi k -subgroup L of G .

As in the subgroup case, we say that \mathfrak{h} is G -completely reducible (resp., G -irreducible, G -indecomposable) if it is G -completely reducible over \bar{k} (resp., G -irreducible over \bar{k} , G -indecomposable over \bar{k}).

For $k = \bar{k}$, this notion is due to McNinch, see [29] and also [13, §5.3].

Example 3.5. Let k , G , \mathfrak{h} and \mathfrak{m} be as in Example 2.7. Then \mathfrak{h} is G -ir but the ideal \mathfrak{m} of \mathfrak{h} is not G -cr.

Example 3.6. If p is fabulous for G and $0 \neq \mathfrak{h}$ consists of nilpotent elements then \mathfrak{h} is not G -cr: for $\mathfrak{h} \subseteq \text{Lie}(B)$ for some Borel subgroup B of G , but clearly $\mathfrak{h} \not\subseteq \text{Lie}(T)$ for any maximal torus T of B .

Remark 3.7. The notion of G -complete reducibility for subgroups and subalgebras generalizes the concept of semisimplicity from representation theory in the following sense. If H is a subgroup of $\text{GL}(V)$ or $\text{SL}(V)$, or \mathfrak{h} is a subalgebra of $\mathfrak{gl}(V)$ or $\mathfrak{sl}(V)$, then H (resp. \mathfrak{h}) is $\text{GL}(V)$ -cr or $\text{SL}(V)$ -cr if and only if V is a semisimple H -module (resp. \mathfrak{h} -module). See [35], [36, (3.2.2)] for the case of subgroups; the case of subalgebras is almost identical.

Remark 3.8. Let H be a subgroup of G and \mathfrak{h} a subalgebra of \mathfrak{g} . If k'/k is an algebraic field extension then we may also regard G as a k' -group, and it therefore makes sense to ask whether H and \mathfrak{h} are G -cr over k' as well as whether they are G -cr over k .

Remark 3.9. Note that Definition 3.4 makes sense even if \mathfrak{h} is not k -defined. We also note that since $\mathfrak{p}_{g\lambda} = g \cdot \mathfrak{p}_\lambda$ and $\mathfrak{l}_{g\lambda} = g \cdot \mathfrak{l}_\lambda$ for any $\lambda \in Y(G)$ and any $g \in G$ (see, e.g., [7, §6]), it follows that \mathfrak{h} is G -cr over k if and only if every $\text{Ad}(G(k))$ -conjugate of \mathfrak{h} is. More generally, one can show that if \mathfrak{h} is G -cr over k (resp., G -ir over k , resp., G -ind over k) then so is $d\phi(\mathfrak{h})$, for any k -defined automorphism ϕ of G . Similar observations hold for subgroups.

We now recall the link to GIT. For this, we need the following definition.

Definition 3.10. Let \mathfrak{h} be a Lie algebra. We call $\mathbf{x} = (x_1, \dots, x_m) \in \mathfrak{h}^m$, for some $m \in \mathbb{N}$, a *generating tuple* for \mathfrak{h} if x_1, \dots, x_m is a generating set for \mathfrak{h} as a Lie algebra.

The next theorem shows the relevance of the previous definition to the study of complete reducibility. It is [5, Thm. 5.4]; see also [29, Thm. 1(i)] for the case $k = \bar{k}$.

Theorem 3.11. *Let \mathfrak{h} be a subalgebra of $\mathfrak{g} = \text{Lie}(G)$. Let $\mathbf{x} \in \mathfrak{g}^m$ be a generating tuple for \mathfrak{h} , and let G act on \mathfrak{g}^m by simultaneous conjugation. Then \mathfrak{h} is G -completely reducible over k if and only if the $G(k)$ -orbit of \mathbf{x} is cocharacter-closed in \mathfrak{g}^m over k .*

The next result is [5, Thm. 5.5]. Note that if k is perfect, then part (i) implies the equivalence of G -complete reducibility over k and G -complete reducibility (over \bar{k}), because the extension \bar{k}/k is separable for perfect k .

Proposition 3.12. *Let \mathfrak{h} be a subalgebra of \mathfrak{g} .*

- (i) *Suppose \mathfrak{h} is k -defined, and let k'/k be a separable algebraic field extension. Then \mathfrak{h} is G -completely reducible over k' if and only if \mathfrak{h} is G -completely reducible over k .*
- (ii) *Let S be a k -defined torus of $C_G(\mathfrak{h})$ and set $L = C_G(S)$. Then \mathfrak{h} is G -completely reducible over k if and only if \mathfrak{h} is L -completely reducible over k .*

The final ingredient we need for our first main result is the connection between complete reducibility of subgroups of G and the notion of complete reducibility in the spherical building of G . Recall that the spherical building $\Delta_k = \Delta_k(G)$ of G over k can be identified as a simplicial complex with the poset of k -parabolic subgroups of G , with inclusion reversed [38]; the parabolic subgroup G corresponds to the empty simplex. For each k -parabolic subgroup P of G , we let σ_P denote the corresponding simplex in Δ_k . We note that k -points of G induce simplicial automorphisms of Δ_k : for $g \in G(k)$ and a k -parabolic subgroup P of G , we can define $g \cdot \sigma_P = \sigma_{gPg^{-1}}$. Recall that the simplicial complex Δ_k has a *geometric realisation*, which we denote by $\overline{\Delta}_k$ [1, A.1.1]. We have the following definitions.

Definition 3.13. ([35, 2.1.4, 2.1.5, 2.2.1])

- (i) Two simplices σ, τ in Δ_k are called *opposite* if when we write $\sigma = \sigma_P$ and $\tau = \sigma_Q$ for k -parabolic subgroups P and Q of G , then P and Q are opposite in G ; that is, $P \cap Q$ is a common Levi subgroup of P and Q .
- (ii) A subcomplex Σ of Δ_k is called *convex* if the corresponding subset $\overline{\Sigma} \subseteq \overline{\Delta}_k$ of the geometric realisation is convex.
- (iii) A convex subcomplex Σ of Δ_k is said to be Δ_k -completely reducible (Δ_k -cr) if for every $\sigma \in \Sigma$, there exists $\tau \in \Sigma$ with τ opposite σ .

The following argument forms part of the proof of [29, Lem. 4]; it provides the key link between G -complete reducibility and the building Δ_k . We give the idea of the proof for completeness.

Proposition 3.14. *Suppose \mathfrak{h} is a subalgebra of \mathfrak{g} . Then $\Delta_k^{\mathfrak{h}} := \{\sigma_P \mid \mathfrak{h} \subseteq \text{Lie}(P)\}$ is a convex subcomplex of Δ_k , and it is Δ_k -completely reducible if and only if \mathfrak{h} is G -completely reducible over k .*

Sketch Proof. For the first assertion, the key observation is that the intersection of two k -parabolic subgroups P and Q of G is smooth, and hence $\text{Lie}(P \cap Q) = \text{Lie}(P) \cap \text{Lie}(Q)$, e.g., [24, §10.3]. Now the result follows from Serre's criterion [36, Prop. 3.1] for recognising a convex subcomplex of Δ_k : a collection Σ of simplices is a convex subcomplex if whenever P, Q, R are k -parabolic subgroups of G with $\sigma_P, \sigma_Q \in \Sigma$ and $P \cap Q \subseteq R$, then $\sigma_R \in \Sigma$.

To see that \mathfrak{h} is G -cr over k if and only if $\Delta_k^{\mathfrak{h}}$ is Δ_k -cr, the key again is the smoothness of the intersection of two k -parabolic subgroups. This implies that for two k -parabolic subgroups

P and Q of G , we can detect whether or not they are opposite (and hence correspond to opposite simplices of Δ_k) on the level of their Lie algebras. \square

Remark 3.15. An analogous result holds for subgroups, with a very similar proof [36].

Since each subalgebra \mathfrak{h} of \mathfrak{g} gives rise to a subcomplex $\Delta_k^{\mathfrak{h}}$ of Δ_k , we can use the so-called Centre Conjecture of Tits, which in fact is a theorem for subcomplexes, see [31] (G of classical type or type G_2), [26] (G of type F_4 or E_6) and [33] (G of type E_7 or E_8). We give a version which is suitable for our purposes.

Theorem 3.16 (Tits' Centre Theorem). *Let Δ_k be a thick spherical building, and Σ a convex subcomplex of Δ_k . Then (at least) one of the following holds:*

- (i) Σ is Δ_k -completely reducible;
- (ii) there exists a nonempty simplex $\sigma \in \Sigma$ which is fixed by all simplicial automorphisms of Δ_k which stabilize Σ .

The typical use of this theorem (as we see in the next proof below) is in guaranteeing the existence of a simplex as in part (ii) when Σ is *not* Δ_k -cr; such a simplex is often referred to as a *centre* of Σ . We can now prove the main result of this section. Over \bar{k} , this is [29, Thm. 1(ii)]; see also [13, Thm. 5.27, Ex. 5.29]. However, the proofs given there, which use the technology of optimal destabilising subgroups, do not go through over arbitrary k . Instead, we use the Centre Theorem.

Theorem 3.17. *Suppose H is a k -defined subgroup of G . If H^0 is G -completely reducible over k , then so is $\text{Lie}(H)$.*

Proof. Since $\text{Lie}(H) = \text{Lie}(H^0)$, and the hypothesis is that H^0 is G -cr, it is no loss to assume that H is connected. Let $\mathfrak{h} = \text{Lie}(H)$. Using Proposition 3.12(i) (and its analogue for subgroups [8, Thm. 1.1]), it is enough to prove the result over the separable closure k_s . These reductions imply that we may assume that $H(k)$ is dense in H , because this is true for a smooth connected k -group over a separably closed field k [30, Prop. 1.26(b), A.48]. Now suppose L is a minimal Levi k -subgroup of G containing H . Since $L = L_\lambda = C_G(\text{Im}(\lambda))$ for some $\lambda \in Y_k(C_G(H))$, Proposition 3.12(ii) (and its analogue for subgroups [5, Thm. 1.4]) implies that H and \mathfrak{h} are G -cr if and only if they are L -cr, so we may replace G with L . Then H is G -cr over k but is not contained in any proper Levi k -subgroup of G , so H is G -ir over k .

We now proceed with the proof of the theorem by contradiction. Suppose that \mathfrak{h} is not G -cr over k . The subgroup $H(k)$ acts on the building Δ_k by simplicial automorphisms, and the subcomplex $\Delta_k^{\mathfrak{h}}$ is stabilized by the action of $H(k)$ since H stabilizes its own Lie algebra. We are assuming that \mathfrak{h} is not G -cr over k , so this subcomplex is not Δ_k -cr, by Proposition 3.14, and hence there is a nonempty $H(k)$ -fixed simplex σ in $\Delta_k^{\mathfrak{h}}$ by Theorem 3.16. This simplex has the form σ_P for some proper k -parabolic subgroup P of G , and the fact that σ_P is $H(k)$ -fixed translates into the fact that P is normalized by $H(k)$. Since parabolic subgroups are self-normalizing, this shows that $H(k) \subseteq P$, which in turn gives $H \subseteq P$ because $H(k)$ is dense in H . This contradicts the conclusion of the first paragraph, that H is G -ir over k , so we are done. \square

Remarks 3.18. (i). We observe that the converse of Theorem 3.17 is false, already in the algebraically closed case. See the counterexample due to the third author in [29, §1]; see also Example 5.13 below.

(ii). If k is perfect, then Theorem 3.17 also holds with the hypothesis that H (instead of H^0) is G -cr. For, as has already been observed, if k is perfect then H is G -cr over k if and only if H is G -cr, and the same for $\text{Lie}(H)$. Now over \bar{k} , if H is G -cr then H^0 is G -cr too [7, Thm. 3.10], and the hypotheses of the theorem hold.

4. MAPS INDUCED BY INCLUSION

In this section we assume k is algebraically closed. We need some material on quotients in GIT. Recall that if G acts on an affine variety X then we can form the quotient variety $X//G$. The coordinate algebra $k[X//G]$ is by definition the subring of invariants $k[X]^G$ of the coordinate ring $k[X]$, and the inclusion of $k[X]^G$ in $k[X]$ induces a morphism π_X (or $\pi_{X,G}$) from X to $X//G$. If $x \in X$, there is a unique closed G -orbit $C(x)$ contained in the closure of the orbit $G \cdot x$. By the Hilbert-Mumford Theorem, there exists $\lambda \in Y(G)$ such that $\lim_{a \rightarrow 0} \lambda(a) \cdot x$ belongs to $C(x)$. Given $x_1, x_2 \in X$, we have $\pi_X(x_1) = \pi_X(x_2)$ if and only if $C(x_1) = C(x_2)$ if and only if $f(x_1) = f(x_2)$ for all $f \in k[X]^G$. In particular, if $G \cdot x_1$ and $G \cdot x_2$ are closed then $C(x_1) = G \cdot x_1$ and $C(x_2) = G \cdot x_2$, so $\pi_X(x_1) = \pi_X(x_2)$ if and only if x_1 and x_2 lie in the same G -orbit. Hence points of $X//G$ correspond to closed G -orbits in X . For background on quotient varieties and GIT, see [32, Ch. 3] or [3]. Here is a particular instance of this set-up. The group G acts on G^m by simultaneous conjugation, so we can form the quotient $G^m//G$. Likewise we can form the quotient $\mathfrak{g}^m//G$. There is a direct connection with G -complete reducibility arising from Theorem 3.11 as follows: points in $\mathfrak{g}^m//G$ correspond to closed G -orbits $G \cdot (x_1, \dots, x_m)$, and $(x_1, \dots, x_m) \in \mathfrak{g}^m$ yields a closed orbit if and only if the subalgebra \mathfrak{h} generated by the x_i is G -cr. Analogous statements hold for $G^m//G$.

Now let H be a reductive subgroup of G . The inclusion ι of H^m in G^m induces a map $\Psi : H^m//H \rightarrow G^m//G$. The third author proved that Ψ is a finite morphism of varieties [28, Thm. 1.1]; this has various consequences for the theory of G -completely reducible subgroups (see, e.g., [7, Cor. 3.8]). It follows that the image of Ψ is closed. In particular, $\pi_{G^m}(\iota(H^m))$ is a closed subset of $G^m//G$. Now consider the analogous situation for Lie algebras. The derivative $d\iota$ of ι maps \mathfrak{h}^m to \mathfrak{g}^m . Now $d\iota$ does give rise to a map ψ from $\mathfrak{h}^m//H$ to $\mathfrak{g}^m//G$ mapping $\pi_{\mathfrak{h}^m,H}(x_1, \dots, x_m)$ to $\pi_{\mathfrak{g}^m,G}(d\iota(x_1, \dots, x_m))$, but we see in Example 4.2 below that this map need not be finite.

First we need a preliminary result.

Proposition 4.1. *Let $(x_1, \dots, x_m) \in \mathfrak{g}^m$ and let \mathfrak{m} be the subalgebra spanned by the x_i . Then the following are equivalent:*

- (a) $\pi_{\mathfrak{g}^m,G}(x_1, \dots, x_m) = \pi_{\mathfrak{g}^m,G}(0, \dots, 0)$.
- (b) For every nonconstant homogeneous $f \in k[\mathfrak{g}^m]^G$, we have $f(x_1, \dots, x_m) = 0$.
- (c) There exists $\lambda \in Y(G)$ such that $\lim_{a \rightarrow 0} \lambda(a) \cdot (x_1, \dots, x_m) = (0, \dots, 0)$.
- (d) There exists a maximal unipotent subgroup U of G such that $\mathfrak{m} \subseteq \text{Lie}(U)$.

Proof. The equivalence of (a), (b) and (c) follows from the results given at the start of the section. The equivalence of (c) with (d) follows from Lemma 2.4(iii), since a maximal unipotent subgroup of G is the unipotent radical of a Borel subgroup. \square

Example 4.2. Let $p = 2$ and let $G = \mathrm{SL}_3(k)$. Let $H = \mathrm{PGL}_2(k)$. The adjoint representation of $\mathrm{SL}_2(k)$ on its Lie algebra gives rise to an embedding i of H in G ; with a suitable choice

of basis, this takes the form $i \left(\overline{\begin{pmatrix} a & b \\ c & d \end{pmatrix}} \right) = \overline{\begin{pmatrix} a^2 & c^2 & 0 \\ b^2 & d^2 & 0 \\ ac & bd & 1 \end{pmatrix}}$.

For $a \neq 0$, define $x_1 = \overline{\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}}$ and $x_2(a) = \overline{\begin{pmatrix} 0 & 0 \\ a & 0 \end{pmatrix}}$. Then the pairs $(x_1, x_2(a))$ span the subalgebra \mathfrak{h} from Example 2.7, which is H -ir by Example 3.5. It follows from Theorem 3.11 that $H \cdot (x_1, x_2(a))$ is closed for each a . It is easily seen that the $(x_1, x_2(a))$ are pairwise non-conjugate under the H -action. Hence the points $\pi_{\mathfrak{h}^2, H}(x_1, x_2(a))$ are pairwise distinct in $\mathfrak{h}^2 // H$. On the other hand, $di(\mathfrak{h}) \subseteq \mathrm{Lie}(B)$, where B is the Borel subgroup of upper triangular matrices in G , so $\pi_{\mathfrak{g}^2, G}(x_1, x_2(a)) = \pi_{\mathfrak{g}^2, G}(0, 0)$ for all a , by Proposition 4.1. This shows that the fibre $\psi^{-1}(\pi_{\mathfrak{g}^2, G}(0, 0))$ is infinite, so ψ cannot be finite.

If we put a restriction on p then the situation improves. Recall that if V is a rational G -module then $v \in V$ is said to be *unstable* if 0 belongs to the closure of $G \cdot v$, and *semistable* otherwise.

Proposition 4.3. *Suppose p is fabulous for G . Then the image of ψ is closed.*

Proof. It is enough by [4, Prop. 5.2] to show that if $(x_1, \dots, x_m) \in \mathfrak{h}^m$ is semistable for the H -action then it is semistable for the G -action. We prove the contrapositive. Suppose (x_1, \dots, x_m) is unstable for the G -action. Then the subalgebra \mathfrak{m} generated by the x_i consists of nilpotent elements, by Proposition 4.1. The hypothesis on p implies that $\mathfrak{m} \subseteq \mathrm{Lie}(U)$ for some maximal unipotent subgroup U of H . It follows from Proposition 4.1 that (x_1, \dots, x_m) is unstable for the H -action, so we are done. \square

Remark 4.4. Suppose p is fabulous for G . Choose any $f_1, \dots, f_t \in k[\mathfrak{g}^m // G]$ such that $\psi(\mathfrak{h}^m // H) = \{z \in \mathfrak{h}^m // G \mid f_1(z) = \dots = f_t(z) = 0\}$; such f_i exist by Proposition 4.3. Note that $\pi_{\mathfrak{g}^m, G}(\mathfrak{h}^m) = \psi(\mathfrak{h}^m // H)$ by construction. Regarding the f_i as elements of $k[\mathfrak{g}^m]^G$, we see that for any $(y_1, \dots, y_m) \in \mathfrak{g}^m$, $\pi_{\mathfrak{g}^m, G}(y_1, \dots, y_m)$ belongs to $\pi_{\mathfrak{g}^m, G}(\mathfrak{h}^m)$ if and only if $\pi_{\mathfrak{g}^m, G}(y_1, \dots, y_m)$ belongs to $\psi(\mathfrak{h}^m // H)$ if and only if $f_i(y_1, \dots, y_m) = 0$ for $1 \leq i \leq t$.

5. k -SEMISIMPLIFICATION

First, we recall the notion of k -semisimplification for subgroups of G and the main theorem from [10].

Definition 5.1. Let H be a subgroup of G . We say that a subgroup H' of G is a *k -semisimplification of H (for G)* if there exists a parabolic k -subgroup P of G and a Levi k -subgroup L of P such that $H \subseteq P$, $H' = c_L(H)$ and H' is G -completely reducible (or equivalently L -completely reducible, by [10, Prop. 3.6]) over k . We say *the pair (P, L) yields H'* .

Remarks 5.2. (i). Let H be a subgroup of G . If H is G -cr over k then clearly H is a k -semisimplification of itself, yielded by the pair (G, G) .

(ii). Given any subgroup H of G , [10, Rem. 4.3] guarantees the existence of a k -semisimplification of H .

Here is the main result [10, Thm. 4.5] from [10], which was proved in the special case $k = \bar{k}$ in [13, Prop. 5.14(i)], cf. [36, Prop. 3.3(b)]. The uniqueness asserted in Theorem 5.3 is akin to the theorem of Jordan–Hölder.

Theorem 5.3. *Let H be a subgroup of G . Then any two k -semisimplifications of H are $G(k)$ -conjugate.*

We now come to the analogue of Definition 5.1 for subalgebras of \mathfrak{g} .

Definition 5.4. Let \mathfrak{h} be a Lie subalgebra of \mathfrak{g} . We say that a Lie subalgebra \mathfrak{h}' of \mathfrak{g} is a k -semisimplification of \mathfrak{h} (for G) if there exists a parabolic k -subgroup P of G and a Levi k -subgroup L of P such that $\mathfrak{h} \subseteq \text{Lie}(P)$, $\mathfrak{h}' = c_{\text{Lie}(L)}(\mathfrak{h})$ and \mathfrak{h}' is G -completely reducible (or equivalently, by Proposition 3.12(ii), L -completely reducible) over k . We say *the pair* (P, L) *yields* \mathfrak{h}' .

Remarks 5.5. (i). Let \mathfrak{h} be a subalgebra of \mathfrak{g} . If \mathfrak{h} is already G -cr over k then clearly \mathfrak{h} is a k -semisimplification of itself, yielded by the pair (G, G) .

(ii). Suppose (P, L) yields a k -semisimplification \mathfrak{h}' of \mathfrak{h} . Let L_1 be another Levi k -subgroup of P . Then $L_1 = uLu^{-1}$ for some $u \in R_u(P)(k)$ by Lemma 2.3(iii), so consequently $c_{\text{Lie}(L_1)}(\mathfrak{h}) = u \cdot c_{\text{Lie}(L)}(\mathfrak{h})$. Hence (P, L_1) also yields a k -semisimplification of \mathfrak{h} . Because of this, when the choice of L doesn't matter we simply say that P yields a k -semisimplification of \mathfrak{h} .

(iii). It is straightforward to check that if ϕ is an automorphism of G (as a k -group), \mathfrak{h} is a subalgebra of \mathfrak{g} and (P, L) yields a k -semisimplification \mathfrak{h}' of \mathfrak{h} then $d\phi(\mathfrak{h}')$ is a k -semisimplification of $d\phi(\mathfrak{h})$, yielded by $(\phi(P), \phi(L))$.

The following is immediate from Lemma 2.1.

Lemma 5.6. *Suppose that \mathfrak{h}' is a k -semisimplification of \mathfrak{h} . Then there is $\lambda \in Y_k(G)$ such that \mathfrak{h}' is yielded by the pair (P_λ, L_λ) .*

As in the group case (Remark 5.2(ii)) we always have the existence of a k -semisimplification of an arbitrary subalgebra of \mathfrak{g} , due to the rational Hilbert-Mumford Theorem 3.2, as the following remark shows.

Remark 5.7. Suppose \mathfrak{h} is a subalgebra of \mathfrak{g} . Let $\mathbf{h} = (h_1, \dots, h_m) \in \mathfrak{h}^m$ be a generating tuple for \mathfrak{h} . Then $c_\lambda(\mathbf{h}) = (c_\lambda(h_1), \dots, c_\lambda(h_m))$ is a generating tuple for $c_\lambda(\mathfrak{h})$, for any $\lambda \in Y_k(G)$, and hence $c_\lambda(\mathfrak{h})$ is a k -semisimplification of \mathfrak{h} if and only if $G(k) \cdot c_\lambda(\mathbf{h})$ is cocharacter-closed over k , by Theorem 3.11. It follows from Theorem 3.2 that \mathfrak{h} admits at least one k -semisimplification: for we can choose $\lambda \in Y_k(G)$ such that $G(k) \cdot c_\lambda(\mathbf{h})$ is cocharacter-closed over k , so $c_\lambda(\mathfrak{h})$ is a k -semisimplification of \mathfrak{h} , yielded by (P_λ, L_λ) .

Here is the analogue of the main result Theorem 4.5 from [10] in the Lie algebra setting, which can be viewed as a kind of Jordan–Hölder theorem. Since the adjoint action is k -linear, the proof is easier to the one in [10], where a descending chain argument is needed.

Theorem 5.8. *Let \mathfrak{h} be a subalgebra of \mathfrak{g} . Then any two k -semisimplifications of \mathfrak{h} are $\text{Ad}(G(k))$ -conjugate.*

Proof. Let $\mathfrak{h}_1, \mathfrak{h}_2$ be k -semisimplifications of \mathfrak{h} . By Lemma 5.6, there exist $\lambda_1, \lambda_2 \in Y_k(G)$ such that $(P_{\lambda_1}, L_{\lambda_1})$ realizes \mathfrak{h}_1 and $(P_{\lambda_2}, L_{\lambda_2})$ realizes \mathfrak{h}_2 . Let $\mathbf{x} \in \mathfrak{h}^m$ be a generating tuple

for \mathfrak{h} . Then $c_{\lambda_i}(\mathbf{x})$ is a generating tuple for \mathfrak{h}_i for $i = 1, 2$, and each orbit $G(k) \cdot c_{\lambda_i}(\mathbf{x})$ is cocharacter-closed over k . It follows from the uniqueness result in the rational Hilbert-Mumford Theorem 3.2 that the two orbits $G(k) \cdot c_{\lambda_1}(\mathbf{x})$ and $G(k) \cdot c_{\lambda_2}(\mathbf{x})$ have to be equal. Thus there exists $g \in G(k)$ such that $g \cdot c_{\lambda_1}(\mathbf{x}) = c_{\lambda_2}(\mathbf{x})$. This means that the spanning set $c_{\lambda_1}(\mathbf{x})$ of \mathfrak{h}_1 is $\text{Ad}(G(k))$ -conjugate to the one of \mathfrak{h}_2 . Since the adjoint action is k -linear, \mathfrak{h}_1 and \mathfrak{h}_2 are also $\text{Ad}(G(k))$ -conjugate. \square

Next we study the connection between the notions of k -semisimplifications for subgroups and subalgebras. It turns out that they are compatible in a natural fashion.

Theorem 5.9. *Let H be a subgroup of G and let H' be a k -semisimplification of H . Then $\text{Lie}(H')$ is a k -semisimplification of $\text{Lie}(H)$.*

Proof. By Lemma 2.1, there is a $\lambda \in Y_k(G)$ such that (P_λ, L_λ) yields $H' = c_\lambda(H)$. It follows from Theorem 3.17 that $\text{Lie}(H')$ is G -cr over k . Since the differential of conjugation is the adjoint action, we have $dc_{L_\lambda} = c_{\lambda}$. Using Lemma 2.4, it follows that $\text{Lie}(H') = c_{\lambda}(\text{Lie}(H))$, and so the pair (P_λ, L_λ) also yields $\text{Lie}(H')$. \square

Next we revisit the example of Remark 3.18(i) in the context of Theorems 5.8 and 5.9, cf. [29].

Example 5.10. Suppose $\text{char}(k) = p > 0$. Let H be a non-trivial connected semisimple group and let $\varrho_i : H \rightarrow \text{SL}(V_i)$ be representations of H for $i = 1, 2$ with ϱ_1 semisimple and ϱ_2 not semisimple. Let $\varrho : H \rightarrow G := \text{SL}(V_1 \oplus V_2)$ be the representation given by $h \mapsto \varrho_1(h) \oplus \varrho_2(F(h))$, where $F : H \rightarrow H$ is the Frobenius endomorphism of H . Let J be the image of H under ϱ in G . Since $V_1 \oplus V_2$ is not semisimple as a J -module, J is not G -cr (cf. Remark 3.7). However, $\text{Lie}(J) = \text{Im } d\varrho_1 \oplus 0 \subseteq \mathfrak{g}$ is G -cr, showing that the converse of Theorem 3.17 fails.

Now let J' be a k -semisimplification of J . It follows from Theorem 5.9 that $\text{Lie}(J')$ is a k -semisimplification of $\text{Lie}(J)$ and thus, by Remark 5.5(a) and Theorem 5.8, that $\text{Lie}(J')$ and $\text{Lie}(J)$ are $\text{Ad}(G(k))$ -conjugate.

The following is the analogue of [10, Def. 4.6].

Definition 5.11. Let \mathfrak{h} be a subalgebra of \mathfrak{g} . We define $\mathcal{D}_k(\mathfrak{h})$ to be the set of $\text{Ad}(G(k))$ -conjugates of any k -semisimplification of \mathfrak{h} in \mathfrak{g} . This is well-defined by Theorem 5.8.

In the following two examples we show that not every element of $\mathcal{D}_k(\mathfrak{h})$ need be a k -semisimplification of \mathfrak{h} and that there is no direct relation between the notions of k -semisimplification and \bar{k} -semisimplification of a subalgebra of \mathfrak{g} .

Example 5.12. Let \mathfrak{h} be a subalgebra of \mathfrak{g} . As noted in Remark 5.5(a), if \mathfrak{h} is G -cr over k then \mathfrak{h} is a k -semisimplification of itself, yielded by the pair (G, G) . If \mathfrak{h} is G -ir, then \mathfrak{h} is not contained in $\text{Lie}(P)$ for any proper parabolic subgroup P of G , so \mathfrak{h} is the **only** k -semisimplification of itself.

Example 5.13. There are many examples in the literature of the following: a reductive group G over an imperfect field k , and a subgroup H of G such that H is G -cr over k but not G -cr, or H is G -cr but not G -cr over k ; see [2, Thm. 1.3], for example. Not all of these instances give rise to similar ones on the level of Lie algebras, even when the subgroup H is

connected, because of problems like that in Example 5.10. However, one of the most basic families of examples does work, as we now describe.

Let k be an imperfect field of characteristic p , and let $a \in k \setminus k^p$. Let t be a p^{th} root of a in \bar{k} ; then the extension $k' = k(t)$ is purely inseparable over k of degree p . Given a k' -group H' , we may form the *Weil restriction* $R_{k'/k}(H')$, which is a k -group, see [18, A.5] for example. (It is easiest to describe Weil restriction functorially: if we view H' as a functor from k' -algebras to groups, then $R_{k'/k}(H')$ is the corresponding functor from k -algebras to groups with $R_{k'/k}(H')(A) := H'(A \otimes_k k')$ for each k -algebra A .) Now let $H' = \mathbb{G}_m$ be the multiplicative group over k' . Then $H := R_{k'/k}(\mathbb{G}_m)$ is a p -dimensional abelian k -group with a $(p-1)$ -dimensional unipotent radical, but no connected normal unipotent k -subgroup (it is a basic example of a so-called *pseudo-reductive group*, see [18, Ex. 1.1.3]). The natural action of H' on k' by multiplication Weil restricts to give a p -dimensional representation of H over k which is irreducible: in terms of coordinates, this arises by writing down a k -basis for k' and interpreting the action through that basis, see [14, §5.2]. However, after base changing to \bar{k} , this module becomes indecomposable and not irreducible [14, Rem. 4.7(i)]. This means that H is GL_p -cr over k but not GL_p -cr. This example is due to McNinch (see [7, Ex. 5.11]).

If we turn our attention to the Lie algebras, we may identify the Lie algebra $\text{Lie}(H')$ with the additive group over k' , and we may view the multiplicative group H' as an open subset. Doing this is compatible with the action of H' and $\text{Lie}(H')$ on k' by multiplication. Therefore, after Weil restricting, we may identify the Lie algebra of H with a p -dimensional abelian Lie algebra over k containing H as an open subset, with the same compatibility between the corresponding representations (see [18, A.7.6] for more details on the Lie algebra of a Weil restriction). Therefore, in this case, we also have that $\text{Lie}(H)$ is GL_p -cr over k but not GL_p -cr. Note that $C_{\text{GL}_p}(\text{Lie}(H))$ is smooth.

The following theorem shows that under a mild restriction on k , the process of k -semisimplification behaves well under passing to ideals.

Theorem 5.14. *Suppose p is fabulous for G . Let \mathfrak{h} be a subalgebra of \mathfrak{g} and let \mathfrak{m} be an ideal of \mathfrak{h} . Then:*

- (a) *If \mathfrak{h} is G -completely reducible over k then so is \mathfrak{m} .*
- (b) *Every parabolic subgroup P of G which yields a k -semisimplification of \mathfrak{h} also yields one for \mathfrak{m} . In particular, there exist k -semisimplifications \mathfrak{h}' of \mathfrak{h} and \mathfrak{m}' of \mathfrak{m} such that \mathfrak{m}' is an ideal in \mathfrak{h}' .*

Proof. For (a), by the same reductions as at the start of the proof of Theorem 3.17, we may reduce to the case that $k = k_s$ and \mathfrak{h} is G -ir over k . We again proceed from this point using a contradiction argument invoking Tits' Centre Theorem 3.16. So suppose \mathfrak{m} is not G -cr over k . Then the subcomplex $\Delta_k^{\mathfrak{m}}$ of the building Δ_k is not Δ_k -cr, by Proposition 3.14, and hence there is a proper k -parabolic subgroup P of G such that $\sigma_P \in \Delta_k^{\mathfrak{m}}$ is fixed by all simplicial automorphisms of Δ_k stabilizing $\Delta_k^{\mathfrak{m}}$, by Theorem 3.16. Since $N_G(\mathfrak{m})(k)$ clearly stabilizes $\Delta_k^{\mathfrak{m}}$, we can conclude that $N_G(\mathfrak{m})(k) \subseteq P$. Now, because p is fabulous, $N_G(\mathfrak{h})$ is smooth, and because $k = k_s$ this means that the k -points of $N_G(\mathfrak{m})$ are dense by [15, AG.13.3 Cor.]. Therefore, we may conclude that $N_G(\mathfrak{m}) \subseteq P$. Finally, this gives

$$\mathfrak{m} \subseteq \mathfrak{h} \subseteq \mathfrak{n}_{\mathfrak{g}}(\mathfrak{m}) = \text{Lie}(N_G(\mathfrak{m})) \subseteq \text{Lie}(P),$$

where the equality in the middle follows from smoothness of $N_G(\mathfrak{m})$ again. This gives the required contradiction, as we had reduced to the case that \mathfrak{h} is G -ir over k .

For (b), pick any $\lambda \in Y_k(G)$ such that (P_λ, L_λ) yields a k -semisimplification $\mathfrak{h}' := c_\lambda(\mathfrak{h})$ of \mathfrak{h} . Then $c_\lambda(\mathfrak{h})$ is G -cr over k and, as c_λ is a Lie algebra homomorphism, $c_\lambda(\mathfrak{m})$ is an ideal in $c_\lambda(\mathfrak{h})$. Now $c_\lambda(\mathfrak{h})$ and $c_\lambda(\mathfrak{m})$ satisfy the hypotheses of the theorem, so $c_\lambda(\mathfrak{m})$ is G -cr over k by (a). Hence (P_λ, L_λ) yields a semisimplification $\mathfrak{m}' := c_\lambda(\mathfrak{m})$ of \mathfrak{m} as well, and \mathfrak{m}' is an ideal in \mathfrak{h}' . \square

Remark 5.15. Both parts of Theorem 5.14 are false without the assumption on p : see Example 3.5.

6. G -TORAL AND SOLVABLE SUBALGEBRAS

We assume in this section that k is algebraically closed. We study G -complete reducibility properties of solvable and G -toral subalgebras \mathfrak{h} of \mathfrak{g} (see Definition 6.4 for the latter).

Definition 6.1. We call a Lie subalgebra \mathfrak{h} of \mathfrak{g} *Jordan-closed* if for every $x \in \mathfrak{h}$, the semisimple part x_s and nilpotent part x_n of x both belong to \mathfrak{h} . We define the *Jordan closure* \mathfrak{h}^J of \mathfrak{h} to be the smallest Jordan-closed Lie subalgebra of \mathfrak{g} that contains \mathfrak{h} .

Remarks 6.2. (a). It is clear that \mathfrak{h}^J is well-defined. Here is an explicit construction. We define an increasing chain of subalgebras \mathfrak{h}_i of \mathfrak{g} as follows. Set $\mathfrak{h}_0 = \mathfrak{h}$. Given \mathfrak{h}_i , let \mathfrak{h}_{i+1} be the subalgebra generated by the elements of the form x_s and x_n for $x \in \mathfrak{h}_i$. For dimension reasons, the chain becomes stationary and we have $\mathfrak{h}_n = \mathfrak{h}^J$ for n sufficiently large.

(b). If \mathfrak{h} is algebraic then \mathfrak{h} is Jordan-closed. In particular, if P is a parabolic subgroup of G and L is a Levi subgroup of P then $\text{Lie}(P)$ and $\text{Lie}(L)$ are Jordan-closed. It follows easily that for any subalgebra \mathfrak{m} of \mathfrak{g} , \mathfrak{m} is G -cr (resp., G -ir, resp., G -ind) if and only if \mathfrak{m}^J is. Also, if $\text{char}(k) = 0$, \mathfrak{h} is a subalgebra of \mathfrak{g} and \mathfrak{h} is semisimple then \mathfrak{h} is algebraic (see [34, Lem. 3.2]), so \mathfrak{h} is Jordan-closed.

Remark 6.3. Let \mathfrak{h} be a subalgebra of \mathfrak{g} and let $f: G \rightarrow M$ be an epimorphism of connected reductive groups. We have $df(x)_s = df(x_s)$ and $df(x)_n = df(x_n)$ for any $x \in \mathfrak{h}$, so $df(\mathfrak{h})$ is Jordan-closed if \mathfrak{h} is. The converse also holds if f is an embedding, for then df is injective.

Definition 6.4. A subalgebra \mathfrak{h} of \mathfrak{g} is *G -toral* if $\mathfrak{h} \subseteq \text{Lie}(S)$ for some torus S of G .

Remark 6.5. Recall that a Lie algebra \mathfrak{h} is said to be *toral* if every element of \mathfrak{h} is ad-semisimple; in this case, \mathfrak{h} is abelian [22, Lem. 8.1]. Clearly, if \mathfrak{h} is G -toral then \mathfrak{h} is toral, but the converse is false: e.g., take \mathfrak{h} to be a nonzero abelian subalgebra consisting of nilpotent elements.

The following is the counterpart of [24, Lem. 11.24] for subalgebras of \mathfrak{g} .

Lemma 6.6. *Let \mathfrak{h} be a G -toral subalgebra of \mathfrak{g} . Then \mathfrak{h} is G -completely reducible.*

Proof. Choose a torus S such that $\mathfrak{h} \subseteq \text{Lie}(S)$. By Proposition 3.12(ii) it suffices to prove that \mathfrak{h} is $C_G(S)$ -cr. But this is clear because S is central in $C_G(S)$, so $\text{Lie}(S)$ is contained in $\text{Lie}(P)$ and $\text{Lie}(L)$ for every parabolic subgroup P and every Levi subgroup L of $C_G(S)$. \square

Let \mathfrak{m} be a G -toral subalgebra of \mathfrak{g} : say, $\mathfrak{m} \subseteq \text{Lie}(S)$, where S is a torus of G . Then $N_G(\mathfrak{m})$ is reductive. To see this, note that $N_G(\mathfrak{m})$ contains T , where T is a maximal torus of G containing S , so $N_G(\mathfrak{m})^0$ is generated by T and by certain root groups U_α with respect

to T . But if α is a root, $g \in U_\alpha$ and $x \in \text{Lie}(S)$ then $g \cdot x - x \in \text{Lie}(U_\alpha)$, so U_α normalises \mathfrak{m} if and only if U_α centralises \mathfrak{m} if and only if $d\alpha$ annihilates \mathfrak{m} . We deduce that $U_\alpha \subseteq N_G(\mathfrak{m})$ if and only if $U_{-\alpha} \subseteq N_G(\mathfrak{m})$, and reductivity of $N_G(\mathfrak{m})$ follows. This argument also shows that $N_G(\mathfrak{m})^0 = C_G(\mathfrak{m})^0$.

Suppose further that p is fabulous for G . Then centralisers and normalisers are smooth, so

$$(6.7) \quad \mathfrak{m} \subseteq \mathfrak{n}_{\mathfrak{g}}(\mathfrak{m}) = \mathfrak{c}_{\mathfrak{g}}(\mathfrak{m}) = \text{Lie}(C_G(\mathfrak{m})^0).$$

Moreover, let $K = C_G(\mathfrak{m})^0$, let $Z = C_G(K)^0$ and let $L = C_G(Z)$. As p is fabulous for G , Z is smooth. We have $Z \subseteq Z(K)^0$: for if $g \in C_G(K)$ then g centralises \mathfrak{m} since $\mathfrak{m} \subseteq \text{Lie}(K)$ by (6.7), so $C_G(K)^0$ is a connected subgroup of $C_G(\mathfrak{m})$. This implies that Z is a torus, as K is reductive. Hence L is a Levi subgroup of G . We have $\mathfrak{m} \subseteq \text{Lie}(Z)$ by smoothness of Z since K centralises \mathfrak{m} . It follows that $\mathfrak{m} \subseteq \text{Lie}(L)$ and that if $\mathfrak{m} \not\subseteq \mathfrak{z}(\mathfrak{g})$ then L is proper.

Lemma 6.8. *Let \mathfrak{h} be a subalgebra of \mathfrak{g} such that every element of \mathfrak{h} is semisimple. Then \mathfrak{h} is G -toral.*

Proof. Note that \mathfrak{h} is abelian by Remark 6.5. We use induction on $\dim(G)$. The result holds trivially if $\dim(G) = 0$, so let G be arbitrary. If $\mathfrak{h} \subseteq \mathfrak{z}(\mathfrak{g})$ then $\mathfrak{h} \subseteq \text{Lie}(T)$ for any maximal torus T of G , so we are done. Otherwise there exists $x \in \mathfrak{h}$ such that $C_G(x)^0$ is a proper reductive subgroup of G . Now $C_G(x)^0$ is smooth by [15, 9.1 Prop.] since x is semisimple, so $\mathfrak{h} \subseteq \text{Lie}(C_G(x)^0)$. By our induction hypothesis, \mathfrak{h} is $C_G(x)^0$ -toral, so \mathfrak{h} is G -toral. \square

Lemma 6.9. *Let \mathfrak{h} be a G -toral subalgebra of \mathfrak{g} and let $x \in \mathfrak{g}$ such that x is semisimple. Suppose that x centralises \mathfrak{h} , or that $\text{char}(k) = 0$ and x normalises \mathfrak{h} . Then $k \cdot x + \mathfrak{h}$ is G -toral.*

Proof. Choose an embedding of G in GL_n for some $n \in \mathbb{N}$. If x centralises \mathfrak{h} then $k \cdot x \cup \mathfrak{h}$ consists of pairwise commuting semisimple matrices. Hence by a standard theorem from linear algebra, the matrices in $k \cdot x \cup \mathfrak{h}$ are simultaneously diagonalisable in $M_n(k)$. This implies that every element of $k \cdot x + \mathfrak{h}$ is semisimple, so $k \cdot x + \mathfrak{h}$ is G -toral by Lemma 6.8.

Now suppose $\text{char}(k) = 0$ and x normalises \mathfrak{h} . Since x is semisimple, x acts semisimply on \mathfrak{h} . Let $y \in \mathfrak{h}$ be any nonzero eigenvector of $\text{ad}(x)$, with eigenvalue a . A simple calculation shows that $\text{ad}(x)(y^t) = tay^t$ for any $t \in \mathbb{N}$, where y^t denotes the usual matrix power of y in $M_n(k) = \text{Lie}(\text{GL}_n)$. But $y^t \neq 0$ for any $t \in \mathbb{N}$ as y is semisimple, which forces $ta = 0$ for all but finitely many t . This implies that $a = 0$ since we are in characteristic 0. We deduce that x centralises \mathfrak{h} , and the result follows from the previous paragraph. \square

Lemma 6.10. *Suppose p is fabulous for G . Let \mathfrak{h} be an abelian G -cr subalgebra of \mathfrak{g} . Then \mathfrak{h} is G -toral.*

Proof. Let $x \in \mathfrak{h}$. Then $\mathfrak{c}_{\mathfrak{g}}(x) = \text{Lie}(C_G(x))$, so if $y \in \mathfrak{c}_{\mathfrak{g}}(x)$ then $y_s \in \mathfrak{c}_{\mathfrak{g}}(x)$ and $y_n \in \mathfrak{c}_{\mathfrak{g}}(x)$, so y_s and y_n centralise x . By a similar argument, y_s and y_n centralise x_s and x_n . It follows from the construction described in Remark 6.2(a) that \mathfrak{h}^J is abelian. Now \mathfrak{h}^J is G -cr by Remark 6.2(b), and it is enough to prove that \mathfrak{h}^J is G -toral. Hence we can assume without loss that \mathfrak{h} is Jordan-closed.

By Lemma 6.8, it suffices to show that \mathfrak{h} consists of semisimple elements. Suppose not. Then there exists $0 \neq y \in \mathfrak{h}$ such that y is nilpotent. The 1-dimensional subspace \mathfrak{m} spanned by y is an ideal of \mathfrak{h} , so \mathfrak{m} is G -cr by Theorem 5.14. But this is impossible by Example 3.6. This completes the proof. \square

Lemma 6.11. *Suppose p is fabulous for G . Let \mathfrak{h} be a solvable G -completely reducible subalgebra of \mathfrak{g} . Then \mathfrak{h} is G -toral.*

Proof. We use induction on $\dim(G) + \dim(\mathfrak{h})$. The ideal $[\mathfrak{h}, \mathfrak{h}]$ is G -cr by Theorem 5.14 and $[\mathfrak{h}, \mathfrak{h}] \subseteq [\mathfrak{g}, \mathfrak{g}] = \text{Lie}([G, G])$. Now $[\mathfrak{h}, \mathfrak{h}]$ is a proper subalgebra of \mathfrak{h} , so $[\mathfrak{h}, \mathfrak{h}]$ is G -toral by our induction hypothesis. If $[\mathfrak{h}, \mathfrak{h}] = 0$ then \mathfrak{h} is abelian, so \mathfrak{h} is G -toral by Lemma 6.10. Otherwise $0 \neq [\mathfrak{h}, \mathfrak{h}]$ is not contained in $\mathfrak{z}(\mathfrak{g})$ as $[G, G]$ is semisimple (see Remarks 2.6(v)), so by the discussion following (6.7), $\mathfrak{h} \subseteq N_{\mathfrak{g}}([\mathfrak{h}, \mathfrak{h}]) \subseteq \text{Lie}(L)$ for some proper Levi subgroup L of G . Then \mathfrak{h} is L -cr by Proposition 3.12(ii) and $\dim(L) < \dim(G)$, so \mathfrak{h} is L -toral by our induction hypothesis. Hence \mathfrak{h} is G -toral, as required. \square

We can now give a classification result for maximal solvable subalgebras of \mathfrak{g} when p is fabulous for G : compare [21, Thm. D(b)].

Proposition 6.12. *Suppose p is fabulous for G . Let \mathfrak{h} be a solvable subalgebra of \mathfrak{g} . Then $\mathfrak{h} \subseteq \mathfrak{h}^J \subseteq \text{Lie}(B)$ for some Borel subgroup B of G . In particular, a maximal solvable subalgebra of \mathfrak{g} is the Lie algebra of some Borel subgroup.*

Proof. Since $\text{Lie}(B)$ is Jordan-closed for any Borel subgroup B of G (Remarks 6.2(b)), it suffices to prove that $\mathfrak{h} \subseteq \text{Lie}(B)$ for some Borel subgroup B . Let $\lambda \in Y(G)$ such that (P_λ, L_λ) yields a k -semisimplification of \mathfrak{h} . Then $\mathfrak{s} := c_\lambda(\mathfrak{h}) \subseteq \mathfrak{l}_\lambda$ is solvable and G -cr and $\mathfrak{h} \subseteq \mathfrak{s} + \text{Lie}(R_u(P_\lambda))$. Now \mathfrak{s} is L_λ -cr by Proposition 3.12(ii), so \mathfrak{s} is L_λ -toral by Lemma 6.8: say, $\mathfrak{s} \subseteq \text{Lie}(S)$ for some torus S of L_λ . It follows that $\mathfrak{h} \subseteq \text{Lie}(S) + \text{Lie}(R_u(P_\lambda)) = \text{Lie}(SR_u(P_\lambda))$. But $SR_u(P_\lambda)$ is contained in a Borel subgroup of G , so the first assertion follows. The second assertion is immediate. \square

Corollary 6.13. *Suppose p is fabulous for G . Let \mathfrak{h} be a Jordan-closed solvable subalgebra of \mathfrak{g} and let \mathfrak{n} be the set of nilpotent elements of \mathfrak{h} . Let $\lambda \in Y(G)$ such that (P_λ, L_λ) yields a semisimplification of \mathfrak{h} . Then:*

- (a) $\mathfrak{n} = \mathfrak{h} \cap \text{Lie}(R_u(P_\lambda))$. In particular, \mathfrak{n} is an ideal of \mathfrak{h} .
- (b) There is a G -toral subalgebra \mathfrak{s} of \mathfrak{h} such that $\mathfrak{h} = \mathfrak{s} \oplus \mathfrak{n}$.

Proof. (a). Since $c_\lambda(\mathfrak{h})$ is solvable and G -cr, $c_\lambda(\mathfrak{h})$ consists of semisimple elements by Lemma 6.11. Hence c_λ kills every nilpotent element of \mathfrak{h} , so $\mathfrak{n} \subseteq \text{Lie}(R_u(P_\lambda))$. But $\text{Lie}(R_u(P_\lambda))$ consists of nilpotent elements, so $\mathfrak{n} = \mathfrak{h} \cap \text{Lie}(R_u(P_\lambda))$. This is an ideal of \mathfrak{h} because $\mathfrak{h} \subseteq \mathfrak{p}_\lambda$.

(b). Let \mathfrak{s} be a maximal G -toral subalgebra of \mathfrak{h} . We claim that $c_\lambda(\mathfrak{s}) = c_\lambda(\mathfrak{h})$. Suppose not. Now \mathfrak{s} is abelian and consists of ad-semisimple elements, so \mathfrak{s} acts completely reducibly on \mathfrak{h} . Let $\mathfrak{h}_0 = \mathfrak{c}_{\mathfrak{h}}(\mathfrak{s})$ be the trivial weight space for the action of \mathfrak{s} on \mathfrak{h} ; then $c_\lambda(\mathfrak{h}_0) = c_\lambda(\mathfrak{h})$. Note that $\mathfrak{s} \subseteq \mathfrak{h}_0$ as \mathfrak{s} is abelian.

By hypothesis, there exists $x \in \mathfrak{h}_0$ such that $c_\lambda(x) \notin c_\lambda(\mathfrak{s})$. As p is fabulous for G , $\mathfrak{c}_{\mathfrak{g}}(\mathfrak{s}) = \text{Lie}(C_G(\mathfrak{s}))$ is Jordan-closed, by Remarks 6.2(b), and \mathfrak{h} is Jordan-closed by assumption, so $\mathfrak{h}_0 = \mathfrak{c}_{\mathfrak{h}}(\mathfrak{s}) = \mathfrak{c}_{\mathfrak{g}}(\mathfrak{s}) \cap \mathfrak{h}$ is Jordan-closed. Hence $x_s \in \mathfrak{h}_0$ and $c_\lambda(x_s) = c_\lambda(x) \notin c_\lambda(\mathfrak{s})$; in particular, $x_s \notin \mathfrak{s}$. Since x_s is semisimple and commutes with \mathfrak{s} , the subalgebra $\mathfrak{s} \oplus k \cdot x_s$ of \mathfrak{h} is toral by Lemma 6.9. But this contradicts the choice of \mathfrak{s} . We deduce that $c_\lambda(\mathfrak{s}) = c_\lambda(\mathfrak{h})$, which implies that $\mathfrak{h} = \mathfrak{s} \oplus \mathfrak{n}$. \square

7. CHARACTERISTIC 0

We keep our assumption that k is algebraically closed. If H is a G -cr subgroup of G then H is reductive, and the converse also holds if $\text{char}(k) = 0$ [36, Prop. 4.2]. This gives an intrinsic characterisation of G -cr subgroups in characteristic 0: a subgroup H is G -cr if and only if it is reductive. (Note that a reductive subgroup of G need not be G -cr in positive characteristic: e.g., see [7, Ex. 3.45].) The situation for Lie algebras is not so clear-cut.

Example 7.1. Consider the 1-dimensional Lie algebra k . We may embed k in \mathfrak{sl}_2 as the subalgebra of traceless diagonal matrices, or as the subalgebra of strictly upper triangular matrices. It is clear that in the first case, the image of k is SL_2 -cr, while in the second it is not.

This example shows that even in characteristic 0, we cannot determine whether a subalgebra \mathfrak{h} of \mathfrak{g} is G -cr by looking only at the intrinsic properties of \mathfrak{h} . We can trace the problem to the lack of a Jordan decomposition for \mathfrak{h} : there is no intrinsic notion of semisimple and nilpotent elements. We do have notions of ad-semisimple and ad-nilpotent elements, but they don't detect properties of $\mathfrak{z}(\mathfrak{h})$.

There is an intrinsic characterisation for when a restricted subalgebra \mathfrak{h} of \mathfrak{g} is G -cr when the characteristic is positive and sufficiently large, using the notion of “ p -reductive” subalgebras: see [21, Cor. 10.3]. In *loc. cit.* one makes use of the extra structure arising from the p -power map on \mathfrak{h} . Below we give a counterpart to [21, Cor. 10.3] by providing a characterisation of G -cr subalgebras of \mathfrak{g} in characteristic 0 (Theorem 7.3), and we give an explicit description for a semisimplification of an arbitrary subalgebra in characteristic 0. Different methods from those of *op. cit.* are required, as there is no notion of restricted structure here.

The content of Theorem 7.3 in characteristic 0 follows quickly from work of Richardson — see Remark 7.8 below. We give an independent proof, however, as some of our results hold under the weaker hypothesis that p is fabulous for G .

The next result is a Lie algebra counterpart of [7, Thm. 3.46]; the analogues to the separability conditions in *loc. cit.* hold here because of the assumption on p .

Proposition 7.2. *Suppose p is fabulous for G . Let \mathfrak{h} be a subalgebra of \mathfrak{g} such that \mathfrak{h} acts semisimply on \mathfrak{g} . Then \mathfrak{h} is G -completely reducible.*

Proof. Suppose for a contradiction that \mathfrak{h} is not G -cr. Choose $\mathbf{x} = (x_1, \dots, x_m) \in \mathfrak{h}^m$ for some $m \in \mathbb{N}$ such that the x_i span \mathfrak{h} . The orbit $G \cdot \mathbf{x}$ is not closed (Theorem 3.11), so there exists $\lambda \in Y(G)$ such that $\mathbf{x}' := \lim_{a \rightarrow 0} \lambda(a) \cdot \mathbf{x}$ exists and $G \cdot \mathbf{x}'$ is closed. This implies that $\dim(G \cdot \mathbf{x}) > \dim(G \cdot \mathbf{x}')$. Consequently, letting \mathfrak{h}' be the subalgebra generated by the components of \mathbf{x}' , we have $\dim(C_G(\mathfrak{h}')) > \dim(C_G(\mathfrak{h}))$, so

$$\dim(\mathfrak{c}_{\mathfrak{g}}(\mathfrak{h}')) = \dim(\text{Lie}(C_G(\mathfrak{h}'))) > \dim(\text{Lie}(C_G(\mathfrak{h}))) = \dim(\mathfrak{c}_{\mathfrak{g}}(\mathfrak{h})),$$

where the equalities hold because p is fabulous for G .

Now $\text{ad}_G(\mathfrak{h})$ is $\text{GL}(\mathfrak{g})$ -cr, so $\text{GL}(\mathfrak{g}) \cdot \text{ad}_G(\mathbf{x})$ is closed. Hence $\text{ad}_G(\mathbf{x}') = \text{ad}_G(\lim_{a \rightarrow 0} \lambda(a) \cdot \mathbf{x}) = \lim_{a \rightarrow 0} \lambda(a) \cdot \text{ad}_G(\mathbf{x})$ is $\text{GL}(\mathfrak{g})$ -conjugate to $\text{ad}_G(\mathbf{x})$. But $\mathfrak{c}_{\mathfrak{g}}(\mathfrak{h}')$ and $\mathfrak{c}_{\mathfrak{g}}(\mathfrak{h})$ are precisely the subsets of \mathfrak{g} annihilated by $\text{ad}_G(\mathfrak{h}')$ and $\text{ad}_G(\mathfrak{h})$, respectively, so $\mathfrak{c}_{\mathfrak{g}}(\mathfrak{h}')$ and $\mathfrak{c}_{\mathfrak{g}}(\mathfrak{h})$ are $\text{GL}(\mathfrak{g})$ -conjugate, which implies they have the same dimension. This gives a contradiction. We conclude that \mathfrak{h} must be G -cr after all, so we are done. \square

Before we proceed to Theorem 7.3, we need to recall some standard Lie algebra theory. If \mathfrak{h} is a Lie algebra then we define $\text{rad}(\mathfrak{h})$ to be the *solvable radical* of \mathfrak{h} : that is, the unique largest solvable ideal of \mathfrak{h} . We say that \mathfrak{h} is *semisimple* if $\text{rad}(\mathfrak{h}) = 0$, [22, §3.1]. If $\text{char}(k) = 0$ then any finite-dimensional representation of a semisimple Lie algebra is completely reducible [22, Thm. 6.3], and any Lie algebra \mathfrak{h} has a *Levi decomposition* $\mathfrak{h} = \mathfrak{k} \oplus \text{rad}(\mathfrak{h})$, where \mathfrak{k} is a semisimple subalgebra of \mathfrak{h} (not necessarily an ideal) [16, §6.8 Thm. 5]. In this case, if $\text{rad}(\mathfrak{h})$ is G -toral then $\text{rad}(\mathfrak{h}) = \mathfrak{z}(\mathfrak{h})$. For if $x \in \mathfrak{h}$ is semisimple then x centralises $\text{rad}(\mathfrak{h})$ by Lemma 6.9; but the set of semisimple elements of \mathfrak{k} is dense as \mathfrak{k} is semisimple, so \mathfrak{k} centralises $\text{rad}(\mathfrak{h})$, so $\text{rad}(\mathfrak{h}) \subseteq \mathfrak{z}(\mathfrak{h})$. The reverse inclusion follows because $\mathfrak{z}(\mathfrak{k}) = 0$.

Theorem 7.3. *Let \mathfrak{h} be a subalgebra of \mathfrak{g} . Consider the following conditions:*

- (i) \mathfrak{h} acts semisimply on \mathfrak{g} .
- (ii) \mathfrak{h} is G -completely reducible.
- (iii) $\text{rad}(\mathfrak{h})$ is G -toral.

If p is fabulous for G then (i) \implies (ii) \implies (iii). If $\text{char}(k) = 0$ then (i)–(iii) are equivalent.

Proof. Suppose p is fabulous for G . If (i) holds then (ii) holds by Proposition 7.2, while if (ii) holds then (iii) holds by Theorem 5.14 and Lemma 6.11. Now suppose $\text{char}(k) = 0$. To complete the proof, it's enough to show that (iii) implies (i). So suppose $\text{rad}(\mathfrak{h})$ is G -toral. Write $\mathfrak{h} = \mathfrak{k} \oplus \text{rad}(\mathfrak{h})$, where \mathfrak{k} is semisimple. Since $\text{char}(k) = 0$, \mathfrak{k} acts semisimply on \mathfrak{g} . Now $\text{rad}(\mathfrak{h})$ consists of pairwise commuting ad-semisimple elements, so $\text{rad}(\mathfrak{h})$ acts semisimply on \mathfrak{g} . But $\text{rad}(\mathfrak{h})$ commutes with \mathfrak{k} , so it follows easily that \mathfrak{h} acts semisimply on \mathfrak{g} , as required. \square

Let $f: G \rightarrow M$ be a homomorphism of connected reductive groups. We say that f is *non-degenerate* if $\ker(f)^0$ is a torus. The following is the Lie algebra counterpart of [36, Cor. 4.3], see also [7, Lem. 2.12(ii)].

Corollary 7.4. *Suppose $\text{char}(k) = 0$. Let $f: G \rightarrow M$ be a homomorphism of connected reductive groups. Let \mathfrak{h} be a subalgebra of \mathfrak{g} . If \mathfrak{h} is G -completely reducible, then $df(\mathfrak{h})$ is M -completely reducible. Conversely, if f is non-degenerate and $df(\mathfrak{h})$ is M -completely reducible then \mathfrak{h} is G -completely reducible.*

Proof. Write $\mathfrak{h} = \mathfrak{k} \oplus \mathfrak{z}$, where \mathfrak{k} is semisimple and $\mathfrak{z} = \text{rad}(\mathfrak{h})$. Then $df(\mathfrak{k})$ is semisimple and $df(\mathfrak{z})$ is solvable, so $df(\mathfrak{h}) = df(\mathfrak{k}) \oplus df(\mathfrak{z})$. It follows that $\text{rad}(df(\mathfrak{h})) = df(\mathfrak{z})$. If \mathfrak{h} is G -cr then \mathfrak{z} is G -toral by Theorem 7.3, so there is a torus S of G such that $\mathfrak{z} \subseteq \mathfrak{s}$. Then $df(\mathfrak{z})$ belongs to the Lie algebra of the torus $f(S)$, so $df(\mathfrak{z})$ is G -toral. Hence $df(\mathfrak{h})$ is M -cr by Theorem 7.3.

Conversely, suppose f is non-degenerate and $df(\mathfrak{h})$ is M -cr. Then $df(\mathfrak{z}) = \text{rad}(df(\mathfrak{h}))$ is M -toral by Theorem 7.3. Since $\ker(df) = \text{Lie}(\ker(f))$ is central in \mathfrak{g} and consists of semisimple elements, it follows that \mathfrak{z} consists of semisimple elements. Hence \mathfrak{z} is G -toral (Lemma 6.8), so \mathfrak{h} is G -cr by Theorem 7.3. \square

Remark 7.5. Applying Theorem 3.11, we can translate Corollary 7.4 into more geometric language. Let $m \in \mathbb{N}$ and let $(x_1, \dots, x_m) \in \mathfrak{g}^m$. It follows from Corollary 7.4 applied to the subalgebra \mathfrak{h} generated by the x_i that if $G \cdot (x_1, \dots, x_m)$ is closed then $M \cdot df(x_1, \dots, x_m)$ is closed, and that the converse also holds if f is non-degenerate.

Corollary 7.6. *Suppose $\text{char}(k) = 0$. Let \mathfrak{h} be a G -completely reducible subalgebra of \mathfrak{g} . Then \mathfrak{h} is Jordan-closed.*

Proof. We can write $\mathfrak{h} = \mathfrak{k} \oplus \text{rad}(\mathfrak{h})$ for some semisimple subalgebra \mathfrak{k} of \mathfrak{g} . Now $\text{rad}(\mathfrak{h})$ is G -toral by Theorem 7.3, so $\text{rad}(\mathfrak{h})$ is Jordan-closed and $\text{rad}(\mathfrak{h}) = \mathfrak{z}(\mathfrak{h})$. By [34, Lem. 3.2], \mathfrak{k} is algebraic, so \mathfrak{k} is Jordan-closed. It now follows easily that \mathfrak{h} is Jordan-closed. \square

Remark 7.7. The equivalence of (ii) and (iii) in Theorem 7.3 can fail in positive characteristic. For example, let p , G and \mathfrak{h} be as in Example 2.7. We observed earlier that \mathfrak{h} is G -ir, but it is easy to check that every element of \mathfrak{h} is nilpotent.

Conversely, let k be algebraically closed of characteristic 3, let $M = \text{SL}_2$, let V be the natural module for M , and consider the action of M on the third symmetric power $W := S^3V$ —this gives an embedding of M inside $G := \text{GL}_4$ and gives rise to a faithful representation of the Lie algebra \mathfrak{m} . We claim that W is not semisimple as an \mathfrak{m} -module, so \mathfrak{m} is not G -cr even though it has trivial radical. The four-dimensional module W has a basis x^3, x^2y, xy^2, y^3 , where x and y can be identified with the standard basis vectors for V . The basis vectors x^3 and y^3 are killed by the action of \mathfrak{m} , so span a two-dimensional submodule with a trivial \mathfrak{m} -action. The quotient by this submodule is a copy of the natural module V , which is simple, but there is no submodule of W isomorphic as an \mathfrak{m} -module to V , as direct calculation with the standard generators for $\mathfrak{m} \cong \mathfrak{sl}_2$ will easily verify.

Corollary 7.6 can also fail in positive characteristic. For, let $\text{char}(k) = 2$ and let $G = \text{PGL}_2(k) \times k^* \times k^*$ and let \mathfrak{h} be the subalgebra of \mathfrak{g} spanned by the elements of the form $\left(\left(\begin{smallmatrix} 0 & a \\ 0 & 0 \end{smallmatrix} \right), a, 0 \right)$ and $\left(\left(\begin{smallmatrix} 0 & 0 \\ b & 0 \end{smallmatrix} \right), 0, b \right)$ for $a, b \in k$. Clearly \mathfrak{h} is G -cr but is not Jordan-closed.

Remark 7.8. Richardson’s seminal paper [34] laid the foundations for the study via GIT of G -complete reducibility for subgroups and subalgebras. We explain how to obtain Theorem 7.3 in the characteristic 0 case from Richardson’s results. Suppose $\text{char}(k) = 0$. If \mathfrak{h} is a subalgebra of \mathfrak{g} then there is a unique smallest subalgebra $\mathfrak{h}^{\text{alg}}$ of \mathfrak{g} such that $\mathfrak{h} \subseteq \mathfrak{h}^{\text{alg}}$ and $\mathfrak{h}^{\text{alg}}$ is algebraic, and there is a unique connected subgroup $A(\mathfrak{h})$ of G such that $\mathfrak{h}^{\text{alg}} = \text{Lie}(A(\mathfrak{h}))$. In fact, the map $K \mapsto \text{Lie}(K)$ gives an inclusion-preserving bijection from the set of connected subgroups of G to the set of algebraic subalgebras of \mathfrak{g} [23, 13.1 Thm.].

Now let $m \in \mathbb{N}$, let $x_1, \dots, x_m \in \mathfrak{g}$, let \mathfrak{h} be the subalgebra of \mathfrak{g} generated by the x_i and let $M = A(\mathfrak{h})$. Richardson showed that the following are equivalent (see [34, Lem. 3.5 and Thm. 3.6]): (a) $G \cdot (x_1, \dots, x_m)$ is closed; (b) M is reductive; (c) M acts semisimply on \mathfrak{g} . These conditions are equivalent to \mathfrak{h} being G -cr, by Theorem 3.11. Note that if M is reductive then $\text{rad}(\mathfrak{h}^{\text{alg}}) = \text{rad}(\text{Lie}(M)) = \text{Lie}(Z(M)^0)$ is G -toral, and it is not hard to see that the converse holds. Moreover, it is also straightforward to show that $\text{rad}(\mathfrak{h}^{\text{alg}}) = \text{rad}(\mathfrak{h})^{\text{alg}}$, and that $\text{rad}(\mathfrak{h})^{\text{alg}}$ is G -toral if and only if $\text{rad}(\mathfrak{h})$ is G -toral. The characteristic 0 case of Theorem 7.3 now follows.

Example 7.9. Assume $\text{char}(k) = 0$. We now give an explicit description of the k -semisimplification of a subalgebra \mathfrak{h} of \mathfrak{g} . Write $\mathfrak{h} = \mathfrak{k} \oplus \mathfrak{m}$, where \mathfrak{k} is semisimple and $\mathfrak{m} = \text{rad}(\mathfrak{h})$. Then \mathfrak{k} is Jordan-closed (see the proof of Corollary 7.6), and it follows easily that $\mathfrak{h}^J = \mathfrak{k} \oplus \mathfrak{m}^J$. Now \mathfrak{m}^J is solvable by Proposition 6.12. Write $\mathfrak{m}^J = \mathfrak{s} \oplus \mathfrak{n}$ as in Corollary 6.13. We claim that $\mathfrak{k} \oplus \mathfrak{s}$ is a semisimplification of both \mathfrak{h} and \mathfrak{h}^J .

To see this, choose a parabolic subgroup P of G such that P yields a semisimplification of \mathfrak{h}^J . Since $\mathfrak{k} \oplus \mathfrak{s}$ is G -cr by Theorem 7.3, we can choose $\lambda \in Y(G)$ such that $P_\lambda = P$ and λ centralises $\mathfrak{k} \oplus \mathfrak{s}$. Now \mathfrak{m}^J is an ideal of \mathfrak{h}^J , so (P_λ, L_λ) also yields a semisimplification of \mathfrak{m}^J by Theorem 5.14(b). Hence c_λ kills \mathfrak{n} by Corollary 6.13(a). It follows that $c_\lambda(\mathfrak{h}^J) = \mathfrak{k} \oplus \mathfrak{s}$.

If $a \in k^*$ then $\lambda(a) \cdot \mathfrak{m} \subseteq \mathfrak{p}_\lambda$ and $c_\lambda(\lambda(a) \cdot \mathfrak{m}) = c_\lambda(\mathfrak{m})$ since λ centralises \mathfrak{s} . Hence $c_\lambda(\mathfrak{m}) = c_\lambda(\mathfrak{m}')$, where \mathfrak{m}' is the subspace of \mathfrak{p}_λ spanned by the subspaces $\lambda(a) \cdot \mathfrak{m}$ for $a \in k^*$. It is clear that \mathfrak{m}' contains \mathfrak{m}^J , since \mathfrak{n} is contained in the sum of the nonzero weight spaces for λ acting on \mathfrak{p}_λ . Hence $c_\lambda(\mathfrak{m}) = c_\lambda(\mathfrak{m}') = c_\lambda(\mathfrak{m}^J) = \mathfrak{s}$, and we conclude that $c_\lambda(\mathfrak{h}) = \mathfrak{k} \oplus \mathfrak{s}$. But $\mathfrak{k} \oplus \mathfrak{s}$ is G -cr, so the rest of the claim follows.

We finish by giving a refinement of Remark 4.4 in characteristic 0.

Proposition 7.10. *Let k, H, G, ψ and f_1, \dots, f_t be as in Remark 4.4. We suppose further that $\text{char}(k) = 0$ and we keep our assumption that k is algebraically closed. Let $(y_1, \dots, y_m) \in \mathfrak{g}^m$ such that the subalgebra \mathfrak{m} generated by the y_i is G -completely reducible. Then (y_1, \dots, y_m) belongs to $G \cdot d\iota(\mathfrak{h}^m)$ if and only if $f_i(y_1, \dots, y_m) = 0$ for $1 \leq i \leq t$.*

Proof. Suppose (y_1, \dots, y_m) belongs to $G \cdot d\iota(\mathfrak{h}^m)$. Then there exist $(x_1, \dots, x_m) \in \mathfrak{h}^m$ and $g \in G$ such that $g \cdot (y_1, \dots, y_m) = d\iota(x_1, \dots, x_m)$. So

$$\pi_{\mathfrak{g}^m, G}(y_1, \dots, y_m) = \pi_{\mathfrak{g}^m, G}(d\iota(x_1, \dots, x_m)) = \psi(\pi_{\mathfrak{h}^m, H}(x_1, \dots, x_m)) \in \psi(\mathfrak{h}^m // H).$$

Remark 4.4 implies that $f_i(y_1, \dots, y_m) = 0$ for $1 \leq i \leq t$.

Conversely, suppose $f_i(y_1, \dots, y_m) = 0$ for $1 \leq i \leq t$. Then

$$\pi_{\mathfrak{g}^m, G}(y_1, \dots, y_m) = \psi(\pi_{\mathfrak{h}^m, H}(x_1, \dots, x_m)) = \pi_{\mathfrak{g}^m, G}(d\iota(x_1, \dots, x_m))$$

for some $(x_1, \dots, x_m) \in \mathfrak{h}^m$ by the choice of the f_i . Without loss we can assume that the orbit $H \cdot (x_1, \dots, x_m)$ is closed. Then $G \cdot d\iota(x_1, \dots, x_m)$ is closed by Remark 7.5. Now $G \cdot (y_1, \dots, y_m)$ is closed by Theorem 3.11. It follows that $G \cdot d\iota(x_1, \dots, x_m) = G \cdot (y_1, \dots, y_m)$, so $(y_1, \dots, y_m) \in G \cdot d\iota(\mathfrak{h}^m)$. \square

Proposition 7.10 will be used in forthcoming work of the third author with A.R. Gover.

Acknowledgments: The research of this work was supported in part by the DFG (Grant #RO 1072/22-1 (project number: 498503969) to G. Röhrle).

REFERENCES

- [1] P. Abramenko, K. Brown, *Buildings. Theory and applications*, Graduate Texts in Mathematics, **248**, Springer, New York, 2008. xxii+747 pp.
- [2] F. Bannuscher, A. Litterick, T. Uchiyama, *Complete reducibility of subgroups of reductive algebraic groups over non-perfect fields IV: an F_4 example*. J. Group Theory **25** (2022), no. 3, 527–541.
- [3] P. Bardsley, R.W. Richardson, *Étale slices for algebraic transformation groups in characteristic p* , Proc. London Math. Soc. (3) **51** (1985), no. 2, 295–317.
- [4] M. Bate, H. Geranios, B. Martin, *Orbit closures and invariants*, Math. Z. **293** (2019), no. 3-4, 1121–1159.
- [5] M. Bate, S. Herpel, B. Martin, G. Röhrle, *Cocharacter-closure and spherical buildings*, Pacific J. Math. **279** (2015), no. 1–2, 65–85.
- [6] ———, *Cocharacter-closure and the rational Hilbert-Mumford theorem*, Math. Z. **287** (2017), no. 1–2, 39–72.
- [7] M. Bate, B. Martin, G. Röhrle, *A geometric approach to complete reducibility*, Invent. Math. **161**, no. 1 (2005), 177–218.

- [8] ———, *Complete reducibility and separable field extensions*, C. R. Acad. Sci. Paris Ser. I Math. **348** (2010), 495–497.
- [9] ———, *The strong Centre Conjecture: an invariant theory approach*, J. Algebra **372** (2012), 505–530.
- [10] ———, *Semisimplification for subgroups of reductive algebraic groups*, Forum Math. Sigma **8** (2020), Paper No. e43, 10 pp.
- [11] M. Bate, B. Martin, G. Röhrle, R. Tange, *Complete reducibility and separability*, Trans. Amer. Math. Soc. **362** (2010), no. 8, 4283–4311.
- [12] ———, *Complete reducibility and conjugacy classes of tuples in algebraic groups and Lie algebras*, Math. Z., **269** (2011), no. 3-4, 809–832.
- [13] ———, *Closed orbits and uniform S -instability in geometric invariant theory*, Trans. Amer. Math. Soc. **365** (2013), no. 7, 3643–3673.
- [14] M. Bate, D.I. Stewart, *Irreducible modules for pseudo-reductive groups*, J. Eur. Math. Soc. (JEMS) **24** (2022), no. 7, 2533–2553.
- [15] A. Borel, *Linear algebraic groups*, Graduate Texts in Mathematics, **126**, Springer-Verlag 1991.
- [16] N. Bourbaki, *Lie groups and Lie algebras*. Chapter 1, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 2007
- [17] ———, *Lie groups and Lie algebras*. Chapters 7–9, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 2005
- [18] B. Conrad, O. Gabber, G. Prasad, *Pseudo-reductive groups*, second ed., New Mathematical Monographs, vol. 26, Cambridge University Press, Cambridge, 2015.
- [19] M. Demazure, P. Gabriel. *Groupes algébriques. Tome I: Géométrie algébrique, généralités, groupes commutatifs*. Éditeurs, Paris; North-Holland Publishing Co., Amsterdam, 1970.
- [20] S. Herpel, *On the smoothness of centralizers in reductive groups*. Trans. Amer. Math. Soc. **365** (2013), no. 7, 3753–3774.
- [21] S. Herpel, D.I. Stewart, *On the smoothness of normalisers, the subalgebra structure of modular Lie algebras, and the cohomology of small representations*. Doc. Math. **21** (2016), 1–37.
- [22] J.E. Humphreys, *Introduction to Lie algebras and representation theory*, Graduate Texts in Math. Vol. 9, Springer-Verlag, New York, 1972.
- [23] ———, *Linear Algebraic Groups*, Springer-Verlag, New York, 1975.
- [24] J.C. Jantzen, *Nilpotent orbits in representation theory. Lie theory*, Progr. Math., 228, Birkhäuser Boston, Boston, MA, 2004.
- [25] G.R. Kempf, *Instability in invariant theory*, Ann. Math. **108** (1978), 299–316.
- [26] B. Leeb, C. Ramos-Cuevas, *The center conjecture for spherical buildings of types F_4 and E_6* , Geom. Funct. Anal. **21** (2011), no. 3, 525–559.
- [27] P. Levy, G. McNinch, D. Testerman, *Nilpotent subalgebras of semisimple Lie algebras*. C. R. Math. Acad. Sci. Paris **347** (2009), no. 9-10, 477–482.
- [28] B. Martin, *Reductive subgroups of reductive groups in nonzero characteristic*, J. Algebra **262**, (2003), no. 2, 265–286.
- [29] G. McNinch, *Completely reducible Lie subalgebras*, Transform. Groups **12** (2007), no. 1, 127–135.
- [30] J.S. Milne, *Algebraic groups. The theory of group schemes of finite type over a field*. Cambridge Studies in Advanced Mathematics, **170**. Cambridge University Press, Cambridge, 2017.
- [31] B. Mühlherr, J. Tits, *The Centre Conjecture for non-exceptional buildings*, J. Algebra **300** (2006), no. 2, 687–706.
- [32] P. E. Newstead, *Introduction to moduli problems and orbit spaces*, Tata Institute of Fundamental Research Lectures on Mathematics and Physics **51**. Tata Institute of Fundamental Research, Bombay, 1978.
- [33] C. Ramos-Cuevas, *The center conjecture for thick spherical buildings*, Geom. Dedicata **166** (2013), 349–407.
- [34] R.W. Richardson, *Conjugacy classes of n -tuples in Lie algebras and algebraic groups*, Duke Math. J. **57** (1988), no. 1, 1–35.
- [35] J-P. Serre, *La notion de complète réductibilité dans les immeubles sphériques et les groupes réductifs*, Séminaire au Collège de France, résumé dans [40, pp. 93–98.] (1997).
- [36] ———, *Complète réductibilité*, Séminaire Bourbaki, 56ème année, 2003–2004, n° 932.

- [37] T.A. Springer, *Linear algebraic groups*, Second edition. Progress in Mathematics, 9. Birkhäuser Boston, Inc., Boston, MA, 1998.
- [38] J. Tits, *Buildings of spherical type and finite BN-pairs*, Lecture Notes in Math. **386**, Springer-Verlag (1974).
- [39] ———, *Unipotent elements and parabolic subgroups of reductive groups. II*. Algebraic groups Utrecht 1986, 265–284, Lecture Notes in Math., 1271, Springer, Berlin, 1987.
- [40] ———, *Résumés des Cours au Collège de France, (1973–2000)*, Documents Mathématiques **12**, Soc. Math. France (2013).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, YORK YO10 5DD, UNITED KINGDOM
Email address: michael.bate@york.ac.uk

FAKULTÄT FÜR MATHEMATIK, RUHR-UNIVERSITÄT BOCHUM, D-44780 BOCHUM, GERMANY
Email address: soeren.boehm@rub.de

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ABERDEEN, KING'S COLLEGE, FRASER NOBLE BUILDING, ABERDEEN AB24 3UE, UNITED KINGDOM
Email address: b.martin@abdn.ac.uk

FAKULTÄT FÜR MATHEMATIK, RUHR-UNIVERSITÄT BOCHUM, D-44780 BOCHUM, GERMANY
Email address: gerhard.roehrle@rub.de

FAKULTÄT FÜR MATHEMATIK, RUHR-UNIVERSITÄT BOCHUM, D-44780 BOCHUM, GERMANY
Email address: laura.voggesberger@rub.de