

**ORIGINAL RESEARCH ARTICLE**

# Balloon catheter for induction of labor in women with one previous cesarean and an unfavorable cervix

Claartje M. A. Huisman<sup>1</sup> | Mieke L. G. ten Eikelder<sup>2</sup> | Kelly Mast<sup>3</sup> |  
 Katrien Oude Rengerink<sup>4</sup> | Marta Jozwiak<sup>5</sup> | Frédérique van Dunné<sup>1</sup> |  
 Johannes J. Duvekot<sup>6</sup> | Jim van Eyck<sup>7</sup> | Ingrid Gaugler-Senden<sup>8</sup> |  
 Christianne J. M. de Groot<sup>9</sup> | Maureen T. M. Franssen<sup>10</sup> | Nicolette van Gemund<sup>11</sup> |  
 Josje Langenveld<sup>12</sup> | Jan Willem de Leeuw<sup>13</sup> | Eefje J. Oude Lohuis<sup>14,7</sup> |  
 Martijn A. Oudijk<sup>9</sup> | Dimitri Papatsonis<sup>15</sup> | Mariëlle van Pampus<sup>16</sup> |  
 Martina Porath<sup>17</sup> | Sabina Rombout-de Weerd<sup>18</sup> | Jos J. van Roosmalen<sup>2</sup> |  
 Paulien C. M. van der Salm<sup>19</sup> | Hubertina C. J. Scheepers<sup>3</sup> | Marko J. Sikkema<sup>20</sup> |  
 Jan Sporken<sup>21</sup> | Rob H. Stigter<sup>22</sup> | Wim J. van Wijngaarden<sup>1</sup> | Mallory Woiski<sup>23</sup> |  
 Ben Willem J. Mol<sup>24</sup> | Kitty W. M. Bloemenkamp<sup>25</sup> | The PROBAAT-S project group

<sup>1</sup>Obstetrics and Gynecology, Haaglanden Medical Center, The Hague, the Netherlands

<sup>2</sup>Obstetrics and Gynecology, Leiden University Medical Center, Leiden, the Netherlands

<sup>3</sup>Obstetrics and Gynecology, Academic Hospital Maastricht, Maastricht, the Netherlands

<sup>4</sup>Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, the Netherlands

<sup>5</sup>Obstetrics and Gynecology, St. Antonius Hospital, Nieuwegein, the Netherlands

<sup>6</sup>Obstetrics and Gynecology, Erasmus University Medical Center, Rotterdam, the Netherlands

<sup>7</sup>Obstetrics and Gynecology, Isala Clinics, Zwolle, the Netherlands

<sup>8</sup>Obstetrics and Gynecology, Jeroen Bosch Hospital, Den Bosch, the Netherlands

<sup>9</sup>Obstetrics and Gynecology, Amsterdam UMC, University of Amsterdam, Amsterdam, the Netherlands

<sup>10</sup>Obstetrics and Gynecology, University Medical Center Groningen, Groningen, the Netherlands

<sup>11</sup>Obstetrics and Gynecology, St. Franciscus Gasthuis, Rotterdam, the Netherlands

<sup>12</sup>Obstetrics and Gynecology, Zuyderland Medical Center, Heerlen, the Netherlands

<sup>13</sup>Obstetrics and Gynecology, Ikazia Hospital, Rotterdam, the Netherlands

<sup>14</sup>Obstetrics and Gynecology, Medical Spectrum Twente, Enschede, the Netherlands

<sup>15</sup>Obstetrics and Gynecology, Amphia Hospital, Breda, the Netherlands

<sup>16</sup>Obstetrics and Gynecology, Onze Lieve Vrouwe Gasthuis, Amsterdam, the Netherlands

<sup>17</sup>Obstetrics and Gynecology, Maxima Medical Center, Veldhoven, the Netherlands

<sup>18</sup>Obstetrics and Gynecology, Albert Schweitzer Hospital, Dordrecht, the Netherlands

<sup>19</sup>Obstetrics and Gynecology, Meander Medical Center, Amersfoort, the Netherlands

<sup>20</sup>Obstetrics and Gynecology, Hospital Group Twente (ZGT), Almelo, the Netherlands

<sup>21</sup>Obstetrics and Gynecology, Canisius Hospital, Nijmegen, the Netherlands

<sup>22</sup>Obstetrics and Gynecology, Deventer Hospital, Deventer, the Netherlands

**Abbreviations:** aOR, adjusted odds ratios; BMI, body mass index; CI, confidence interval; RR, relative risk; VBAC, vaginal birth after cesarean.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2019 The Authors. *Acta Obstetrica et Gynecologica Scandinavica* published by John Wiley & Sons Ltd on behalf of Nordic Federation of Societies of Obstetrics and Gynecology (NFOG)

<sup>23</sup>Obstetrics and Gynecology, Radboud University Medical Center, Nijmegen, the Netherlands

<sup>24</sup>Department of Obstetrics and Gynecology, Monash University, Melbourne, Australia

<sup>25</sup>Division Women and Baby, Department of Obstetrics, Birth Center Wilhelmina Children's Hospital, University Medical Center Utrecht, Utrecht, the Netherlands

#### Correspondence

Claartje Huisman, Department of Obstetrics and Gynecology, Haaglanden Medical Center, Postbus 432, 2501 CK Den Haag, the Netherlands.

Email: c.m.a.huisman@gmail.com

#### Funding information

Financial support of €25,000 was given by the Perinatology and Maternal Disease workgroup of the Dutch Society of Obstetrics and Gynecology NVOG and €25,000 by the Dutch Obstetric Consortium.

## Abstract

**Introduction:** When women with a previous cesarean section and an unfavorable cervix have an indication for delivery, the choice is to induce labor or to perform a cesarean section. This study aims to assess the effectiveness and safety of a balloon catheter as a method of induction of labor in women with one previous cesarean section and an unfavorable cervix compared with an elective repeat cesarean section.

**Material and methods:** We performed a prospective cohort study in 51 hospitals in the Netherlands on term women with one previous cesarean section, a live singleton fetus in cephalic position, an unfavorable cervix and an indication for delivery. We recorded obstetric, maternal and neonatal characteristics. We compared the outcome of women who were induced with a balloon catheter with the outcome of women who delivered by elective repeat cesarean section. Main outcomes were maternal and neonatal morbidity. Mode of delivery was a secondary outcome for women who were induced. Adjusted odds ratios (aOR) were calculated using logistic regression, adjusted for potential confounders.

**Results:** Analysis was performed on 993 women who were induced and 321 women who had a repeat cesarean section (August 2011 until September 2012). Among the women who were induced, 560 (56.4%) delivered vaginally and 11 (1.1%) sustained a uterine rupture. Composite adverse maternal outcome (uterine rupture, severe postpartum hemorrhage or postpartum infection) occurred in 73 (7.4%) in the balloon and 14 (4.5%) women in the repeat cesarean section group (aOR 1.58, 95% confidence interval [CI] 0.85-2.96). Composite adverse neonatal outcome (Apgar score <7 at 5 minutes or umbilical pH <7.10) occurred in 57 (5.7%) and 10 (3.2%) neonates, respectively (aOR 1.40, 95% CI 0.87-3.48). Women who were induced had a shorter postpartum admission time (2.0 vs 3.0 days ( $P < 0.0001$ )).

**Conclusions:** In women with a previous cesarean section and a need for delivery, induction of labor with a balloon catheter does not result in a significant increase in adverse maternal and neonatal outcomes as compared with planned cesarean section.

#### KEYWORDS

balloon catheter, cervical ripening, induction of labor, repeat cesarean, vaginal birth after cesarean

## 1 | INTRODUCTION

Among women attempting vaginal birth after a previous cesarean section, labor is induced in 18%-27%.<sup>1-3</sup> Previous studies have shown that 60%-80% of women with one previous cesarean section will deliver vaginally if a trial of labor is allowed, even when induced.<sup>3-5</sup> When delivery is indicated, a decision must be made whether to

induce labor or perform primary cesarean section, and the risks and benefits of this choice must be weighed carefully.

Various publications suggest that induction of labor in women after previous cesarean section increases the risk of uterine rupture, especially after induction with prostaglandins.<sup>3,6,7</sup> In women with a previous cesarean section, balloon catheters have also been proven effective and safe, with vaginal delivery rates of 55.7%-71% and

uterine rupture rates of 0.3%-1.6%.<sup>3,8-10</sup> Guidelines now discourage the use of prostaglandins but suggest use of the balloon catheter for cervical ripening.<sup>11-13</sup>

With rising cesarean rates worldwide, repeat cesarean section with corresponding maternal and neonatal morbidity are increasing as well.<sup>14</sup> The national cesarean section rate in the Netherlands was 16% (28 713/176 155) in 2012.<sup>15</sup> In a national study including 4569 women with a previous cesarean section, 72% (3274/4569) attempted a trial of labor.<sup>1</sup> Although prostaglandins are still used for cervical ripening, the use of mechanical methods in this population is increasing.<sup>16</sup>

When an indication for delivery arises, the decision between induction of labor or a repeat cesarean section may be a difficult one, and one that many clinicians face daily. However, so far, a comparison between the two in terms of effectiveness and safety of the method has not been reported.

The objective of this study is to assess the effectiveness and safety of the balloon catheter in women with one previous cesarean section and an unfavorable cervix compared with an elective repeat cesarean section.

## 2 | MATERIAL AND METHODS

We performed a prospective cohort study in 51 hospitals in the Netherlands. We included term, pregnant women (between 37 and 42 weeks' gestational age) with one previous cesarean section, a live singleton pregnancy in cephalic presentation, an unfavorable cervix and an indication for delivery. The cervix was considered unfavorable if amniotomy was not possible and induction by a cervical priming method such as a balloon catheter was necessary. Women with contraindications for vaginal delivery, such as placenta previa, previous classical uterine incision or obstructing cervical fibroids were not included.

We compared women undergoing an induction of labor by a balloon catheter (with subsequent amniotomy and/or oxytocin augmentation if necessary) with women who had an elective repeat cesarean section performed for one or more reasons. Because of rapid recruitment of women in the elective repeat cesarean section group, with the previous cesarean section as a single reason, recruitment in this group was stopped a few months into the study period. We considered this a homogeneous group and further recruitment was not possible due to lack of resources. Recruitment in the induction of labor group continued until the target number of 1000 women was reached. Some women were excluded because they were preterm, resulting in a cohort of 993 women induced by balloon catheter.

In women undergoing induction of labor, indications for induction of labor were hypertensive disorders, threatening post-term pregnancy (before 42 weeks), insulin-dependent diabetes, intrauterine growth restriction or oligohydramnion, suspected fetal compromise of a non-acute nature (decreased fetal movement and/or suboptimal fetal heart rate tracing) or other maternal or neonatal reasons (ie, elective, intrahepatic cholestasis of pregnancy, pelvic

### Key message

In women with a previous cesarean section and a need for delivery, induction of labor with a balloon catheter does not result in a significant increase in adverse maternal and neonatal outcomes as compared with planned cesarean section.

instability, social or psychological reasons, gestational diabetes, obstetric history or suspected macrosomia).

Insertion of the balloon catheter was performed according to local protocol in the participating hospitals. Generally, the catheter was placed transcervically either manually or using a speculum, followed by fetal heart rate tracing. A single balloon (Foley) catheter (16F or 18F,  $n = 847$ ), a double balloon (Cook) catheter ( $n = 125$ ) or a prostatectomy catheter (20F) ( $n = 21$ ) was used. After passing the internal os, the single balloon and prostatectomy catheters were most commonly filled with 30-50 mL of sterile saline and the double balloon catheters with 60-80 mL. Evaluation of cervical ripeness was done as per hospital protocol, generally after 12-24 hours. When the balloon catheter was expelled or removed, and the cervix was judged to be "ripe", amniotomy was performed and continuous fetal monitoring was started. If uterine activity was insufficient ( $<3$  contractions per 10 minutes), intravenous oxytocin was infused until three to four contractions per 10 minutes or adequate progression occurred.

Primary maternal outcome was a composite maternal morbidity that consisted of uterine rupture (defined as clinical symptoms such as abdominal pain, abnormal fetal heart rate pattern, acute loss of contractions or vaginal blood loss that led to an emergency cesarean section, at which the presumed diagnosis of uterine rupture (with complete rupture of the uterine wall and serosa) was confirmed; or peripartum hysterectomy or laparotomy for uterine rupture after vaginal birth), severe postpartum hemorrhage ( $\geq 2$  L blood loss, blood transfusion, hysterectomy or re-laparotomy for bleeding) or postpartum infection (defined as treated urinary tract infection, endometritis, pneumonia, wound infection or any other unspecified suspected maternal infection requiring treatment).<sup>17</sup> The components of composite adverse maternal outcome were also assessed separately.

Secondary outcomes included suspected maternal intrapartum infection (defined as fever of  $\geq 38^\circ\text{C}$  during labor or fetal tachycardia (a persistent fetal heart rate of more than 150 bpm) and start of broad-spectrum intravenous antibiotics for suspected infection during labor), the amount of postpartum hemorrhage (mL) and postpartum blood transfusion. We also noted the length of maternal postpartum admission, as it is common practice in the Netherlands to be discharged within 4 hours after labor in case of an uneventful delivery. We also collected data on silent uterine ruptures noted during cesarean section (defined as a complete separation of the uterine wall and serosa without clinical symptoms), uterine scar dehiscence (separation of a preexisting scar

but with an intact uterine serosa, as seen during cesarean section, with or without clinical symptoms) and uterine hyperstimulation (more than six contractions in 10 minutes over a minimal period of two periods of 10 minutes, or a contraction lasting more than 3 minutes with fetal heart rate changes). For the induction group, secondary outcomes were the use of epidural analgesia, oxytocin augmentation, mode of delivery, indications for operative delivery and time from start of induction to delivery.

Primary neonatal outcome was an adverse neonatal composite outcome of an Apgar score <7 at 5 minutes or an umbilical arterial pH <7.10. Secondary neonatal outcomes were birthweight, neonatal death, indications for and length of neonatal admissions to the ward, medium and intensive care.

Trained research nurses identified eligible women. From their charts we collected demographics, obstetric history, reasons for induction of labor or cesarean section, intrapartum and postpartum information. Neonatal data until discharge from the hospital were extracted from the corresponding neonatal files. All data were collected in a web-based case-record form using consistency checks (Oracle Clinical version 4.5.3, www.oracle.com). We reviewed all operative reports of all cases with uterine rupture or scar dehiscence to ensure the accuracy of the diagnosis and reclassified if necessary.

## 2.1 | Statistical analyses

Women were analyzed according to the treatment that was planned, meaning that women who started induction but had to have a (emergency) cesarean, were analyzed in the induction group. For maternal outcome, adjusted odds ratios (aOR) with 95% confidence intervals (CI) were calculated after correction for the potential confounding variables maternal age >35, body mass index (BMI) >30, prior vaginal delivery, (un)planned previous cesarean section, reasons for labor induction and prolonged rupture of membranes. For neonatal outcome, aORs with 95% CI were calculated after correction for the potential confounders gestational age, maternal age >35, reasons for labor induction, known fetal (congenital) disease and prolonged rupture of membranes. *P* values of less than 0.05 and confidence intervals of relative risks and aORs that do not include 1 were considered to be statistically significant. All analyses were done with SPSS version 23 (Statistical Package for Social Sciences, Chicago, IL, USA).

## 2.2 | Ethical approval

The protocol was approved by the Medical Ethical Committee of Leiden University Medical Center and the board of directors of each of the participating hospitals approved local execution of the study (reference number p11.023; 11 March 2011).

## 3 | RESULTS

From 1 August 2011 until 30 September 2012, we enrolled 1305 women in the study, of whom 993 were induced with a balloon

catheter and 312 women underwent an elective repeat cesarean section (see Figure 1, flow diagram).


Table 1 shows the baseline characteristics of the two groups. Women in the elective repeat cesarean section group were slightly older: 34.0 vs 33.2 mean years of age ( $P = 0.006$ ). Women in the balloon catheter group had a higher gestational age: 40.0 vs 39.0 weeks ( $P < 0.0001$ ). Hypertension and threatening post-term pregnancy as an indication for delivery were more common in the induction group: 21.3% vs 10.9%, (relative risk [RR] 1.90, 95% CI 1.37-2.64,  $P < 0.0001$ ) and 35.9% vs 9.6%, (RR 3.95, 95% CI 2.76-5.64,  $P < 0.0001$ ), respectively. Of the women planned for induction of labor by balloon catheter, 27 switched to another method for various reasons (Figure 1).

Composite adverse maternal outcome (uterine rupture, severe postpartum hemorrhage or suspected postpartum infection) occurred in 7.4% and 4.5% in the balloon catheter group and the repeat cesarean section group, respectively (aOR 1.58, 95% CI 0.85-2.96,  $P = 0.15$ ).

Uterine rupture occurred 11 times in the balloon catheter group and once in the elective repeat cesarean section group. The latter was a silent rupture without clinical signs. Two of the women with a uterine rupture had severe postpartum hemorrhage and one woman also had bladder injury during cesarean section. Uterine dehiscence occurred seven times in the balloon catheter group and three times in the elective repeat cesarean section group.

Table 2 shows maternal and neonatal outcomes. Although maternal temperature of  $\geq 38^{\circ}\text{C}$  during labor occurred more often in the balloon catheter group (8.4% vs 1.6%; aOR 7.00, 95% CI 2.73-17.95,  $P < 0.0001$ ), maternal suspected intrapartum infection and postpartum infection was comparable between the two groups. The higher occurrence rate of pyrexia is most likely due to the use of epidural analgesia: 15.9% (73/459) women with epidural analgesia developed pyrexia, accounting for 88% (73/83) of all women with pyrexia.

Serious adverse events in the balloon catheter group were umbilical cord prolapse ( $n = 1$ ), partial placental abruption ( $n = 1$ , several hours after removing the balloon catheter), bladder injury ( $n = 4$ , during cesarean section), uterine inversion after vaginal birth ( $n = 1$ ) and relaparotomy ( $n = 1$ ). In two women, cesarean section was performed due to abnormal blood loss after catheter placement. One woman had an unknown amount of blood loss directly after trans-cervical balloon catheter insertion and, after removal, immediate cesarean section was performed. Another woman suddenly had approximately 1000 mL of blood loss 10 hours after insertion, after which emergency cesarean section was performed. In both cases neonatal outcome was good. The woman undergoing relaparotomy suffered from severe postpartum hemorrhage after an emergency cesarean section. Total blood loss was 7.5 L, for which she received 11 units of blood and was admitted to the intensive care unit. There were no other women with massive transfusion (defined as 10 units of red cells in 24 hours), no women with hysterectomy or organ failure and no maternal deaths. Three women were admitted to the intensive care unit, all due to severe hemorrhage (7.5, 2.8 and 2.6 L, respectively). Hyperstimulation occurred in 14 women who were induced



**FIGURE 1** Flow chart. A total of 1305 women were included in the study, of whom 993 were planned for induction of labor by balloon catheter and 312 for repeat cesarean section. \*The previous cesarean section was the sole reason to opt for a repeat cesarean section

by a balloon catheter, leading to a uterine rupture once and postpartum hemorrhage twice. Neonatal outcomes were not affected. In the repeat cesarean section group, there was one maternal adverse event: a uterine inversion during cesarean section. There were no women with massive blood transfusion, hysterectomy, organ failure and no maternal deaths. One woman was admitted to a maternal intensive care unit due to spinal muscular atrophy, for which she was observed post-cesarean.

In the balloon catheter group, 21.6% of women were not admitted (adjusted  $P < 0.0001$ ) during the postpartum period (ie, discharged within 4 hours after delivery) and those admitted had a shorter median maternal postpartum admission in the hospital (2.0 days vs 3.0 days, adjusted  $P < 0.0001$ ) (Table 2). However, median total admission time was 3.0 days for both groups.

The cesarean section rate was 43.6% (433 women) in the balloon catheter group. Further delivery characteristics for women induced by balloon catheter can be seen in Table 3.

Composite adverse neonatal outcome of an Apgar score  $<7$  at 5 minutes or an umbilical arterial pH  $<7.10$  was not significantly different between the groups, it occurred in 5.7% and 3.2% (aOR 1.40, 95% CI 0.67-2.93,  $P = 0.38$ ) after induction by balloon catheter and repeat cesarean section, respectively. Specific neonatal outcomes can be seen in Table 2. There was one case of neonatal death in the balloon catheter group due to a prenatally detected congenital heart disease. Of the 11 women with a uterine scar rupture, there were

two neonates with low Apgar scores (1/2/5 and 0/3/4 after 1, 5 and 10 minutes) and an umbilical artery pH below 7.0; they were admitted to the neonatal intensive care unit. One neonate was cooled for 72 hours with good cerebral function monitoring at discharge after 8 days; the (neurological) outcome of other neonate is unknown, discharge was after 20 days.

## 4 | DISCUSSION

The results of this prospective cohort study show that in women with one previous cesarean section and an unfavorable cervix, induction of labor with a balloon catheter is a safe method to induce delivery compared with women who deliver by means of an elective repeat cesarean section. Between the two groups, there was no significant difference in maternal morbidity such as uterine scar rupture or dehiscence, postpartum hemorrhage or infection. Postpartum admission time was shorter in the induction of labor group. Neonatal morbidity with regard to umbilical cord pH and Apgar scores was similar in both groups.

In the induction of labor group, the cesarean section rate was 43.6%, which is comparable to other studies such as the systematic review and meta-analysis by Kehl et al (n = 144, cesarean section rate 43.6%)<sup>18</sup> but higher than a recent study by Kruit et al (n = 361, cesarean section rate 38%)<sup>9</sup> and the study by Jozwiak et al (n = 208,

**TABLE 1** Baseline characteristics for method of induction/delivery

	Balloon catheter (n = 993) (%)	Repeat CS (n = 312) (%)	RR (95% CI)	P value
Maternal age mean, SD	33.2 (4.5)	34.0 (4.5)	NA	0.006
<25 y	48 (4.8)	7 (2.2)	1.92 (0.95-3.86)	0.05
25-35 y	589 (59.3)	174 (55.8)	1.12 (0.92-1.36)	0.27
>35 y	356 (35.9)	131 (42.0)	0.82 (0.68-1.00)	0.05
SES				
Low	337 (33.9)	95 (30.4)	1.13 (0.92-1.40)	0.25
Middle	372 (37.5)	131 (42.0)	0.87 (0.71-1.05)	0.15
High	253 (25.5)	78 (25.0)	1.02 (0.82-1.28)	0.87
Unknown	31 (3.1)	8 (2.6)	1.17 (0.63-2.19)	0.61
Caucasian	715 (72.0)	235 (75.3)	0.96 (0.90-1.03)	0.25
BMI, kg/m <sup>2</sup> ; median (IQR)	25.9 (23.0 - 30.0) <sup>a</sup>	26.7 (23.0 -31.2) <sup>b</sup>	NA	0.22
BMI >30	216 (21.8)	78 (25.0)	0.87 (0.70-1.09)	0.23
Parity				
1	801 (80.7)	258 (82.7)		
≥2	192 (19.3)	54 (17.3)	1.11 (0.86-1.44)	0.42
Previous vaginal births before CS				
0	880 (88.6)	270 (86.5)	1.05 (0.95-1.16)	0.32
1	88 (8.9)	35 (11.2)	0.82 (0.61-1.11)	0.21
≥2	25 (2.5)	7 (2.2)	1.10 (0.57-2.12)	0.79
Previous vaginal births after CS				
0	882 (88.8)	287 (92.0)	0.92 (0.85-1.01)	0.11
1	90 (9.1)	21 (6.7)	1.29 (0.86-1.92)	0.20
≥2	21 (2.1)	4 (1.3)	1.50 (0.61-3.71)	0.35
Previous unplanned CS	608 (61.7)	215 (69.6)	0.76 (0.62-0.95)	0.01
Gestational age wk; median, IQR	40.0 (38.6-41.1)	39.0 (38.4-39.7)	NA	<0.0001 <sup>d</sup>
Indications for induction of labor				
Hypertensive disorders	212 (21.3)	34 (10.9)	1.90 (1.37-2.64)	<0.0001
Threatening post-term pregnancy	356 (35.9)	30 (9.6)	3.95 (2.76-5.64)	<0.0001
Insulin-dependent diabetes	70 (7.0)	32 (10.3)	0.74 (0.55-1.01)	0.07
Intrauterine growth restriction or oligohydramnion	81 (8.2)	17 (5.4)	1.41 (0.90-2.20)	0.11
Fetal distress	96 (9.7)	27 (8.7)	1.10 (0.78-1.56)	0.59
Other <sup>c</sup>	367 (37.0)	224 (71.8)	0.51 (0.46-0.57)	<0.0001

BC, balloon catheter; CS, cesarean section; IQR, interquartile range; NA, not applicable; RCS, repeat cesarean section; SES, socioeconomic status; RR, relative risk.

<sup>a</sup>13% missing.

<sup>b</sup>17% missing.

<sup>c</sup>Other reasons for induction: other maternal/neonatal disease not mentioned in any of the above options.

<sup>d</sup>Mann-Whitney U test.

cesarean section rate 28.8%).<sup>10</sup> All of these studies specifically looked at balloon catheter for induction of labor at term after previous cesarean section.

We found a uterine rupture rate after induction of labor of 1.1%, which is comparable to the incidence quoted in the literature of 0.3%-1.2%.<sup>9,18</sup> Notable in our study is the relative good neonatal outcome after uterine rupture. The uterine rupture rate is high when compared with rates of 0.7%-0.8% seen in spontaneous vaginal birth after cesarean section

(VBAC).<sup>9,18</sup> This percentage must be included in counseling women who may be opting for an induced trial of labor. It is unclear whether there is a selection bias in the population or it is the balloon catheter that increases the risk of uterine rupture. Since most uterine ruptures occur in the active phase of labor, it is unlikely that the balloon catheter itself, used for cervical ripening, explains this increased risk.

Oxytocin for augmentation of labor was used in 77.5% of women, in keeping with rates quoted in the literature (68.4-85.3%).<sup>9,18</sup> The

**TABLE 2** Maternal and neonatal outcome

	Balloon catheter (n = 993) (%)	Repeat CS (n = 312) (%)	adjusted OR (95% CI)	adjusted P value
<b>Maternal outcome</b>				
Composite maternal morbidity <sup>a</sup>	73 (7.4)	14 (4.5)	1.58 (0.85-2.96)	0.15
Suspected intrapartum infection <sup>b</sup>	25 (2.5)	4 (1.3)	2.04 (0.66-6.34)	0.22
Postpartum hemorrhage <sup>c</sup>	30 (3.0)	5 (1.6)	1.34 (0.49-3.68)	0.57
Postpartum infection <sup>d</sup>	37 (3.7)	8 (2.6)	1.66 (0.73-3.81)	0.23
Uterine rupture	11 (1.1)	1 (0.3) <sup>e</sup>	3.01 (0.36-25.03)	0.31
Maternal length of admission days; median (IQR)	2.0 (1.0-3.0)	3.0 (2.25-3.0)	Beta: -1.09	<0.0001
<b>Neonatal outcome</b>				
Composite neonatal morbidity	57 (5.7)	10 (3.2)	1.40 (0.67-2.93)	0.38
<b>Apgar score &lt;7</b>				
1 min	68 (6.9)	11 (3.5)	1.74 (0.87-3.48)	0.12
5 min	21 (2.1)	4 (1.3)	1.31 (0.41-4.17)	0.65
pH <7.10	40 (5.9) <sup>f</sup>	8 (3.7) <sup>g</sup>	1.08 (0.46-2.55)	0.86
Neonatal birthweight, g; mean (SD)	3515 (527)	3627 (558)	NA	<0.0001
<b>Neonatal admission</b>				
Ward	252 (25.4)	72 (23.1)	1.24 (0.89-1.73)	0.20
Medium care	85 (8.6)	26 (8.3)	1.13 (0.69-1.85)	1.13
Intensive care	26 (2.6)	2 (0.6)	6.20 (1.31-29.38)	0.02
<b>Reason for admission</b>				
Suspected infection	42 (4.2)	9 (2.9)	1.35 (0.62-2.94)	0.45
Asphyxia	10 (1.0)	2 (0.6)	1.32 (0.27-6.54)	0.73
Dysmaturity	40 (4.0)	11 (3.5)	1.08 (0.50-2.33)	0.84
Hypoglycemia	16 (1.6)	10 (3.2)	0.46 (0.19-1.10)	0.08
Glucose protocol	160 (16.1)	60 (19.2)	0.99 (0.67-1.46)	0.95
IRDS	1 (0.1)	2 (0.6)	0.05 (0.00-0.62)	0.02
Meconium aspiration	4 (0.4)	0	NA	0.99
Pneumothorax	2 (0.2)	0	NA	0.99
Apnea	6 (0.6)	0	NA	0.99
Other or unknown <sup>h</sup>	174 (17.5)	33 (10.6)	2.18 (1.39-3.42)	0.001
Neonatal length of admission (d)	2.0 (1.0-4.0)	3.0 (3.0-5.0)	NA	<0.0001

CS, cesarean section; NA, not applicable. RR, relative risk.

<sup>a</sup>Suspected postpartum infection or severe postpartum hemorrhage or uterine rupture.

<sup>b</sup>Body temperature during labor  $\geq 38^{\circ}\text{C}$  or fetal tachycardia (a persistent fetal heart rate of  $>150$  bpm) and start of broad-spectrum antibiotics due to suspected infection.

<sup>c</sup> $>2000$  mL or blood transfusion.

<sup>d</sup>Defined as treated urinary tract infection, endometritis, pneumonia, wound infection or other unspecified suspected maternal infection.

<sup>e</sup>Silent rupture.

<sup>f</sup>31% missing values.

<sup>g</sup>34% missing values.

<sup>h</sup>Other reasons including: lung disease, pneumothorax, apnea, intraventricular hemorrhage, periventricular leucomalacia.

rate of intra- and postpartum infection after induction of labor of 2.5% and 3.7%, respectively, is comparable to that found by Kruit et al (2.8% and 2.2%, respectively).

Of all cesarean sections after induction of labor, 50.8% were because of failure to progress in the first stage. Unfortunately, we were not able to record cervical dilation at the time of cesarean section.

It is possible that many women in whom failure to progress in first stage was diagnosed, had not yet entered the accelerative phase, as commonly used standards to evaluate adequate progress of labor may not apply in women who are induced.<sup>19,20</sup>

The strength of our study is that it is the largest cohort of inductions by balloon catheter to date, including almost 1000 women.

**TABLE 3** Delivery characteristics for women induced by balloon catheter

	n = 993 (%)
<b>Intrapartum information</b>	
Epidural analgesia	457 (46.2)
Oxytocin augmentation (Y/N)	770 (77.5)
Hyperstimulation	14 (1.4)
Time from start of induction to birth (median, IQR)	30.8 (21.9-39.0)
<b>Mode of delivery</b>	
Spontaneous	469 (47.2)
Vaginal instrumental	91 (9.2)
Cesarean delivery	433 (43.6)
<b>Indication for cesarean delivery</b>	
Failure to progress in first stage	221 (50.8)
Failure to progress in second stage	24 (5.5)
Fetal distress	127(29.2)
Maternal reason	13 (3.0)
Other or Unknown	50 (11.5)
<b>Indication for vaginal instrumental delivery</b>	
Failure to progress in second stage	38 (38.8)
Fetal distress	44 (44.9)
Failure to progress in second stage AND fetal distress	13 (13.3)
Maternal complication	3 (3.1)
Operative deliveries for fetal distress <sup>a</sup>	182 (18.3)

<sup>a</sup>Two women had a cesarean delivery for fetal distress after failed ventouse extraction.

By collecting information prospectively, and not relying on ICD 10 codes, all outcomes were studied specifically.

Limitations are the short follow-up period (until discharge), so long-term effects of possible consecutive cesarean sections were not included. Another weakness is the lack of randomization. Confounding by indication—which occurs when the clinical indication for selecting a particular treatment also affects the outcome—probably meant that the groups were not comparable at baseline for important prognostic factors. Although we have corrected for numerous possible confounders, there may still be some residual confounders that we could not correct for, such as hospital induction protocols and expected fetal weight. No randomized controlled trials, however, have been performed which compare induction of labor and planned repeat cesarean sections in women with a prior cesarean section.<sup>21</sup> We considered randomization also to be unfeasible, after a randomized controlled trial on this subject in another country was stopped due to low participation rates. A large cohort study is then the best alternative.<sup>22</sup>

While cerebral palsy is thought to occur more often at an arterial umbilical cord pH <7.00, we chose to use the cut-off of pH <7.10 for the adverse neonatal composite outcome. If we had chosen for a pH

value of <7.00 or 7.05, it would mean that the incidence of adverse neonatal composite outcome would be even lower.

Another limitation is the difference in group size. The balloon catheter group is more than three times as large as the cesarean section group, partly because recruitment of women undergoing a repeat cesarean with the previous cesarean section as a single reason, was stopped after a few months.

With an unknown denominator of potentially eligible women meeting the inclusion criteria for the study in either the induction group or the repeat cesarean group, there is risk of selection bias or risk identification. For example, despite the fact that the median gestational age in the cesarean section group was shorter, mean neonatal birthweight was higher. This could possibly be the result of risk identification, where women carrying large-for-gestational-age babies were not offered induction of labor.

We consider our findings to have value for low- and middle-income settings. Although a more expectant management instead of induction may be preferred, regarding the higher rate of uterine rupture after induction compared with spontaneous vaginal birth after cesarean section, induction should still be considered if adequate fetal and maternal monitoring can be provided and emergency cesarean sections can be performed. If there is a medical indication for delivery, and there is enough time to start induction, this group of women will profit from the opportunity to have a vaginal birth instead of another cesarean, with high risks of abnormal invasive placentation in following pregnancies. Although not focused on in this manuscript, induction with balloon catheter compared with prostaglandins is cheaper and is associated with fewer abnormalities of contraction pattern, and fewer maternal side effects, and is thus well suited in resource-poor settings.

With this large cohort study, more evidence is provided that induction of labor after cesarean section is an effective and safe option when an indication for delivery arises. Labor should be closely monitored for signs of uterine rupture, and facilities for emergency cesarean section are imperative. The rates of uterine rupture, emergency cesarean section and complications are useful for clinicians and pregnant women in the shared decision-making process when facing this dilemma. Individual previous experiences, possible long-term effects and future family planning have to be discussed. Individualized management in women with a previous cesarean section and an indication for delivery need to be considered. Both women's preferences and the a priori chance of vaginal delivery in case of induction can be helpful. Decision aids combined with a prediction model that includes induction as a variable can be considered.<sup>23,24</sup>

## 5 | CONCLUSION

In women with a previous cesarean section and a need for delivery, induction of labor with a balloon catheter does not result in a significant increase in adverse maternal and neonatal outcome as compared with planned cesarean section.



## CONFLICT OF INTEREST

B.W.M. is supported by a NHMRC Practitioner Fellowship (GNT1082548). B.W.M. is a consultant for ObsEva, Merck and Guerbet.

## ORCID

Claartje M. A. Huisman  <https://orcid.org/0000-0003-0616-2847>

Jan Willem Leeuw  <https://orcid.org/0000-0001-5028-8055>

Mariëlle Pampus  <https://orcid.org/0000-0002-3020-8908>

Jos J. van Roosmalen  <https://orcid.org/0000-0002-6746-3259>

Kitty W. M. Bloemenkamp  <https://orcid.org/0000-0002-1377-4625>

## REFERENCES

1. Kwee A, Bots ML, Visser GH, Bruinse HW. Obstetric management and outcome of pregnancy in women with a history of caesarean section in the Netherlands. *Eur J Obstet Gynecol Reprod Biol.* 2007;132:171-176.
2. Locatelli A, Regalia AL, Ghidini A, Ciriello E, Biffi A, Pezzullo JC. Risks of induction of labour in women with a uterine scar from previous low transverse caesarean section. *BJOG.* 2004;111:1394-1399.
3. Ravasia DJ, Wood SL, Pollard JK. Uterine rupture during induced trial of labor among women with previous cesarean delivery. *Am J Obstet Gynecol.* 2000;183:1176-1179.
4. Durnwald C, Rouse DJ, Leveno KJ, et al. The maternal-fetal medicine units cesarean registry: safety and efficacy of a trial of labor in preterm pregnancy after a prior cesarean delivery. *Am J Obstet Gynecol.* 2006;195:1119-1126.
5. Dekker G, Chan A, Luke CG, et al. Risk of uterine rupture in Australian women attempting vaginal birth after one prior caesarean section: a retrospective population-based cohort study. *BJOG.* 2010;117:1358-1365.
6. Kayani SI, Alfirevic Z. Induction of labour with previous caesarean delivery: where do we stand? *Curr Opin Obstet Gynecol.* 2006;18:636-641.
7. Ophir E, Odeh M, Hirsch Y, Bornstein J. Uterine rupture during trial of labor: controversy of induction's methods. *Obstet Gynecol Surv.* 2012;67:734-745.
8. Bujold E, Blackwell SC, Gauthier RJ. Cervical ripening with transcervical foley catheter and the risk of uterine rupture. *Obstet Gynecol.* 2004;103:18-23.
9. Kruit H, Wilkman H, Tekay A, Rahkonen L. Induction of labor by Foley catheter compared with spontaneous onset of labor after previous cesarean section: a cohort study. *J Perinatol.* 2017;37:787-792.
10. Jozwiak M, van de Lest HA, Burger NB, Dijksterhuis MG, De Leeuw JW. Cervical ripening with Foley catheter for induction of labor after cesarean section: a cohort study. *Acta Obstet Gynecol Scand.* 2014;93:296-301.
11. American College of Obstetricians and Gynecologists. ACOG Practice Bulletin no. 115: Vaginal birth after previous cesarean delivery. *Obstet Gynecol.* 2010;116:450-463.
12. Birth after Previous Caesarean Birth (Green-top Guideline No. 45). 2015. <https://www.rcog.org.uk/en/guidelines-research-services/guidelines/gtg45/>
13. Zwangerschap en Bevalling na een Voorgaande Sectio Caesarea. 2010. <https://www.nvog.nl/wp-content/uploads/2017/12/Zwangerschap-en-bevalling-na-een-voorgaande-sectio-caesarea-1.0-04-06-2010.pdf>. Accessed 7 May 2018.
14. Betran AP, Ye J, Moller AB, Zhang J, Gulmezoglu AM, Torloni MR. The Increasing Trend in Caesarean Section Rates: Global, Regional and National Estimates: 1990-2014. *PLoS ONE.* 2016;11:e0148343.
15. The Netherlands Perinatal Registry. *Trends 1999-2012.* Utrecht: The Netherlands Perinatal Registry; 2013.
16. Huisman CM, Jozwiak M, de Leeuw JW, Mol BW, Bloemenkamp KW. Cervical ripening in the Netherlands: a survey. *Obstet Gynecol Int.* 2013;2013:745159.
17. Zwart JJ, Richters JM, Ory F, de Vries JI, Bloemenkamp KW, van Roosmalen J. Uterine rupture in The Netherlands: a nationwide population-based cohort study. *BJOG.* 2009;116:1069-1078; discussion 78-80.
18. Kehl S, Weiss C, Rath W. Balloon catheters for induction of labor at term after previous cesarean section: a systematic review. *Eur J Obstet Gynecol Reprod Biol.* 2016;204:44-50.
19. Harper LM, Caughey AB, Odibo AO, Roehl KA, Zhao Q, Cahill AG. Normal progress of induced labor. *Obstet Gynecol.* 2012;119:1113-1118.
20. Rinehart BK, Terrone DA, Hudson C, Isler CM, Larmon JE, Perry KG Jr. Lack of utility of standard labor curves in the prediction of progression during labor induction. *Am J Obstet Gynecol.* 2000;182:1520-1526.
21. Dodd JM, Crowther CA, Grivell RM, Deussen AR. Elective repeat caesarean section versus induction of labour for women with a previous caesarean birth. *Cochrane Database Syst Rev.* 2017;(7):CD004906.
22. Crowther CA, Dodd JM, Hiller JE, Haslam RR, Robinson JS, Birth After Caesarean Study Group. Planned vaginal birth or elective repeat caesarean: patient preference restricted cohort with nested randomised trial. *PLoS Med.* 2012;9:e1001192.
23. Schoorel EN, van Kuijk SM, Melman S, et al. Vaginal birth after a caesarean section: the development of a Western European population-based prediction model for deliveries at term. *BJOG.* 2014;121:194-201; discussion.
24. Schoorel EN, Vankan E, Scheepers HC, et al. Involving women in personalised decision-making on mode of delivery after caesarean section: the development and pilot testing of a patient decision aid. *BJOG.* 2014;121:202-209.

**How to cite this article:** Huisman CMA, ten Eikelder MLG, Mast K, et al. The PROBAAT-S project group. Balloon catheter for induction of labor in women with one previous cesarean and an unfavorable cervix. *Acta Obstet Gynecol Scand.* 2019;98:920-928. <https://doi.org/10.1111/aogs.13558>